



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 04:05 AM UTC

PDB ID : 3CO9 / pdb_00003co9
Title : Crystal structure of HCV NS5B polymerase with a novel pyridazinone inhibitor
Authors : Han, Q.; Showalter, R.E.; Zhao, Q.; Kissinger, C.R.
Deposited on : 2008-03-27
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	9% 86% 10% ..
1	B	578	10% 89% 8% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9397 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	562	Total	C	N	O	S	0	0	0
			4367	2755	770	811	31			
1	B	562	Total	C	N	O	S	0	0	0
			4367	2755	770	811	31			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	544	GLN	ARG	engineered mutation	UNP P26663
A	571	LEU	-	expression tag	UNP P26663
A	572	GLU	-	expression tag	UNP P26663
A	573	HIS	-	expression tag	UNP P26663
A	574	HIS	-	expression tag	UNP P26663
A	575	HIS	-	expression tag	UNP P26663
A	576	HIS	-	expression tag	UNP P26663
A	577	HIS	-	expression tag	UNP P26663
A	578	HIS	-	expression tag	UNP P26663
B	544	GLN	ARG	engineered mutation	UNP P26663
B	571	LEU	-	expression tag	UNP P26663
B	572	GLU	-	expression tag	UNP P26663
B	573	HIS	-	expression tag	UNP P26663
B	574	HIS	-	expression tag	UNP P26663
B	575	HIS	-	expression tag	UNP P26663
B	576	HIS	-	expression tag	UNP P26663
B	577	HIS	-	expression tag	UNP P26663
B	578	HIS	-	expression tag	UNP P26663

- Molecule 2 is N-{3-[4-hydroxy-1-(3-methylbutyl)-2-oxo-1,2-dihydropyrrolo[1,2-b]pyridazin-3-yl]-1,1-dioxido-2H-1,2,4-benzothiadiazin-7-yl}methanesulfonamide (CCD ID: 3MS) (formula: C₂₀H₂₃N₅O₆S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			33	20	5	6	2		
2	B	1	Total	C	N	O	S	0	0
			33	20	5	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	309	Total	O	0	0
			309	309		
3	B	288	Total	O	0	0
			288	288		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.70Å 106.42Å 126.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.19 – 2.10 39.19 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.19-2.10) 99.1 (39.19-2.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.218 , 0.266 0.221 , 0.266	Depositor DCC
R_{free} test set	3353 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9397	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.93 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5501e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3MS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	0/4463	0.82	0/6057
1	B	0.60	0/4463	0.82	3/6057 (0.0%)
All	All	0.59	0/8926	0.82	3/12114 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	552	VAL	N-CA-C	-6.46	106.52	111.62
1	B	148	GLN	CA-C-N	5.56	126.80	119.84
1	B	148	GLN	C-N-CA	5.56	126.80	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4367	0	4387	36	0
1	B	4367	0	4387	24	0
2	A	33	0	23	1	0
2	B	33	0	23	1	0
3	A	309	0	0	4	0
3	B	288	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9397	0	8820	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:303:CYS:SG	1:B:313:MET:HE1	2.02	0.99
1:B:95:HIS:H	1:B:95:HIS:CD2	2.13	0.66
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.79	0.65
1:A:43:ARG:HH11	1:A:43:ARG:HB3	1.64	0.62
1:B:445:CYS:HB2	3:B:665:HOH:O	2.00	0.61
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.83	0.60
1:A:209:LYS:HE3	3:A:852:HOH:O	2.01	0.59
1:A:313:MET:HG2	1:A:320:LEU:HD11	1.87	0.56
1:A:488:CYS:HB2	3:A:866:HOH:O	2.05	0.56
1:A:205:VAL:HG12	1:A:209:LYS:HE2	1.87	0.56
1:B:309:GLN:HB3	1:B:325:GLU:HB2	1.87	0.56
1:B:505:ARG:HH22	1:B:531:LYS:HG3	1.70	0.56
1:B:303:CYS:HB3	1:B:311:CYS:SG	2.46	0.55
1:B:22:PRO:HG2	1:B:401:ARG:HG3	1.89	0.54
1:A:118:HIS:O	1:A:121:SER:HB3	2.08	0.53
1:A:95:HIS:H	1:A:95:HIS:CD2	2.26	0.53
1:B:236:GLU:CD	1:B:280:ARG:HH22	2.16	0.52
1:A:43:ARG:HH11	1:A:43:ARG:CB	2.24	0.51
1:A:508:ARG:HH22	1:A:534:LEU:CD1	2.24	0.50
1:B:200:ARG:HH12	1:B:316:ASN:HD21	1.59	0.50
1:A:22:PRO:HG2	1:A:401:ARG:HG3	1.93	0.49
1:A:43:ARG:HH11	1:A:43:ARG:CG	2.26	0.49
1:B:95:HIS:CD2	1:B:95:HIS:N	2.80	0.48
1:B:12:THR:HG21	1:B:270:LYS:HE2	1.96	0.48
1:B:268:ASN:HB3	1:B:274:CYS:SG	2.53	0.48
1:B:200:ARG:HH12	1:B:316:ASN:ND2	2.12	0.48
1:B:213:ASN:ND2	3:B:835:HOH:O	2.46	0.48
1:A:257:THR:HA	1:A:261:TYR:HB2	1.96	0.47
1:A:144:VAL:HB	1:A:394:ARG:HG2	1.97	0.47
1:A:445:CYS:HB2	3:A:811:HOH:O	2.15	0.47
1:B:248:GLU:HB3	3:B:749:HOH:O	2.14	0.47
1:B:359:ASP:HB3	1:B:362:LEU:HD12	1.97	0.47
1:A:237:GLU:OE2	1:A:253:ILE:HG22	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:HIS:O	1:A:122:VAL:HG23	2.16	0.46
1:A:90:LYS:HA	1:A:90:LYS:HD2	1.62	0.45
2:B:579:3MS:O26	2:B:579:3MS:N21	2.47	0.45
1:A:303:CYS:HB3	1:A:311:CYS:SG	2.56	0.45
1:A:485:VAL:O	1:A:489:LEU:HG	2.17	0.45
1:B:429:PHE:O	1:B:433:LEU:HG	2.17	0.45
1:A:200:ARG:HH12	1:A:316:ASN:ND2	2.14	0.45
1:A:309:GLN:HB3	1:A:325:GLU:HB2	2.00	0.44
1:A:303:CYS:CB	1:A:311:CYS:HG	2.31	0.43
1:A:314:LEU:HB3	1:A:321:VAL:CG1	2.48	0.43
1:A:200:ARG:HH12	1:A:316:ASN:HD21	1.67	0.43
1:A:398:GLU:OE1	1:A:408:TRP:HD1	2.02	0.43
1:A:510:ARG:NH1	3:A:721:HOH:O	2.46	0.43
1:A:26:LEU:HD13	1:A:432:ILE:HG12	2.02	0.42
1:A:101:PHE:O	1:A:114:LYS:HE3	2.20	0.42
1:A:187:MET:HE2	1:A:293:LEU:HD23	2.02	0.41
1:B:182:LEU:HD12	1:B:243:CYS:SG	2.60	0.41
1:B:300:SER:HA	1:B:313:MET:HE3	2.01	0.41
1:A:160:ILE:HA	1:A:282:SER:OG	2.21	0.41
1:B:257:THR:HA	1:B:261:TYR:HB2	2.02	0.41
1:A:466:LEU:HD22	1:A:551:PHE:HE2	1.86	0.41
2:A:579:3MS:N21	2:A:579:3MS:O26	2.51	0.41
1:B:144:VAL:HB	1:B:394:ARG:HG2	2.03	0.41
1:A:346:TYR:O	1:A:347:SER:HB3	2.21	0.41
1:B:182:LEU:N	1:B:183:PRO:CD	2.84	0.41
1:A:313:MET:CG	1:A:322:VAL:HG22	2.51	0.41
1:B:1:SER:N	3:B:675:HOH:O	2.52	0.41
1:A:405:VAL:HG21	1:A:446:GLN:HG3	2.03	0.40
1:A:79:LYS:HA	1:A:244:ASP:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	560/578 (97%)	547 (98%)	12 (2%)	1 (0%)	43	44
1	B	560/578 (97%)	546 (98%)	14 (2%)	0	100	100
All	All	1120/1156 (97%)	1093 (98%)	26 (2%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	153	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	478/493 (97%)	465 (97%)	13 (3%)	39	45
1	B	478/493 (97%)	470 (98%)	8 (2%)	53	62
All	All	956/986 (97%)	935 (98%)	21 (2%)	45	53

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	CYS
1	A	43	ARG
1	A	50	LYS
1	A	90	LYS
1	A	114	LYS
1	A	151	LYS
1	A	182	LEU
1	A	313	MET
1	A	352	ASP
1	A	381	VAL
1	A	473	SER
1	A	533	LYS
1	A	536	LEU
1	B	90	LYS
1	B	150	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	160	ILE
1	B	352	ASP
1	B	440	GLU
1	B	466	LEU
1	B	532	THR
1	B	545	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	142	ASN
1	A	316	ASN
1	A	475	HIS
1	A	544	GLN
1	B	117	ASN
1	B	316	ASN
1	B	374	HIS
1	B	436	GLN
1	B	475	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3MS	A	579	-	34,36,36	1.81	3 (8%)	42,56,56	2.10	11 (26%)
2	3MS	B	579	-	34,36,36	1.90	3 (8%)	42,56,56	1.83	12 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3MS	A	579	-	-	3/10/29/29	0/3/4/4
2	3MS	B	579	-	-	4/10/29/29	0/3/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	579	3MS	S33-N25	7.02	1.72	1.63
2	B	579	3MS	S32-N23	6.61	1.69	1.62
2	A	579	3MS	S32-N23	6.29	1.68	1.62
2	A	579	3MS	S33-N25	6.19	1.71	1.63
2	B	579	3MS	C12-C14	2.52	1.50	1.44
2	A	579	3MS	C12-C14	2.37	1.50	1.44

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	579	3MS	C17-S33-N25	-6.04	99.77	106.56
2	A	579	3MS	O28-S32-C9	-4.68	103.46	109.30
2	B	579	3MS	C11-C12-C13	4.38	123.90	118.55
2	A	579	3MS	C14-N24-N22	-4.20	119.14	122.54
2	A	579	3MS	O26-C14-N24	4.04	124.09	120.52
2	B	579	3MS	C14-N24-N22	-4.00	119.30	122.54
2	A	579	3MS	O28-S32-O27	3.91	122.41	118.44
2	B	579	3MS	C17-S33-N25	-3.82	102.26	106.56
2	A	579	3MS	C11-C12-C13	3.81	123.20	118.55
2	A	579	3MS	O26-C14-C12	-3.38	117.94	125.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	579	3MS	O26-C14-N24	3.18	123.33	120.52
2	B	579	3MS	O28-S32-C9	-3.13	105.39	109.30
2	B	579	3MS	C12-C14-N24	3.10	117.74	114.59
2	B	579	3MS	O26-C14-C12	-3.08	118.58	125.16
2	A	579	3MS	C12-C14-N24	3.05	117.70	114.59
2	B	579	3MS	O28-S32-O27	3.02	121.51	118.44
2	A	579	3MS	O30-S33-N25	2.69	112.47	107.10
2	B	579	3MS	C7-N21-C13	2.64	122.10	117.58
2	B	579	3MS	C14-C12-C13	-2.58	116.32	120.77
2	B	579	3MS	O30-S33-N25	2.51	112.11	107.10
2	A	579	3MS	O31-C11-C12	2.49	126.14	120.94
2	B	579	3MS	O31-C11-C12	2.21	125.56	120.94
2	A	579	3MS	C14-C12-C13	-2.15	117.08	120.77

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	579	3MS	C20-C18-C19-N24
2	B	579	3MS	C8-N25-S33-O30
2	A	579	3MS	C8-N25-S33-O30
2	A	579	3MS	C8-N25-S33-O29
2	B	579	3MS	C8-N25-S33-O29
2	A	579	3MS	C8-N25-S33-C17
2	B	579	3MS	C8-N25-S33-C17


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	579	3MS	1	0
2	B	579	3MS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	562/578 (97%)	0.71	50 (8%) 15 16	29, 40, 59, 94	0
1	B	562/578 (97%)	0.81	58 (10%) 12 12	29, 39, 62, 100	0
All	All	1124/1156 (97%)	0.76	108 (9%) 13 14	29, 40, 61, 100	0

All (108) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	545	LEU	6.6
1	B	550	TRP	6.2
1	B	545	LEU	6.2
1	B	150	GLU	6.2
1	B	541	ALA	5.6
1	B	532	THR	5.4
1	B	547	LEU	5.4
1	A	534	LEU	5.1
1	A	539	ILE	4.7
1	A	152	GLY	4.7
1	A	532	THR	4.6
1	A	547	LEU	4.6
1	B	536	LEU	4.5
1	B	539	ILE	4.5
1	B	544	GLN	4.4
1	B	534	LEU	4.3
1	A	15	ALA	4.1
1	A	541	ALA	4.1
1	B	546	ASP	4.1
1	A	536	LEU	4.0
1	A	543	SER	3.9
1	B	548	SER	3.9
1	A	14	CYS	3.8
1	A	538	PRO	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	535	LYS	3.7
1	B	95	HIS	3.6
1	A	153	GLY	3.6
1	A	549	GLY	3.6
1	B	533	LYS	3.5
1	A	542	ALA	3.5
1	B	540	PRO	3.5
1	A	550	TRP	3.5
1	B	549	GLY	3.5
1	B	14	CYS	3.4
1	B	535	LYS	3.4
1	B	542	ALA	3.4
1	B	15	ALA	3.4
1	B	329	THR	3.3
1	A	537	THR	3.2
1	B	149	PRO	3.2
1	A	95	HIS	3.2
1	A	548	SER	3.1
1	B	562	HIS	3.1
1	A	381	VAL	3.0
1	B	330	GLN	3.0
1	A	110	ASN	3.0
1	B	16	ALA	3.0
1	A	16	ALA	3.0
1	B	337	ARG	2.9
1	B	537	THR	2.9
1	A	150	GLU	2.9
1	A	533	LYS	2.9
1	B	35	ASN	2.8
1	A	562	HIS	2.8
1	B	509	ALA	2.7
1	A	544	GLN	2.7
1	B	311	CYS	2.7
1	B	327	ALA	2.7
1	A	311	CYS	2.7
1	B	223	CYS	2.7
1	B	153	GLY	2.7
1	B	543	SER	2.6
1	A	530	VAL	2.6
1	B	336	LEU	2.6
1	A	303	CYS	2.6
1	A	376	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	223	CYS	2.6
1	B	309	GLN	2.5
1	A	377	SER	2.5
1	A	546	ASP	2.5
1	B	513	SER	2.5
1	A	531	LYS	2.4
1	B	100	LYS	2.4
1	B	213	ASN	2.4
1	A	382	TYR	2.4
1	B	538	PRO	2.4
1	A	515	GLY	2.4
1	B	552	VAL	2.3
1	B	353	PRO	2.3
1	B	514	GLN	2.3
1	B	381	VAL	2.3
1	A	149	PRO	2.3
1	B	247	PRO	2.3
1	B	332	ASP	2.3
1	A	213	ASN	2.2
1	A	483	ASN	2.2
1	A	8	GLY	2.2
1	B	551	PHE	2.2
1	A	51	LYS	2.2
1	B	530	VAL	2.2
1	A	451	CYS	2.2
1	B	515	GLY	2.2
1	B	377	SER	2.1
1	B	473	SER	2.1
1	A	151	LYS	2.1
1	A	310	ASP	2.1
1	B	522	GLY	2.1
1	A	514	GLN	2.1
1	A	540	PRO	2.1
1	A	469	LEU	2.1
1	B	324	CYS	2.1
1	B	331	GLU	2.1
1	B	160	ILE	2.1
1	A	498	ARG	2.0
1	A	329	THR	2.0
1	B	505	ARG	2.0
1	B	502	HIS	2.0
1	B	382	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	3MS	A	579	33/33	0.97	0.08	32,34,36,36	0
2	3MS	B	579	33/33	0.97	0.07	32,34,38,39	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 3MS A 579:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 3MS B 579:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.