



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:05 PM UTC

PDB ID : 8CO3 / pdb_00008co3
Title : Three dimensional structure of human carbonic anhydrase XII in complex with sulfonamide
Authors : Leitans, J.; Dvinskis, E.; Tars, K.
Deposited on : 2023-02-26
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	 3% 89% 8% ..
1	B	263	 5% 90% 8% .
1	C	263	 5% 92% 6% .
1	D	263	 2% 89% 10% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9646 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	1	0
			2087	1328	355	397	7			
1	B	260	Total	C	N	O	S	0	0	0
			2078	1322	351	398	7			
1	C	260	Total	C	N	O	S	0	0	0
			2080	1323	354	396	7			
1	D	260	Total	C	N	O	S	0	1	0
			2091	1330	355	399	7			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	-	expression tag	UNP O43570
B	2	ALA	-	expression tag	UNP O43570
C	2	ALA	-	expression tag	UNP O43570
D	2	ALA	-	expression tag	UNP O43570

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 5-(5-methyl-6-quinolin-5-yl-pyridin-3-yl)thiophene-2-sulfonamide (CCD ID: V8O) (formula: C₁₉H₁₅N₃O₂S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	19	3	2	2		
3	B	1	Total	C	N	O	S	0	0
			26	19	3	2	2		
3	C	1	Total	C	N	O	S	0	0
			26	19	3	2	2		
3	D	1	Total	C	N	O	S	0	0
			26	19	3	2	2		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	339	Total	O	0	0
			339	339		
4	B	286	Total	O	0	0
			286	286		
4	C	282	Total	O	0	0
			282	282		
4	D	295	Total	O	0	0
			295	295		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carbonic anhydrase 12





- Molecule 1: Carbonic anhydrase 12



- Molecule 1: Carbonic anhydrase 12



- Molecule 1: Carbonic anhydrase 12



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.78Å 67.48Å 81.00Å 81.47° 84.09° 86.45°	Depositor
Resolution (Å)	79.74 – 1.68 79.74 – 1.68	Depositor EDS
% Data completeness (in resolution range)	85.5 (79.74-1.68) 85.5 (79.74-1.68)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.68Å)	Xtriage
Refinement program	REFMAC 5.8.0350	Depositor
R, R_{free}	0.199 , 0.254 0.208 , 0.261	Depositor DCC
R_{free} test set	4734 reflections (4.27%)	wwPDB-VP
Wilson B-factor (Å ²)	11.9	Xtriage
Anisotropy	0.147	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9646	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.03 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.1542e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, V8O

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2153	0.90	2/2933 (0.1%)
1	B	0.64	2/2141 (0.1%)	0.90	0/2919
1	C	0.61	0/2143	0.90	0/2921
1	D	0.65	2/2157 (0.1%)	0.94	4/2938 (0.1%)
All	All	0.63	4/8594 (0.0%)	0.91	6/11711 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	3
1	C	0	5
1	D	0	4
All	All	0	15

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	117	HIS	CE1-NE2	7.16	1.39	1.32
1	D	126	HIS	CE1-NE2	7.10	1.39	1.32
1	B	34	HIS	CE1-NE2	5.27	1.37	1.32
1	D	67	HIS	CE1-NE2	5.09	1.37	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	76	ASP	CA-CB-CG	6.14	118.74	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	226	GLU	CB-CG-CD	5.64	122.19	112.60
1	D	87	LEU	CA-C-N	5.46	127.86	120.38
1	D	87	LEU	C-N-CA	5.46	127.86	120.38
1	A	105	ASP	CA-CB-CG	5.41	118.01	112.60
1	A	126	HIS	CA-CB-CG	5.03	118.83	113.80

There are no chirality outliers.

All (15) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	218	ARG	Sidechain
1	A	246	ARG	Sidechain
1	A	261	ARG	Sidechain
1	B	192	ARG	Sidechain
1	B	200	ARG	Sidechain
1	B	261	ARG	Sidechain
1	C	192	ARG	Sidechain
1	C	198	ARG	Sidechain
1	C	200	ARG	Sidechain
1	C	218	ARG	Sidechain
1	C	246	ARG	Sidechain
1	D	198	ARG	Sidechain
1	D	218	ARG	Sidechain
1	D	246	ARG	Sidechain
1	D	90	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2087	0	1980	15	0
1	B	2078	0	1957	14	0
1	C	2080	0	1964	10	0
1	D	2091	0	1984	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
3	A	26	0	0	0	0
3	B	26	0	0	1	0
3	C	26	0	0	1	0
3	D	26	0	0	1	0
4	A	339	0	0	6	0
4	B	286	0	0	3	0
4	C	282	0	0	3	0
4	D	295	0	0	5	0
All	All	9646	0	7885	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:MET:SD	1:A:225:GLN:HG3	1.76	1.26
1:B:75:SER:HA	4:B:402:HOH:O	1.65	0.94
1:D:140:ASN:HB2	4:D:406:HOH:O	1.76	0.84
1:A:154:MET:SD	1:A:225:GLN:CG	2.65	0.82
1:C:241:ASP:OD2	4:C:1301:HOH:O	2.04	0.73
1:A:225:GLN:HE21	1:A:225:GLN:H	1.41	0.69
1:A:4:LYS:HB2	4:A:420:HOH:O	1.98	0.64
1:A:198:ARG:HD2	1:A:213:LEU:HD11	1.80	0.63
1:B:50:GLN:NE2	4:B:401:HOH:O	2.32	0.62
1:D:226:GLU:HB2	4:D:640:HOH:O	2.00	0.62
1:A:59:GLN:NE2	4:A:401:HOH:O	2.30	0.61
1:B:116:GLN:NE2	1:C:116:GLN:NE2	2.48	0.60
1:A:187:GLU:HG2	4:A:659:HOH:O	2.03	0.58
1:A:57:ASN:ND2	4:A:403:HOH:O	2.37	0.57
1:D:163:LYS:HD2	4:D:627:HOH:O	2.03	0.56
1:D:176:GLN:HE21	1:D:240:MET:HE3	1.70	0.55
1:B:116:GLN:HE21	1:C:116:GLN:NE2	2.04	0.55
1:B:116:GLN:HE21	1:C:116:GLN:HE22	1.56	0.52
1:D:47:LEU:HD22	1:D:82:ILE:CG2	2.39	0.52
1:A:195:GLU:OE2	1:A:218:ARG:NH1	2.26	0.52
1:B:203:LEU:HD22	3:B:302:V8O:S8	2.51	0.51
1:B:93:ALA:N	4:B:402:HOH:O	2.44	0.50
1:B:105:ASP:OD1	1:B:107:HIS:HD2	1.95	0.49
1:D:59:GLN:OE1	4:D:401:HOH:O	2.19	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:73:LEU:HD13	1:C:77:MET:HE1	1.96	0.47
1:A:58:LYS:HE2	1:A:60:PHE:CZ	2.50	0.46
1:D:176:GLN:NE2	1:D:240:MET:HE3	2.30	0.46
1:C:53:ASN:ND2	4:C:1302:HOH:O	2.48	0.45
1:A:259:ASP:O	1:A:261:ARG:HG3	2.16	0.45
1:C:204:THR:OG1	3:C:302:V8O:N1	2.49	0.45
1:A:70:LYS:NZ	4:A:412:HOH:O	2.44	0.45
1:A:29:SER:HB3	1:A:30:PRO:HA	1.99	0.44
4:A:472:HOH:O	1:D:104:ASN:HB3	2.18	0.44
1:C:165:PHE:HA	1:C:168:LEU:HG	1.99	0.44
1:C:141:LYS:HE2	4:C:1312:HOH:O	2.18	0.43
1:B:77:MET:HB3	1:B:93:ALA:HB3	2.00	0.43
1:A:184:ASN:HB3	1:A:187:GLU:HG3	2.01	0.43
1:B:58:LYS:HD3	1:B:60:PHE:CZ	2.54	0.43
1:D:261:ARG:HD3	4:D:501:HOH:O	2.18	0.42
1:B:116:GLN:NE2	1:C:116:GLN:HE21	2.18	0.42
1:B:124:ILE:HG22	1:B:126:HIS:CE1	2.55	0.41
1:D:203:LEU:HD22	3:D:302:V8O:S8	2.60	0.41
1:D:96:LEU:C	1:D:96:LEU:HD12	2.46	0.41
1:A:116:GLN:NE2	1:D:116:GLN:NE2	2.68	0.41
1:D:53:ASN:HA	1:D:184:ASN:HA	2.02	0.40
1:B:259:ASP:O	1:B:261:ARG:HG3	2.22	0.40
1:D:39:GLN:O	1:D:264:TYR:HA	2.21	0.40
1:B:20:TYR:CD2	1:B:206:PRO:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	259/263 (98%)	254 (98%)	5 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	258/263 (98%)	252 (98%)	6 (2%)	0	100	100
1	C	258/263 (98%)	252 (98%)	6 (2%)	0	100	100
1	D	259/263 (98%)	254 (98%)	4 (2%)	1 (0%)	30	13
All	All	1034/1052 (98%)	1012 (98%)	21 (2%)	1 (0%)	48	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	240	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/234 (99%)	227 (98%)	5 (2%)	45	19
1	B	230/234 (98%)	228 (99%)	2 (1%)	70	55
1	C	230/234 (98%)	225 (98%)	5 (2%)	45	19
1	D	233/234 (100%)	231 (99%)	2 (1%)	70	55
All	All	925/936 (99%)	911 (98%)	14 (2%)	57	34

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LYS
1	A	70	LYS
1	A	83	GLN
1	A	95	GLN
1	A	225	GLN
1	B	95	GLN
1	B	143	GLU
1	C	70	LYS
1	C	82	ILE
1	C	95	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	162	ASP
1	C	241	ASP
1	D	162	ASP
1	D	200	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	67	HIS
1	A	88	GLN
1	A	107	HIS
1	A	116	GLN
1	A	225	GLN
1	A	239	HIS
1	B	83	GLN
1	B	107	HIS
1	B	116	GLN
1	C	14	ASN
1	C	50	GLN
1	C	53	ASN
1	C	57	ASN
1	C	67	HIS
1	C	83	GLN
1	C	116	GLN
1	C	219	ASN
1	D	34	HIS
1	D	39	GLN
1	D	57	ASN
1	D	59	GLN
1	D	102	ASN
1	D	104	ASN
1	D	116	GLN
1	D	117	HIS
1	D	250	ASN
1	D	256	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	V8O	B	302	2	28,29,29	1.88	7 (25%)	39,43,43	3.55	16 (41%)
3	V8O	D	302	2	28,29,29	1.89	6 (21%)	39,43,43	3.56	15 (38%)
3	V8O	A	302	2	28,29,29	1.65	4 (14%)	39,43,43	3.80	15 (38%)
3	V8O	C	302	2	28,29,29	1.98	6 (21%)	39,43,43	3.30	14 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	V8O	B	302	2	-	5/13/14/14	0/4/4/4
3	V8O	D	302	2	-	2/13/14/14	0/4/4/4
3	V8O	A	302	2	-	6/13/14/14	0/4/4/4
3	V8O	C	302	2	-	2/13/14/14	0/4/4/4

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	302	V8O	C15-C14	5.65	1.49	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	302	V8O	C15-C14	5.42	1.49	1.40
3	B	302	V8O	C15-C14	5.00	1.48	1.40
3	A	302	V8O	C15-C14	4.90	1.48	1.40
3	B	302	V8O	C24-C23	4.43	1.49	1.42
3	D	302	V8O	C24-C23	4.31	1.49	1.42
3	A	302	V8O	C24-C23	4.18	1.49	1.42
3	C	302	V8O	C24-C23	4.15	1.49	1.42
3	C	302	V8O	S4-N1	3.61	1.66	1.58
3	C	302	V8O	C19-C24	3.53	1.50	1.42
3	B	302	V8O	C7-S4	-3.41	1.70	1.75
3	B	302	V8O	C19-C24	3.16	1.50	1.42
3	D	302	V8O	C7-S4	-3.09	1.70	1.75
3	D	302	V8O	C7-S8	2.90	1.76	1.71
3	A	302	V8O	C19-C24	2.73	1.49	1.42
3	C	302	V8O	C7-S8	2.70	1.76	1.71
3	D	302	V8O	S4-N1	2.69	1.64	1.58
3	C	302	V8O	C7-S4	-2.57	1.71	1.75
3	D	302	V8O	C19-C24	2.55	1.48	1.42
3	A	302	V8O	C7-S4	-2.45	1.71	1.75
3	B	302	V8O	O5-S4	2.43	1.47	1.43
3	B	302	V8O	C7-S8	2.16	1.75	1.71
3	B	302	V8O	S4-N1	2.07	1.63	1.58

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	V8O	O5-S4-N1	14.12	128.05	108.39
3	B	302	V8O	O5-S4-N1	13.93	127.80	108.39
3	D	302	V8O	O6-S4-N1	13.06	126.58	108.39
3	A	302	V8O	O5-S4-O6	-12.33	102.64	120.27
3	C	302	V8O	O6-S4-N1	11.28	124.10	108.39
3	B	302	V8O	O5-S4-O6	-9.52	106.66	120.27
3	C	302	V8O	O5-S4-O6	-8.69	107.85	120.27
3	D	302	V8O	O5-S4-O6	-7.47	109.58	120.27
3	C	302	V8O	C14-C15-N16	-6.75	117.43	123.08
3	D	302	V8O	C19-C15-N16	6.50	127.69	115.27
3	C	302	V8O	S8-C7-S4	6.40	128.97	120.46
3	D	302	V8O	C19-C15-C14	-6.30	112.49	122.20
3	A	302	V8O	C14-C15-N16	-6.19	117.90	123.08
3	B	302	V8O	S8-C7-S4	6.09	128.56	120.46
3	A	302	V8O	S8-C7-S4	5.91	128.32	120.46
3	B	302	V8O	C14-C15-N16	-5.67	118.34	123.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	302	V8O	S8-C7-S4	5.43	127.69	120.46
3	D	302	V8O	C24-C23-N28	-4.99	117.53	122.82
3	D	302	V8O	C7-S8-C9	4.75	94.63	90.41
3	C	302	V8O	C7-S8-C9	4.70	94.58	90.41
3	A	302	V8O	C7-S8-C9	4.52	94.42	90.41
3	D	302	V8O	C11-C7-S8	-4.37	108.20	112.55
3	B	302	V8O	C24-C23-N28	-4.17	118.40	122.82
3	A	302	V8O	C11-C7-S8	-4.06	108.51	112.55
3	A	302	V8O	C24-C23-N28	-4.02	118.56	122.82
3	C	302	V8O	C11-C7-S8	-4.00	108.57	112.55
3	D	302	V8O	C13-C12-C9	-3.92	115.44	120.97
3	D	302	V8O	C14-C15-N16	-3.91	119.81	123.08
3	C	302	V8O	O5-S4-N1	-3.84	103.04	108.39
3	B	302	V8O	O6-S4-N1	-3.81	103.09	108.39
3	B	302	V8O	C12-C9-C10	3.69	135.37	128.47
3	C	302	V8O	C24-C23-N28	-3.49	119.12	122.82
3	A	302	V8O	C17-C12-C9	-3.34	116.85	121.29
3	B	302	V8O	C7-S8-C9	3.28	93.32	90.41
3	C	302	V8O	C27-N28-C23	3.27	121.88	116.93
3	B	302	V8O	C17-C12-C9	-3.26	116.96	121.29
3	D	302	V8O	C27-N28-C23	3.18	121.74	116.93
3	B	302	V8O	C27-N28-C23	2.95	121.39	116.93
3	B	302	V8O	C11-C7-S8	-2.87	109.69	112.55
3	A	302	V8O	C12-C9-C10	2.72	133.55	128.47
3	C	302	V8O	C17-C12-C9	-2.69	117.71	121.29
3	D	302	V8O	C22-C23-N28	2.69	122.89	118.55
3	C	302	V8O	C17-N16-C15	2.68	122.63	118.01
3	B	302	V8O	C7-S4-N1	2.65	110.34	107.51
3	A	302	V8O	C27-N28-C23	2.59	120.85	116.93
3	A	302	V8O	C18-C14-C15	-2.48	118.86	122.34
3	D	302	V8O	C12-C9-C10	2.48	133.10	128.47
3	A	302	V8O	C7-S4-N1	2.45	110.12	107.51
3	B	302	V8O	C12-C9-S8	-2.44	114.81	121.02
3	D	302	V8O	C17-C12-C9	2.42	124.51	121.29
3	B	302	V8O	C22-C23-N28	2.37	122.37	118.55
3	A	302	V8O	C17-N16-C15	2.33	122.04	118.01
3	C	302	V8O	C7-S4-N1	2.32	109.99	107.51
3	B	302	V8O	C11-C7-S4	-2.30	121.74	126.69
3	C	302	V8O	C12-C9-C10	2.30	132.76	128.47
3	A	302	V8O	C26-C25-C24	-2.14	118.02	120.91
3	C	302	V8O	C19-C15-C14	2.12	125.48	122.20
3	D	302	V8O	C13-C12-C17	2.07	120.05	117.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	V8O	C17-N16-C15	2.07	121.58	118.01
3	A	302	V8O	C12-C13-C14	-2.01	119.27	121.74

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	V8O	C11-C7-S4-O6
3	A	302	V8O	S8-C7-S4-O6
3	A	302	V8O	C17-C12-C9-C10
3	A	302	V8O	C17-C12-C9-S8
3	B	302	V8O	C11-C7-S4-O6
3	B	302	V8O	S8-C7-S4-O6
3	C	302	V8O	C11-C7-S4-O5
3	D	302	V8O	C11-C7-S4-O5
3	D	302	V8O	S8-C7-S4-O5
3	A	302	V8O	C13-C12-C9-S8
3	C	302	V8O	S8-C7-S4-O5
3	A	302	V8O	C13-C12-C9-C10
3	B	302	V8O	S8-C7-S4-N1
3	B	302	V8O	C17-C12-C9-S8
3	B	302	V8O	C17-C12-C9-C10


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	V8O	1	0
3	D	302	V8O	1	0
3	C	302	V8O	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/263 (98%)	0.33	7 (2%) 56 63	6, 13, 28, 54	1 (0%)
1	B	260/263 (98%)	0.38	12 (4%) 37 46	5, 13, 29, 48	0
1	C	260/263 (98%)	0.52	13 (5%) 34 42	7, 16, 32, 41	0
1	D	260/263 (98%)	0.36	4 (1%) 72 80	6, 15, 27, 40	1 (0%)
All	All	1040/1052 (98%)	0.40	36 (3%) 47 55	5, 14, 28, 54	2 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	243	PRO	4.3
1	C	243	PRO	3.5
1	C	140	ASN	3.3
1	C	131	LEU	3.1
1	D	8	PHE	3.1
1	B	143	GLU	3.0
1	B	243	PRO	3.0
1	A	260	GLU	3.0
1	B	244	SER	3.0
1	B	8	PHE	2.8
1	C	240	MET	2.8
1	B	260	GLU	2.7
1	C	260	GLU	2.7
1	A	8	PHE	2.7
1	B	179	PHE	2.7
1	C	8	PHE	2.7
1	B	26	LEU	2.5
1	B	245	PRO	2.5
1	C	88	GLN	2.5
1	C	138	ALA	2.5
1	C	259	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	241	ASP	2.4
1	A	242	ASP	2.4
1	C	191	GLU	2.4
1	B	142	SER	2.3
1	B	131	LEU	2.3
1	A	241	ASP	2.3
1	C	142	SER	2.3
1	C	192	ARG	2.2
1	D	244	SER	2.2
1	A	143	GLU	2.1
1	A	4	LYS	2.1
1	B	240	MET	2.1
1	B	140	ASN	2.0
1	D	260	GLU	2.0
1	D	59	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	V8O	C	302	26/26	0.93	0.13	14,28,37,40	0
3	V8O	D	302	26/26	0.93	0.12	11,25,33,33	0
3	V8O	B	302	26/26	0.94	0.12	11,24,36,36	0
3	V8O	A	302	26/26	0.95	0.09	9,22,27,28	0
2	ZN	A	301	1/1	0.99	0.01	7,7,7,7	0
2	ZN	B	301	1/1	0.99	0.02	7,7,7,7	0
2	ZN	C	301	1/1	0.99	0.02	10,10,10,10	0
2	ZN	D	301	1/1	0.99	0.02	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around V8O C 302:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around V8O D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around V8O B 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.