



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:14 AM UTC

PDB ID : 8CPL / pdb_00008cpl
Title : YZw2 a scaffold for cryo-EM of small proteins of interest
Authors : Moche, M.; Friberg, O.; Nygren, P.A.; Nilvebrant, J.
Deposited on : 2023-03-03
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	3% 85% 5% 10%
1	B	501	2% 86% 5% 9%
1	C	501	8% 94% 6%
1	D	501	2% 85% 5% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putrescine aminotransferase,Immunoglobulin G-binding protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	452	Total	C	N	O	S	0	21	0
			3610	2284	631	671	24			
1	B	456	Total	C	N	O	S	0	24	0
			3665	2315	644	681	25			
1	C	499	Total	C	N	O	S	0	20	0
			3973	2511	696	742	24			
1	D	456	Total	C	N	O	S	0	25	0
			3662	2315	641	681	25			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	478	ALA	GLY	conflict	UNP P38507
B	478	ALA	GLY	conflict	UNP P38507
C	478	ALA	GLY	conflict	UNP P38507
D	478	ALA	GLY	conflict	UNP P38507

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	B	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	C	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	D	1	Total	C	N	O	P	0	0
			16	8	1	6	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	534	Total	O	0	0
			534	534		
3	B	582	Total	O	0	0
			582	582		
3	C	510	Total	O	0	0
			510	510		
3	D	568	Total	O	0	0
			568	568		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putrescine aminotransferase,Immunoglobulin G-binding protein A




- Molecule 1: Putrescine aminotransferase,Immunoglobulin G-binding protein A




- Molecule 1: Putrescine aminotransferase,Immunoglobulin G-binding protein A



- Molecule 1: Putrescine aminotransferase,Immunoglobulin G-binding protein A





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.45Å 173.90Å 209.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.44 – 1.60 49.44 – 1.60	Depositor EDS
% Data completeness (in resolution range)	58.4 (49.44-1.60) 58.4 (49.44-1.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.61Å)	Xtrriage
Refinement program	BUSTER 2.10.4 (26-JUL-2023)	Depositor
R, R_{free}	0.188 , 0.210 0.185 , 0.207	Depositor DCC
R_{free} test set	10086 reflections (2.90%)	wwPDB-VP
Wilson B-factor (Å ²)	16.1	Xtrriage
Anisotropy	0.026	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	17168	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.9587e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/3674	0.95	0/4967
1	B	0.79	0/3729	0.94	1/5039 (0.0%)
1	C	0.77	0/4043	0.95	0/5464
1	D	0.78	0/3726	0.94	1/5036 (0.0%)
All	All	0.78	0/15172	0.95	2/20506 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	379	ASP	CA-CB-CG	5.35	117.95	112.60
1	B	379	ASP	CA-CB-CG	5.32	117.92	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3610	0	3634	22	0
1	B	3665	0	3678	24	0
1	C	3973	0	3996	25	0
1	D	3662	0	3674	24	0
2	A	16	0	7	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	16	0	7	4	0
2	C	16	0	7	3	0
2	D	16	0	7	3	0
3	A	534	0	0	1	0
3	B	582	0	0	1	0
3	C	510	0	0	1	0
3	D	568	0	0	2	0
All	All	17168	0	15010	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:331:THR:HG22	1:C:334:GLY:H	1.39	0.87
1:D:331:THR:HG22	1:D:334:GLY:H	1.40	0.86
2:A:601:PLP:O3P	1:B:332[A]:THR:HG23	1.76	0.85
1:C:332[B]:THR:HG22	1:D:93:ILE:HD11	1.59	0.83
1:B:331:THR:HG22	1:B:334:GLY:H	1.40	0.82
1:C:332[B]:THR:CG2	1:D:93:ILE:HD11	2.17	0.74
1:C:274[B]:GLN:NE2	1:C:428:GLU:OE1	2.26	0.69
2:C:601:PLP:O2P	1:D:332[A]:THR:HG23	1.92	0.69
1:C:332[A]:THR:HG23	2:D:601:PLP:O2P	1.93	0.67
1:C:93:ILE:HD11	1:D:332[B]:THR:CG2	2.25	0.67
1:A:93:ILE:HD11	1:B:332[B]:THR:CG2	2.25	0.66
1:C:332[A]:THR:HG21	1:D:300:LYS:HE3	1.79	0.64
1:C:93:ILE:HD11	1:D:332[B]:THR:HG22	1.82	0.62
1:A:332[A]:THR:HG23	2:B:601:PLP:O3P	2.00	0.62
1:A:332[A]:THR:HG21	1:B:300:LYS:HE3	1.81	0.62
1:A:108[B]:GLN:OE1	1:B:108[B]:GLN:OE1	2.18	0.61
1:C:108[B]:GLN:OE1	1:D:108[B]:GLN:OE1	2.21	0.58
1:B:331:THR:HG22	1:B:334:GLY:N	2.17	0.57
1:D:331:THR:HG22	1:D:334:GLY:N	2.17	0.57
1:A:332[B]:THR:CG2	1:B:93:ILE:HD11	2.34	0.57
1:C:331:THR:HG22	1:C:334:GLY:N	2.16	0.56
1:A:331[A]:THR:HG22	1:A:334:GLY:H	1.70	0.55
2:A:601:PLP:O2P	1:B:332[A]:THR:OG1	2.24	0.55
1:C:465:ILE:HD13	1:C:479:PHE:HB3	1.88	0.55
1:C:332[A]:THR:OG1	2:D:601:PLP:O1P	2.25	0.55
1:C:91:PHE:O	1:C:274[B]:GLN:OE1	2.26	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274[B]:GLN:NE2	1:A:428:GLU:OE1	2.41	0.53
1:C:84:PHE:CD2	1:C:414[B]:LEU:HD13	2.43	0.53
1:B:82[B]:GLN:OE1	1:B:412[B]:ARG:HD3	2.10	0.51
1:A:332[A]:THR:OG1	2:B:601:PLP:O2P	2.27	0.51
2:C:601:PLP:O1P	1:D:332[A]:THR:OG1	2.29	0.50
1:D:274[B]:GLN:NE2	1:D:428:GLU:OE1	2.44	0.50
1:B:222[A]:LEU:HD22	1:B:232:VAL:HG21	1.94	0.50
1:B:13[B]:CYS:SG	3:B:1045:HOH:O	2.60	0.50
1:C:222[B]:LEU:CD1	1:C:260:LEU:HG	2.42	0.49
1:C:331:THR:CG2	1:C:334:GLY:H	2.19	0.49
1:D:191:THR:O	1:D:197:ARG:HD2	2.13	0.49
1:C:89:GLY:HA3	3:C:788:HOH:O	2.12	0.48
1:D:82[B]:GLN:OE1	1:D:412[B]:ARG:HD3	2.13	0.48
1:A:93:ILE:HD11	1:B:332[B]:THR:HG22	1.96	0.48
1:C:401:GLY:HA3	1:C:422:ALA:O	2.14	0.47
1:C:300:LYS:HE3	1:D:332[A]:THR:HG21	1.96	0.47
1:C:465:ILE:HD12	1:C:483:LEU:HD11	1.96	0.47
1:A:332[B]:THR:HG23	1:B:93:ILE:HD11	1.97	0.47
1:B:401:GLY:HA3	1:B:422:ALA:O	2.15	0.47
1:B:191:THR:O	1:B:197:ARG:HD2	2.15	0.47
1:A:89:GLY:HA3	3:A:762:HOH:O	2.13	0.46
1:D:274[A]:GLN:NE2	2:D:601:PLP:O3	2.48	0.46
1:A:93:ILE:HD11	1:B:332[B]:THR:HG23	1.98	0.45
1:D:331:THR:CG2	1:D:334:GLY:H	2.21	0.45
1:A:352:GLU:HG2	1:B:11:LEU:HD11	1.98	0.45
1:D:401:GLY:HA3	1:D:422:ALA:O	2.15	0.45
1:A:332[B]:THR:OG1	2:B:601:PLP:O3P	2.29	0.45
1:C:352:GLU:HG2	1:D:11:LEU:HD11	1.98	0.45
1:D:208[B]:ARG:NH2	3:D:713:HOH:O	2.50	0.45
1:A:401:GLY:HA3	1:A:422:ALA:O	2.16	0.45
1:C:92:GLY:HA3	1:C:274[B]:GLN:HE22	1.81	0.45
1:A:41:TYR:HB2	1:B:127:ALA:HB1	1.98	0.44
1:A:300:LYS:HE3	1:B:332[A]:THR:HG21	2.00	0.44
1:C:41:TYR:HB2	1:D:127:ALA:HB1	1.98	0.44
1:C:191:THR:O	1:C:197:ARG:HD2	2.17	0.44
1:A:120:GLU:HG2	1:B:414:LEU:HD11	2.01	0.43
1:A:127:ALA:HB1	1:B:41:TYR:HB2	1.99	0.43
1:B:274[A]:GLN:NE2	2:B:601:PLP:O3	2.50	0.43
1:A:191:THR:O	1:A:197:ARG:HD2	2.19	0.43
1:D:222[B]:LEU:HD12	1:D:260:LEU:HG	2.01	0.42
1:A:11:LEU:HD11	1:B:352:GLU:HG2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:601:PLP:O2P	1:D:332[B]:THR:OG1	2.28	0.42
1:A:76:LEU:HD21	1:A:98:HIS:CE1	2.56	0.41
1:B:331:THR:CG2	1:B:334:GLY:H	2.21	0.40
1:D:20[B]:LEU:HD11	1:D:34:LEU:HD22	2.04	0.40
1:D:447[A]:LYS:NZ	3:D:703:HOH:O	2.36	0.40
1:C:274[A]:GLN:HE21	1:C:274[A]:GLN:HB2	1.71	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/501 (94%)	451 (96%)	19 (4%)	1 (0%)	43	24
1	B	478/501 (95%)	457 (96%)	20 (4%)	1 (0%)	43	24
1	C	517/501 (103%)	495 (96%)	21 (4%)	1 (0%)	43	24
1	D	478/501 (95%)	457 (96%)	20 (4%)	1 (0%)	43	24
All	All	1944/2004 (97%)	1860 (96%)	80 (4%)	4 (0%)	43	24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	ALA
1	B	299	ALA
1	C	299	ALA
1	D	299	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/402 (95%)	371 (97%)	11 (3%)	37	14
1	B	386/402 (96%)	377 (98%)	9 (2%)	44	20
1	C	420/402 (104%)	414 (99%)	6 (1%)	59	38
1	D	386/402 (96%)	378 (98%)	8 (2%)	47	23
All	All	1574/1608 (98%)	1540 (98%)	34 (2%)	53	22

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	GLN
1	A	222[A]	LEU
1	A	222[B]	LEU
1	A	254	LEU
1	A	274[A]	GLN
1	A	274[B]	GLN
1	A	331[A]	THR
1	A	331[B]	THR
1	A	361	GLN
1	A	419	LEU
1	A	458	GLN
1	B	119	GLN
1	B	222[A]	LEU
1	B	222[B]	LEU
1	B	254	LEU
1	B	274[A]	GLN
1	B	274[B]	GLN
1	B	331	THR
1	B	436[A]	GLU
1	B	436[B]	GLU
1	C	119	GLN
1	C	208[A]	ARG
1	C	208[B]	ARG
1	C	254	LEU
1	C	331	THR
1	C	419	LEU
1	D	119	GLN
1	D	222[A]	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	222[B]	LEU
1	D	254	LEU
1	D	331	THR
1	D	419	LEU
1	D	436[A]	GLU
1	D	436[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	80	GLN
1	A	361	GLN
1	A	411	GLN
1	B	74	ASN
1	B	241	GLN
1	C	74	ASN
1	C	80	GLN
1	C	110	GLN
1	D	74	ASN
1	D	80	GLN
1	D	411	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	C	601	-	16,16,16	0.47	0	20,23,23	0.72	0
2	PLP	A	601	-	16,16,16	0.42	0	20,23,23	0.76	0
2	PLP	D	601	-	16,16,16	0.45	0	20,23,23	0.70	0
2	PLP	B	601	-	16,16,16	0.42	0	20,23,23	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	C	601	-	-	2/8/8/8	0/1/1/1
2	PLP	A	601	-	-	2/8/8/8	0/1/1/1
2	PLP	D	601	-	-	2/8/8/8	0/1/1/1
2	PLP	B	601	-	-	2/8/8/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	PLP	C3-C4-C4A-O4A
2	C	601	PLP	C3-C4-C4A-O4A
2	A	601	PLP	C3-C4-C4A-O4A
2	D	601	PLP	C3-C4-C4A-O4A
2	A	601	PLP	C5-C4-C4A-O4A
2	C	601	PLP	C5-C4-C4A-O4A
2	B	601	PLP	C5-C4-C4A-O4A
2	D	601	PLP	C5-C4-C4A-O4A

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	601	PLP	3	0
2	A	601	PLP	2	0
2	D	601	PLP	3	0
2	B	601	PLP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/501 (90%)	-0.13	14 (3%) 51 54	4, 15, 37, 78	21 (4%)
1	B	456/501 (91%)	-0.21	9 (1%) 65 69	6, 16, 30, 89	24 (5%)
1	C	499/501 (99%)	0.18	40 (8%) 18 18	5, 17, 75, 84	20 (4%)
1	D	456/501 (91%)	-0.20	12 (2%) 57 60	6, 16, 31, 86	25 (5%)
All	All	1863/2004 (92%)	-0.08	75 (4%) 42 44	4, 16, 40, 89	90 (4%)

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	462	PHE	7.2
1	C	465	ILE	5.7
1	C	483	LEU	5.5
1	B	463	TYR	5.5
1	D	462	PHE	5.4
1	C	466	LEU	5.3
1	C	494	LEU	5.2
1	D	461	ALA	5.0
1	D	463	TYR	5.0
1	C	462	PHE	4.7
1	C	463	TYR	4.5
1	D	464	GLU	4.5
1	C	493	LEU	4.5
1	C	505	ALA	4.3
1	A	456	VAL	4.3
1	C	491	ALA	4.2
1	B	461	ALA	4.2
1	C	495	ALA	4.2
1	C	471	LEU	4.2
1	C	479	PHE	4.2
1	C	503	ALA	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	500	LEU	3.9
1	C	468	LEU	3.8
1	C	461	ALA	3.7
1	C	506	PRO	3.7
1	C	485	ASP	3.6
1	C	480	ILE	3.5
1	C	469	PRO	3.5
1	C	460	ASN	3.4
1	C	484	LYS	3.4
1	C	478	ALA	3.3
1	C	497	ALA	3.0
1	D	458	GLN	3.0
1	C	472	ASN	3.0
1	C	24[A]	ARG	2.9
1	A	460	ASN	2.9
1	C	487	PRO	2.9
1	C	507	LYS	2.7
1	B	456	VAL	2.7
1	C	490	SER	2.6
1	C	488	SER	2.6
1	A	454	VAL	2.6
1	A	459	GLN	2.6
1	D	456	VAL	2.6
1	B	332[A]	THR	2.5
1	A	59	ALA	2.5
1	B	458	GLN	2.5
1	C	459	GLN	2.5
1	C	481	GLN	2.5
1	A	24[A]	ARG	2.5
1	B	460	ASN	2.5
1	A	53	TYR	2.4
1	C	63	TYR	2.4
1	A	71	GLY	2.4
1	A	331[A]	THR	2.4
1	C	464	GLU	2.3
1	A	368	ASP	2.3
1	C	23[A]	LYS	2.3
1	D	332[A]	THR	2.3
1	D	324	ASP	2.3
1	A	421	ASN	2.2
1	D	331	THR	2.2
1	C	486	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	474	GLU	2.1
1	B	331	THR	2.1
1	A	243	GLU	2.1
1	D	36	ARG	2.1
1	D	460	ASN	2.1
1	D	352	GLU	2.1
1	C	420	ASN	2.1
1	C	492	ASN	2.1
1	A	61	GLY	2.0
1	A	458	GLN	2.0
1	C	504	GLN	2.0
1	B	244	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLP	C	601	16/16	0.91	0.10	17,20,22,23	16
2	PLP	B	601	16/16	0.92	0.09	14,15,18,21	16
2	PLP	A	601	16/16	0.92	0.09	17,18,22,23	16
2	PLP	D	601	16/16	0.92	0.10	17,18,20,22	16

6.5 Other polymers [i](#)

There are no such residues in this entry.