



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:21 AM UTC

PDB ID : 4CQL / pdb_00004cql
Title : Crystal structure of heterotetrameric human ketoacyl reductase complexed with NAD
Authors : Venkatesan, R.; Sah-Teli, S.K.; Awoniyi, L.O.; Jiang, G.; Prus, P.; Kastaniotis, A.J.; Hiltunen, J.K.; Wierenga, R.K.; Chen, Z.
Deposited on : 2014-02-19
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	 82% 13% 6%
1	D	261	 84% 10% 6%
1	E	261	 80% 14% 7%
1	H	261	 82% 10% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	261	 82% 14%
1	L	261	 76% 15% 8%
1	M	261	 75% 19% 6%
1	P	261	 79% 15% 6%
2	B	244	 77% 16% 7%
2	C	244	 73% 15% 11%
2	F	244	 74% 20% 5%
2	G	244	 69% 18% 13%
2	J	244	 73% 20% 6%
2	K	244	 74% 16% 10%
2	N	244	 73% 20% 6%
2	O	244	 73% 18% 8%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 26833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ESTRADIOL 17-BETA-DEHYDROGENASE 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	246	Total 1756	C 1090	N 313	O 342	S 11	0	0	0
1	D	246	Total 1721	C 1070	N 302	O 339	S 10	0	0	0
1	E	244	Total 1733	C 1075	N 308	O 339	S 11	0	0	0
1	H	242	Total 1710	C 1059	N 306	O 335	S 10	0	0	0
1	I	251	Total 1762	C 1097	N 309	O 345	S 11	0	0	0
1	L	240	Total 1697	C 1053	N 302	O 331	S 11	0	0	0
1	M	245	Total 1755	C 1089	N 315	O 341	S 10	0	0	0
1	P	246	Total 1739	C 1080	N 308	O 341	S 10	0	0	0

- Molecule 2 is a protein called CARBONYL REDUCTASE FAMILY MEMBER 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	227	Total 1593	C 1000	N 293	O 291	S 9	0	0	0
2	C	216	Total 1496	C 942	N 266	O 280	S 8	0	0	0
2	F	231	Total 1623	C 1013	N 297	O 304	S 9	0	0	0
2	G	212	Total 1452	C 905	N 263	O 276	S 8	0	0	0
2	J	229	Total 1609	C 1010	N 292	O 298	S 9	0	0	0
2	K	220	Total 1493	C 934	N 271	O 280	S 8	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	229	Total	C	N	O	S	0	0	0
			1616	1014	294	299	9			
2	O	224	Total	C	N	O	S	0	0	0
			1508	946	269	285	8			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	MET	-	expression tag	UNP Q8N4T8
B	-5	HIS	-	expression tag	UNP Q8N4T8
B	-4	HIS	-	expression tag	UNP Q8N4T8
B	-3	HIS	-	expression tag	UNP Q8N4T8
B	-2	HIS	-	expression tag	UNP Q8N4T8
B	-1	HIS	-	expression tag	UNP Q8N4T8
B	0	HIS	-	expression tag	UNP Q8N4T8
C	-6	MET	-	expression tag	UNP Q8N4T8
C	-5	HIS	-	expression tag	UNP Q8N4T8
C	-4	HIS	-	expression tag	UNP Q8N4T8
C	-3	HIS	-	expression tag	UNP Q8N4T8
C	-2	HIS	-	expression tag	UNP Q8N4T8
C	-1	HIS	-	expression tag	UNP Q8N4T8
C	0	HIS	-	expression tag	UNP Q8N4T8
F	-6	MET	-	expression tag	UNP Q8N4T8
F	-5	HIS	-	expression tag	UNP Q8N4T8
F	-4	HIS	-	expression tag	UNP Q8N4T8
F	-3	HIS	-	expression tag	UNP Q8N4T8
F	-2	HIS	-	expression tag	UNP Q8N4T8
F	-1	HIS	-	expression tag	UNP Q8N4T8
F	0	HIS	-	expression tag	UNP Q8N4T8
G	-6	MET	-	expression tag	UNP Q8N4T8
G	-5	HIS	-	expression tag	UNP Q8N4T8
G	-4	HIS	-	expression tag	UNP Q8N4T8
G	-3	HIS	-	expression tag	UNP Q8N4T8
G	-2	HIS	-	expression tag	UNP Q8N4T8
G	-1	HIS	-	expression tag	UNP Q8N4T8
G	0	HIS	-	expression tag	UNP Q8N4T8
J	-6	MET	-	expression tag	UNP Q8N4T8
J	-5	HIS	-	expression tag	UNP Q8N4T8
J	-4	HIS	-	expression tag	UNP Q8N4T8
J	-3	HIS	-	expression tag	UNP Q8N4T8
J	-2	HIS	-	expression tag	UNP Q8N4T8
J	-1	HIS	-	expression tag	UNP Q8N4T8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	0	HIS	-	expression tag	UNP Q8N4T8
K	-6	MET	-	expression tag	UNP Q8N4T8
K	-5	HIS	-	expression tag	UNP Q8N4T8
K	-4	HIS	-	expression tag	UNP Q8N4T8
K	-3	HIS	-	expression tag	UNP Q8N4T8
K	-2	HIS	-	expression tag	UNP Q8N4T8
K	-1	HIS	-	expression tag	UNP Q8N4T8
K	0	HIS	-	expression tag	UNP Q8N4T8
N	-6	MET	-	expression tag	UNP Q8N4T8
N	-5	HIS	-	expression tag	UNP Q8N4T8
N	-4	HIS	-	expression tag	UNP Q8N4T8
N	-3	HIS	-	expression tag	UNP Q8N4T8
N	-2	HIS	-	expression tag	UNP Q8N4T8
N	-1	HIS	-	expression tag	UNP Q8N4T8
N	0	HIS	-	expression tag	UNP Q8N4T8
O	-6	MET	-	expression tag	UNP Q8N4T8
O	-5	HIS	-	expression tag	UNP Q8N4T8
O	-4	HIS	-	expression tag	UNP Q8N4T8
O	-3	HIS	-	expression tag	UNP Q8N4T8
O	-2	HIS	-	expression tag	UNP Q8N4T8
O	-1	HIS	-	expression tag	UNP Q8N4T8
O	0	HIS	-	expression tag	UNP Q8N4T8

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	H	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	I	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	L	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	M	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	P	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	24	Total	O	0	0
			24	24		
4	B	7	Total	O	0	0
			7	7		
4	C	10	Total	O	0	0
			10	10		
4	D	4	Total	O	0	0
			4	4		
4	E	25	Total	O	0	0
			25	25		
4	F	8	Total	O	0	0
			8	8		
4	G	4	Total	O	0	0
			4	4		
4	H	9	Total	O	0	0
			9	9		
4	I	14	Total	O	0	0
			14	14		
4	J	8	Total	O	0	0
			8	8		
4	K	7	Total	O	0	0
			7	7		
4	L	20	Total	O	0	0
			20	20		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	M	38	Total 38	O 38	0	0
4	N	24	Total 24	O 24	0	0
4	O	6	Total 6	O 6	0	0
4	P	10	Total 10	O 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8




- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8




- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8




- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8






- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8

Chain I: 82% 14%




- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8

Chain L: 76% 15% 8%




- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8

Chain M: 75% 19% 6%




- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8


Chain P: 79% 15% 6%



- Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain B: 77% 16% 7%







- Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain C: 73% 15% 11%



- Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain F: 74% 20% 5%




- Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain G: 69% 18% 13%




- Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain J: 73% 20% 6%




- Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain K: 74% 16% 10%




• Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain N: 73% 20% • 6%



• Molecule 2: CARBONYL REDUCTASE FAMILY MEMBER 4

Chain O: 73% 18% • 8%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.15Å 237.11Å 87.35Å 90.00° 94.52° 90.00°	Depositor
Resolution (Å)	49.78 – 2.85 49.78 – 2.85	Depositor EDS
% Data completeness (in resolution range)	91.7 (49.78-2.85) 87.2 (49.78-2.85)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.86Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.183 , 0.236 0.182 , 0.225	Depositor DCC
R_{free} test set	3673 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtrriage
Anisotropy	0.696	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 24.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.308 for l,-k,h	Xtrriage
Reported twinning fraction	0.320 for L,-K,H	Depositor
Outliers	0 of 75315 reflections	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	26833	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/1776	0.71	0/2412
1	D	0.27	0/1742	0.70	0/2373
1	E	0.27	0/1753	0.72	0/2383
1	H	0.27	0/1728	0.70	0/2346
1	I	0.27	0/1783	0.73	0/2427
1	L	0.28	0/1715	0.71	0/2330
1	M	0.28	0/1775	0.72	0/2412
1	P	0.28	0/1758	0.73	2/2387 (0.1%)
2	B	0.26	0/1614	0.72	0/2183
2	C	0.26	0/1511	0.71	0/2045
2	F	0.26	0/1639	0.71	2/2216 (0.1%)
2	G	0.27	0/1461	0.70	0/1971
2	J	0.27	0/1626	0.70	0/2198
2	K	0.27	0/1507	0.72	1/2040 (0.0%)
2	N	0.27	0/1633	0.73	0/2207
2	O	0.26	0/1520	0.71	0/2057
All	All	0.27	0/26541	0.71	5/35987 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	204	THR	CA-C-N	5.70	126.97	119.84
1	P	204	THR	C-N-CA	5.70	126.97	119.84
2	K	46	LEU	CB-CA-C	-5.50	109.75	117.23
2	F	198	ILE	CA-C-N	5.39	124.84	119.24
2	F	198	ILE	C-N-CA	5.39	124.84	119.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1741	21	0
1	D	1721	0	1673	17	0
1	E	1733	0	1707	22	0
1	H	1710	0	1669	15	0
1	I	1762	0	1721	21	0
1	L	1697	0	1662	24	0
1	M	1755	0	1739	30	0
1	P	1739	0	1706	20	0
2	B	1593	0	1549	24	0
2	C	1496	0	1427	23	0
2	F	1623	0	1562	30	0
2	G	1452	0	1375	22	0
2	J	1609	0	1568	33	0
2	K	1493	0	1410	20	0
2	N	1616	0	1591	33	0
2	O	1508	0	1429	26	0
3	A	44	0	26	3	0
3	D	44	0	26	1	0
3	E	44	0	26	3	0
3	H	44	0	26	2	0
3	I	44	0	26	1	0
3	L	44	0	26	3	0
3	M	44	0	26	4	0
3	P	44	0	26	1	0
4	A	24	0	0	0	0
4	B	7	0	0	0	0
4	C	10	0	0	0	0
4	D	4	0	0	0	0
4	E	25	0	0	0	0
4	F	8	0	0	0	0
4	G	4	0	0	0	0
4	H	9	0	0	0	0
4	I	14	0	0	0	0
4	J	8	0	0	0	0
4	K	7	0	0	0	0
4	L	20	0	0	1	0
4	M	38	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	N	24	0	0	0	0
4	O	6	0	0	0	0
4	P	10	0	0	0	0
All	All	26833	0	25737	342	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 342 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:158:PHE:HB2	2:K:150:ALA:HB2	1.65	0.77
2:B:158:PHE:HB2	2:C:150:ALA:HB2	1.69	0.72
1:P:132:PHE:HB2	1:P:179:LEU:HD11	1.73	0.71
2:G:130:ILE:HB	2:G:173:VAL:HG12	1.73	0.71
1:L:187:LEU:HB3	1:L:192:ILE:HB	1.73	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/261 (93%)	228 (94%)	13 (5%)	1 (0%)	30	48
1	D	242/261 (93%)	232 (96%)	10 (4%)	0	100	100
1	E	240/261 (92%)	232 (97%)	8 (3%)	0	100	100
1	H	236/261 (90%)	224 (95%)	12 (5%)	0	100	100
1	I	247/261 (95%)	233 (94%)	13 (5%)	1 (0%)	30	48
1	L	234/261 (90%)	224 (96%)	9 (4%)	1 (0%)	30	48
1	M	241/261 (92%)	230 (95%)	10 (4%)	1 (0%)	30	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	240/261 (92%)	225 (94%)	14 (6%)	1 (0%)	30	48
2	B	223/244 (91%)	215 (96%)	8 (4%)	0	100	100
2	C	208/244 (85%)	197 (95%)	11 (5%)	0	100	100
2	F	223/244 (91%)	208 (93%)	15 (7%)	0	100	100
2	G	198/244 (81%)	181 (91%)	17 (9%)	0	100	100
2	J	223/244 (91%)	211 (95%)	12 (5%)	0	100	100
2	K	212/244 (87%)	200 (94%)	11 (5%)	1 (0%)	24	42
2	N	225/244 (92%)	215 (96%)	10 (4%)	0	100	100
2	O	214/244 (88%)	207 (97%)	7 (3%)	0	100	100
All	All	3648/4040 (90%)	3462 (95%)	180 (5%)	6 (0%)	43	62

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	ALA
1	M	203	ALA
1	P	205	PRO
2	K	57	VAL
1	L	203	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/198 (90%)	179 (100%)	0	100	100
1	D	171/198 (86%)	167 (98%)	4 (2%)	44	68
1	E	176/198 (89%)	174 (99%)	2 (1%)	65	81
1	H	170/198 (86%)	170 (100%)	0	100	100
1	I	175/198 (88%)	171 (98%)	4 (2%)	44	68
1	L	170/198 (86%)	168 (99%)	2 (1%)	63	80
1	M	179/198 (90%)	174 (97%)	5 (3%)	38	62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	174/198 (88%)	167 (96%)	7 (4%)	28	53
2	B	151/193 (78%)	145 (96%)	6 (4%)	28	53
2	C	140/193 (72%)	134 (96%)	6 (4%)	26	50
2	F	154/193 (80%)	147 (96%)	7 (4%)	24	48
2	G	134/193 (69%)	125 (93%)	9 (7%)	15	31
2	J	153/193 (79%)	146 (95%)	7 (5%)	24	48
2	K	134/193 (69%)	128 (96%)	6 (4%)	24	48
2	N	155/193 (80%)	145 (94%)	10 (6%)	15	32
2	O	136/193 (70%)	128 (94%)	8 (6%)	18	37
All	All	2551/3128 (82%)	2468 (97%)	83 (3%)	33	58

5 of 83 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	255	VAL
2	O	71	GLU
2	N	60	GLU
2	N	200	LEU
2	O	237	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	212	GLN
2	O	87	ASN
2	O	78	ASN
2	O	132	ASN
1	H	153	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAD	L	301	-	46,48,48	1.20	4 (8%)	64,73,73	1.58	9 (14%)
3	NAD	H	301	-	46,48,48	1.19	4 (8%)	64,73,73	1.59	10 (15%)
3	NAD	M	301	-	46,48,48	1.18	4 (8%)	64,73,73	1.57	8 (12%)
3	NAD	D	301	-	46,48,48	1.18	4 (8%)	64,73,73	1.55	8 (12%)
3	NAD	P	301	-	46,48,48	1.19	4 (8%)	64,73,73	1.56	9 (14%)
3	NAD	A	301	-	46,48,48	1.18	4 (8%)	64,73,73	1.55	8 (12%)
3	NAD	E	301	-	46,48,48	1.19	4 (8%)	64,73,73	1.57	9 (14%)
3	NAD	I	301	-	46,48,48	1.19	4 (8%)	64,73,73	1.61	9 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	L	301	-	-	12/30/62/62	0/5/5/5
3	NAD	H	301	-	-	9/30/62/62	0/5/5/5
3	NAD	M	301	-	-	14/30/62/62	0/5/5/5
3	NAD	D	301	-	-	10/30/62/62	0/5/5/5
3	NAD	P	301	-	-	11/30/62/62	0/5/5/5
3	NAD	A	301	-	-	9/30/62/62	0/5/5/5
3	NAD	E	301	-	-	10/30/62/62	0/5/5/5
3	NAD	I	301	-	-	13/30/62/62	0/5/5/5

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	NAD	C5A-C4A	4.77	1.47	1.39
3	H	301	NAD	C5A-C4A	4.76	1.47	1.39
3	M	301	NAD	C5A-C4A	4.75	1.47	1.39
3	E	301	NAD	C5A-C4A	4.73	1.47	1.39
3	L	301	NAD	C5A-C4A	4.72	1.47	1.39

The worst 5 of 70 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	301	NAD	C5A-C4A-N3A	-6.00	118.46	126.72
3	E	301	NAD	C5A-C4A-N3A	-5.86	118.65	126.72
3	M	301	NAD	C5A-C4A-N3A	-5.86	118.65	126.72
3	L	301	NAD	C5A-C4A-N3A	-5.84	118.67	126.72
3	A	301	NAD	C5A-C4A-N3A	-5.82	118.71	126.72

There are no chirality outliers.

5 of 88 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	NAD	C5B-O5B-PA-O1A
3	A	301	NAD	C5B-O5B-PA-O3
3	A	301	NAD	O4B-C4B-C5B-O5B
3	A	301	NAD	C5D-O5D-PN-O3
3	D	301	NAD	C5B-O5B-PA-O1A


There are no ring outliers.


8 monomers are involved in 18 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	301	NAD	3	0
3	H	301	NAD	2	0
3	M	301	NAD	4	0
3	D	301	NAD	1	0
3	P	301	NAD	1	0
3	A	301	NAD	3	0
3	E	301	NAD	3	0
3	I	301	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	246/261 (94%)	-1.21	0 100 100	24, 49, 80, 123	0
1	D	246/261 (94%)	-1.26	0 100 100	26, 49, 79, 137	0
1	E	244/261 (93%)	-1.29	0 100 100	23, 45, 70, 120	0
1	H	242/261 (92%)	-1.19	0 100 100	32, 54, 86, 123	0
1	I	251/261 (96%)	-1.27	0 100 100	28, 47, 85, 158	0
1	L	240/261 (91%)	-1.25	0 100 100	28, 49, 88, 156	0
1	M	245/261 (93%)	-1.37	0 100 100	22, 43, 67, 94	0
1	P	246/261 (94%)	-1.20	0 100 100	25, 53, 89, 128	0
2	B	227/244 (93%)	-1.22	0 100 100	22, 53, 89, 114	0
2	C	216/244 (88%)	-1.10	0 100 100	29, 55, 101, 145	0
2	F	231/244 (94%)	-1.19	0 100 100	29, 54, 94, 180	0
2	G	212/244 (86%)	-1.00	0 100 100	35, 62, 100, 153	0
2	J	229/244 (93%)	-1.18	0 100 100	27, 55, 96, 136	0
2	K	220/244 (90%)	-1.05	0 100 100	32, 59, 99, 138	0
2	N	229/244 (93%)	-1.26	0 100 100	28, 51, 87, 141	0
2	O	224/244 (91%)	-1.04	0 100 100	32, 65, 95, 138	0
All	All	3748/4040 (92%)	-1.20	0 100 100	22, 52, 91, 180	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAD	H	301	44/44	0.98	0.06	45,73,108,151	0
3	NAD	L	301	44/44	0.98	0.07	76,100,128,220	0
3	NAD	E	301	44/44	0.99	0.03	5,39,58,68	0
3	NAD	A	301	44/44	0.99	0.04	29,57,80,99	0
3	NAD	I	301	44/44	0.99	0.04	7,42,61,87	0
3	NAD	D	301	44/44	0.99	0.04	19,49,82,93	0
3	NAD	M	301	44/44	0.99	0.04	1,45,58,69	0
3	NAD	P	301	44/44	0.99	0.05	42,74,93,113	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAD H 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD L 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD E 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD I 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD M 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD P 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.