



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:46 PM UTC

PDB ID : 4CQO / pdb_00004cqo
Title : Structure of the human CNOT1 superfamily homology domain in complex with a Nanos1 peptide
Authors : Raisch, T.; Jonas, S.; Weichenrieder, O.; Bhandari, D.; Izaurralde, E.
Deposited on : 2014-02-21
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	535	8% 79% 16% 5%
1	C	535	6% 85% 11% ...
2	B	17	6% 59% 29% 12%
2	D	17	24% 65% 18% 18%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	510	Total	C	N	O	S	0	4	0
			4143	2685	715	718	25			
1	C	516	Total	C	N	O	S	0	2	0
			4189	2710	721	734	24			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1827	GLY	-	expression tag	UNP A5YKK6
A	1828	PRO	-	expression tag	UNP A5YKK6
A	1829	HIS	-	expression tag	UNP A5YKK6
A	1830	MET	-	expression tag	UNP A5YKK6
A	1831	LEU	-	expression tag	UNP A5YKK6
A	1832	GLU	-	expression tag	UNP A5YKK6
C	1827	GLY	-	expression tag	UNP A5YKK6
C	1828	PRO	-	expression tag	UNP A5YKK6
C	1829	HIS	-	expression tag	UNP A5YKK6
C	1830	MET	-	expression tag	UNP A5YKK6
C	1831	LEU	-	expression tag	UNP A5YKK6
C	1832	GLU	-	expression tag	UNP A5YKK6

- Molecule 2 is a protein called NANOS HOMOLOG 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	15	Total	C	N	O	0	0	0
			120	80	17	23			
2	D	14	Total	C	N	O	0	0	0
			110	73	16	21			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	30	Total 30	O 30	0	0
3	C	32	Total 32	O 32	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 1



• Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 1




• Molecule 2: NANOS HOMOLOG 1





● Molecule 2: NANOS HOMOLOG 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.53Å 167.33Å 112.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.45 – 2.80 48.45 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.45-2.80) 99.5 (48.45-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.81Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.3_1479)	Depositor
R, R_{free}	0.222 , 0.243 0.230 , 0.249	Depositor DCC
R_{free} test set	2303 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.925	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8624	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.97 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3838e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/4261	0.70	0/5792
1	C	0.30	0/4303	0.70	2/5853 (0.0%)
2	B	0.23	0/123	0.66	0/168
2	D	0.23	0/113	0.65	0/154
All	All	0.30	0/8800	0.70	2/11967 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2122	PRO	CA-C-N	6.07	125.75	119.56
1	C	2122	PRO	C-N-CA	6.07	125.75	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4143	0	4167	47	0
1	C	4189	0	4194	31	0
2	B	120	0	113	5	0
2	D	110	0	97	2	0
3	A	30	0	0	2	0
3	C	32	0	0	0	0
All	All	8624	0	8571	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1960:ILE:HD11	1:A:2011:ILE:HB	1.81	0.62
1:C:1960:ILE:HD11	1:C:2011:ILE:HB	1.81	0.62
1:C:2195:ASP:O	1:C:2199:ASN:ND2	2.32	0.61
1:A:2121:ILE:O	1:A:2129[B]:ARG:NH1	2.35	0.60
1:C:2193:LEU:O	1:C:2196:LEU:HB2	2.03	0.58
1:A:2195:ASP:O	1:A:2199:ASN:ND2	2.30	0.58
1:A:2195:ASP:HA	1:A:2198:SER:HB2	1.86	0.57
1:A:2145:PRO:HG2	1:A:2323:LEU:HD23	1.86	0.57
1:C:1896:THR:OG1	1:C:1965:LYS:NZ	2.29	0.56
1:C:1854:TYR:OH	1:C:1882:GLN:NE2	2.37	0.55
1:A:2193:LEU:O	1:A:2196:LEU:HB2	2.06	0.55
1:C:2354:GLN:HG3	2:D:49:LEU:HD21	1.92	0.52
1:C:2145:PRO:HG2	1:C:2323:LEU:HD23	1.91	0.51
1:C:1883:MET:HE2	1:C:1889:LEU:HD21	1.92	0.51
1:A:1998:LEU:O	1:A:2002:ASN:ND2	2.32	0.50
1:A:2158:ILE:HD12	1:A:2282:HIS:HB2	1.94	0.50
1:C:1859:TRP:NE1	1:C:1942:LEU:HB2	2.28	0.49
1:A:1854:TYR:OH	1:A:1882:GLN:NE2	2.42	0.49
1:A:1859:TRP:NE1	1:A:1942:LEU:HB2	2.28	0.49
1:A:1871:ASP:HA	1:A:1874:LYS:HE2	1.95	0.48
1:A:2193:LEU:HB3	1:A:2253:ASN:CG	2.37	0.48
1:A:1900:ARG:NE	3:A:3005:HOH:O	2.35	0.48
1:A:1893:ASP:O	1:A:1897:ARG:HG3	2.14	0.48
1:C:1843:ASP:HB3	1:C:1897:ARG:NH2	2.30	0.47
1:A:2263:ARG:HD3	1:A:2300:ILE:HD13	1.97	0.47
1:A:2348:GLU:HA	1:A:2351:LYS:HE2	1.97	0.47
1:A:2354:GLN:HG3	2:B:49:LEU:HD21	1.97	0.46
1:A:2036:TYR:OH	1:A:2087:PRO:HB3	2.15	0.46
1:C:2193:LEU:HB3	1:C:2253:ASN:CG	2.39	0.46
1:A:1843:ASP:HB2	1:A:1897:ARG:HH22	1.79	0.46
1:C:1955:ASN:ND2	1:C:1958:THR:OG1	2.49	0.46
1:A:2024:HIS:CD2	1:A:2027:ARG:HH21	2.34	0.46
1:A:1948:LYS:O	1:A:1959:LYS:NZ	2.46	0.46
1:C:2056:PRO:C	1:C:2058:GLN:H	2.25	0.45
1:A:1939:PHE:CE2	1:A:1943:ILE:HD11	2.51	0.45
2:B:43:TRP:CE2	2:B:47:LEU:HD11	2.52	0.45
1:A:2238:SER:OG	1:A:2239:MET:N	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2140:MET:HE3	1:A:2157:GLU:HG3	1.99	0.45
1:C:2122:PRO:HA	1:C:2123:PRO:HD3	1.75	0.45
1:C:2326:PHE:O	1:C:2330:ILE:HG12	2.17	0.45
1:C:1917:GLN:HB2	1:C:1928:ILE:HG13	1.99	0.45
1:C:2346:ALA:HA	1:C:2347:PRO:HD3	1.86	0.45
1:A:2099:LEU:HD12	1:A:2128:LEU:HD22	2.00	0.44
1:C:2121:ILE:O	1:C:2129[B]:ARG:NH1	2.50	0.44
1:A:2017:THR:HA	1:A:2063:MET:HE1	2.00	0.44
2:B:52:LEU:HD12	1:C:2040:GLU:HG2	2.00	0.44
1:A:2331:LYS:HG3	2:B:43:TRP:CD1	2.52	0.44
1:A:2169:THR:HA	1:A:2172:MET:HE2	2.00	0.44
1:C:2158:ILE:HD12	1:C:2282:HIS:HB2	2.00	0.44
1:A:2104:HIS:HD2	1:C:2317:PRO:HB3	1.82	0.43
1:A:1947:VAL:HA	1:A:1962:LEU:HD23	2.00	0.42
1:A:2309:LEU:O	1:A:2313:ILE:HG13	2.19	0.42
1:A:2290:TYR:CZ	1:A:2294:GLU:HG3	2.53	0.42
2:D:43:TRP:CE2	2:D:47:LEU:HD11	2.54	0.42
1:A:1933:TYR:CE1	1:A:1992:ARG:HG3	2.55	0.42
1:A:2130:ASN:ND2	3:A:3013:HOH:O	2.51	0.42
1:C:1987:GLN:NE2	1:C:2037:ALA:H	2.17	0.42
1:C:1853:GLU:HA	1:C:1905:MET:HE1	2.02	0.42
1:C:2270:ILE:HG23	1:C:2284:PHE:HD1	1.83	0.42
1:C:1844:ASP:CG	1:C:1900:ARG:HH22	2.29	0.41
1:A:1881:GLY:O	1:A:1885[B]:GLN:HG2	2.20	0.41
1:A:2326:PHE:O	1:A:2330:ILE:HG12	2.21	0.41
1:C:1871:ASP:HA	1:C:1874:LYS:HG2	2.02	0.41
1:C:2239:MET:HE1	1:C:2289:LEU:HB2	2.01	0.41
1:A:2332:ASN:HB3	1:A:2335:PHE:HD2	1.85	0.41
1:C:1982:GLN:HB3	1:C:1983:SER:H	1.69	0.41
1:A:2050:ARG:HA	1:A:2050:ARG:HD2	1.93	0.41
1:C:1893:ASP:O	1:C:1897:ARG:HG3	2.20	0.41
1:C:2088:MET:HE3	1:C:2088:MET:HB3	1.96	0.41
1:A:2152:VAL:O	1:A:2158:ILE:HG13	2.21	0.41
1:A:2361:MET:HE1	2:B:46:TYR:HE2	1.86	0.41
1:A:2270:ILE:HG23	1:A:2284:PHE:HD1	1.85	0.40
1:A:2086:LYS:HB2	1:A:2087:PRO:HD3	2.03	0.40
1:A:2184:LEU:HD22	1:A:2247:HIS:CD2	2.56	0.40
1:A:2239:MET:HE1	1:A:2289:LEU:HB2	2.02	0.40
1:C:2212:LEU:HD22	1:C:2216:ASN:HD21	1.86	0.40
1:A:1991:HIS:O	1:A:1995:ILE:HG22	2.21	0.40
1:A:2122:PRO:HB2	1:A:2124:ASN:OD1	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2266:PHE:CZ	1:A:2270:ILE:HD11	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/535 (95%)	499 (98%)	9 (2%)	0	100	100
1	C	514/535 (96%)	502 (98%)	12 (2%)	0	100	100
2	B	13/17 (76%)	12 (92%)	1 (8%)	0	100	100
2	D	12/17 (71%)	12 (100%)	0	0	100	100
All	All	1047/1104 (95%)	1025 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/478 (95%)	448 (98%)	7 (2%)	57	84
1	C	461/478 (96%)	454 (98%)	7 (2%)	57	84
2	B	13/14 (93%)	13 (100%)	0	100	100
2	D	11/14 (79%)	11 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	940/984 (96%)	926 (98%)	14 (2%)	57 84

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1858	GLU
1	A	1987	GLN
1	A	2046	ILE
1	A	2057	GLN
1	A	2111	CYS
1	A	2212	LEU
1	A	2309	LEU
1	C	1858	GLU
1	C	1888	ILE
1	C	2046	ILE
1	C	2057	GLN
1	C	2158	ILE
1	C	2212	LEU
1	C	2309	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1949	HIS
1	A	1987	GLN
1	A	1991	HIS
1	A	2104	HIS
1	A	2213	GLN
1	A	2229	HIS
1	A	2252	GLN
1	A	2282	HIS
1	C	1884	HIS
1	C	1955	ASN
1	C	1982	GLN
1	C	1986	GLN
1	C	1987	GLN
1	C	2012	ASN
1	C	2216	ASN
1	C	2244	HIS
1	C	2273	GLN
1	C	2282	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	510/535 (95%)	0.64	45 (8%)	15 11	25, 56, 102, 146	4 (0%)
1	C	516/535 (96%)	0.31	34 (6%)	24 18	16, 47, 98, 145	2 (0%)
2	B	15/17 (88%)	0.25	1 (6%)	24 17	41, 55, 77, 82	0
2	D	14/17 (82%)	0.77	4 (28%)	1 1	45, 64, 119, 125	0
All	All	1055/1104 (95%)	0.47	84 (7%)	18 13	16, 52, 102, 146	6 (0%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2346	ALA	6.8
1	A	1925	PRO	5.4
1	C	2346	ALA	5.4
1	A	2088	MET	4.9
1	C	1842	TYR	4.8
1	A	1928	ILE	4.8
1	C	1928	ILE	4.6
1	C	1920	ASN	4.3
1	C	1919	HIS	4.3
1	C	2056	PRO	4.3
1	A	2085	THR	4.0
1	A	2082	VAL	4.0
1	A	1842	TYR	3.9
1	C	1867	ALA	3.8
1	C	1926	THR	3.8
1	C	1918	GLN	3.8
1	A	1952	GLU	3.7
1	C	2193	LEU	3.7
1	A	2056	PRO	3.7
1	C	1927	MET	3.7
1	A	1926	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2194	SER	3.6
1	C	1988	LEU	3.5
1	A	2080[A]	ARG	3.4
1	A	1873	THR	3.4
2	D	52	LEU	3.4
1	A	2081	ASN	3.3
1	A	2055	THR	3.3
1	A	1916	GLU	3.2
1	A	2148	PRO	3.2
1	C	2194	SER	3.1
1	A	1927	MET	3.1
2	D	53	ILE	3.1
1	C	1953	ALA	3.1
1	C	2057	GLN	3.1
1	C	2198	SER	3.1
1	A	1953	ALA	3.0
1	C	2148	PRO	3.0
1	C	2055	THR	3.0
1	A	1867	ALA	3.0
1	C	1921	PRO	2.9
2	D	51	THR	2.9
1	C	1874	LYS	2.9
1	A	1892	ASP	2.9
1	A	2193	LEU	2.8
1	A	2087	PRO	2.8
1	C	1932	CYS	2.8
1	A	2296	ASN	2.8
1	A	2241	THR	2.6
1	A	2129[A]	ARG	2.6
1	A	2191	THR	2.6
1	A	1885[A]	GLN	2.6
1	A	2149	ASN	2.5
1	C	1873	THR	2.5
1	C	1870	ARG	2.5
1	C	2361	MET	2.5
1	C	2347	PRO	2.4
1	A	1891	THR	2.4
1	A	2120	VAL	2.4
1	A	2151	LYS	2.4
1	A	2190	VAL	2.3
1	A	1951	GLY	2.3
1	C	2297	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2163	ARG	2.3
2	B	54	THR	2.3
1	A	2086	LYS	2.3
1	A	2054	HIS	2.3
1	C	1933	TYR	2.2
1	C	2296	ASN	2.2
1	A	2153	ASP	2.2
1	C	2241	THR	2.2
2	D	49	LEU	2.2
1	A	1888	ILE	2.1
1	A	1874	LYS	2.1
1	A	1868	ALA	2.1
1	C	1929	ARG	2.1
1	C	2129[A]	ARG	2.1
1	A	2198	SER	2.1
1	A	1872	SER	2.0
1	C	1917	GLN	2.0
1	A	1929	ARG	2.0
1	A	2345	CYS	2.0
1	C	1888	ILE	2.0
1	C	1915	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.