



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:04 AM UTC

PDB ID : 4CQX / pdb_00004cqx
Title : H5 (tyTy) Del133/Ile155Thr Mutant Haemagglutinin in Complex with Human Receptor Analogue 6'SLN
Authors : Xiong, X.; Xiao, H.; Martin, S.R.; Coombs, P.J.; Liu, J.; Collins, P.J.; Vachieri, S.G.; Walker, P.A.; Lin, Y.P.; McCauley, J.W.; Gamblin, S.J.; Skehel, J.J.
Deposited on : 2014-02-21
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	% 94% ...
1	C	327	% 94% ..
1	E	327	% 94% ...
2	B	166	% 89% 9% .
2	D	166	13% 90% 8% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	166	
3	G	3	
3	H	3	
3	I	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	1322	X	-	-	-
4	NAG	E	1322	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12610 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HAEMAGGLUTININ HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2552	1607	443	488	14			
1	C	320	Total	C	N	O	S	0	0	0
			2537	1598	441	484	14			
1	E	322	Total	C	N	O	S	0	0	0
			2549	1606	443	486	14			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ASP	-	expression tag	UNP Q207Z6
A	0	PRO	-	expression tag	UNP Q207Z6
A	?	-	ALA	deletion	UNP Q207Z6
A	150	THR	ILE	engineered mutation	UNP Q207Z6
A	322	ARG	GLY	conflict	UNP Q207Z6
A	324	THR	ARG	conflict	UNP Q207Z6
C	-1	ASP	-	expression tag	UNP Q207Z6
C	0	PRO	-	expression tag	UNP Q207Z6
C	?	-	ALA	deletion	UNP Q207Z6
C	150	THR	ILE	engineered mutation	UNP Q207Z6
C	322	ARG	GLY	conflict	UNP Q207Z6
C	324	THR	ARG	conflict	UNP Q207Z6
E	-1	ASP	-	expression tag	UNP Q207Z6
E	0	PRO	-	expression tag	UNP Q207Z6
E	?	-	ALA	deletion	UNP Q207Z6
E	150	THR	ILE	engineered mutation	UNP Q207Z6
E	322	ARG	GLY	conflict	UNP Q207Z6
E	324	THR	ARG	conflict	UNP Q207Z6

- Molecule 2 is a protein called HAEMAGGLUTININ HA2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	163	Total	C	N	O	S	0	0	0
			1320	822	229	261	8			
2	D	163	Total	C	N	O	S	0	0	0
			1298	808	228	254	8			
2	F	163	Total	C	N	O	S	0	0	0
			1306	815	226	257	8			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	3	Total	C	N	O	0	0	0
			46	25	2	19			
3	H	3	Total	C	N	O	0	0	0
			46	25	2	19			
3	I	3	Total	C	N	O	0	0	0
			46	25	2	19			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	C	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	P	0	0
			5	4	1		
5	E	1	Total	O	P	0	0
			5	4	1		



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	182	Total	O	0	0
			182	182		
6	B	84	Total	O	0	0
			84	84		
6	C	195	Total	O	0	0
			195	195		
6	D	45	Total	O	0	0
			45	45		
6	E	217	Total	O	0	0
			217	217		
6	F	36	Total	O	0	0
			36	36		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: HAEMAGGLUTININ HA1




- Molecule 1: HAEMAGGLUTININ HA1



- Molecule 1: HAEMAGGLUTININ HA1




- Molecule 2: HAEMAGGLUTININ HA2



- Molecule 2: HAEMAGGLUTININ HA2



- Molecule 2: HAEMAGGLUTININ HA2



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.82Å 117.10Å 101.28Å 90.00° 92.58° 90.00°	Depositor
Resolution (Å)	101.17 – 2.30 101.17 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (101.17-2.30) 99.5 (101.17-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0046	Depositor
R, R_{free}	0.186 , 0.226 (Not available) , 0.218	Depositor DCC
R_{free} test set	4539 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.658	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.030 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12610	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NAG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2614	0.70	0/3554
1	C	0.51	0/2598	0.69	0/3531
1	E	0.55	0/2611	0.73	0/3550
2	B	0.52	0/1347	0.73	0/1813
2	D	0.49	0/1324	0.73	0/1784
2	F	0.50	0/1333	0.77	0/1796
All	All	0.51	0/11827	0.72	0/16028

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2552	0	2483	11	0
1	C	2537	0	2473	10	0
1	E	2549	0	2480	10	0
2	B	1320	0	1224	13	0
2	D	1298	0	1196	9	0
2	F	1306	0	1205	8	0
3	G	46	0	40	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	46	0	40	0	0
3	I	46	0	40	1	0
4	A	42	0	39	4	0
4	C	28	0	26	0	0
4	E	56	0	52	0	0
5	A	10	0	0	0	0
5	C	10	0	0	0	0
5	E	5	0	0	1	0
6	A	182	0	0	0	0
6	B	84	0	0	0	0
6	C	195	0	0	1	0
6	D	45	0	0	1	0
6	E	217	0	0	0	0
6	F	36	0	0	1	0
All	All	12610	0	11298	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:ILE:HD11	2:B:122:VAL:HG21	1.63	0.78
1:E:279:THR:HG22	1:E:281:ILE:H	1.48	0.78
1:E:279:THR:HB	1:E:282:GLY:O	1.87	0.73
2:B:156:THR:O	2:B:156:THR:HG22	1.90	0.72
1:A:20:MET:HE1	2:D:110:PHE:CD2	2.34	0.62
1:C:20:MET:HE3	6:F:2010:HOH:O	2.02	0.59
2:B:110:PHE:CD2	1:E:20:MET:HE1	2.38	0.58
1:A:303:LYS:HE2	2:B:60:ASN:HA	1.85	0.58
4:A:1322:NAG:H3	4:A:1322:NAG:C8	2.34	0.58
2:B:124:LEU:HD22	2:F:134:GLY:HA2	1.87	0.57
1:C:19:ILE:HD12	2:D:101:LEU:HB3	1.87	0.56
2:D:60:ASN:HB3	6:D:2013:HOH:O	2.06	0.55
1:C:20:MET:HE2	2:F:51:LYS:HE3	1.90	0.54
4:A:1322:NAG:H3	4:A:1322:NAG:H83	1.91	0.53
2:B:156:THR:O	2:B:156:THR:CG2	2.60	0.49
2:D:84:MET:HE2	2:D:85:GLU:HG2	1.95	0.49
1:C:166:THR:HG22	6:C:2125:HOH:O	2.14	0.47
1:A:273:CYS:SG	1:A:274:ASN:N	2.87	0.47
2:B:51:LYS:HE3	1:E:20:MET:HE3	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:1:NAG:H1	3:I:1:NAG:H82	1.96	0.47
1:E:5:ILE:HG13	2:F:119:TYR:HA	1.98	0.46
1:A:81:GLU:O	1:A:265:LYS:HA	2.16	0.45
1:E:5:ILE:HD11	2:F:122:VAL:HG21	1.97	0.45
1:C:121:SER:CB	1:C:161:ARG:HH22	2.29	0.45
1:C:196:TYR:OH	1:C:241:GLU:OE2	2.31	0.45
1:E:19:ILE:HD12	2:F:101:LEU:HB3	1.99	0.45
1:C:234:PRO:O	1:C:235:ASN:HB2	2.16	0.45
2:B:72:ASN:OD1	2:B:75:ARG:NH2	2.50	0.45
1:C:279:THR:HG22	1:C:297:THR:HG22	1.99	0.45
2:B:102:MET:HE3	2:F:102:MET:HE1	1.99	0.44
1:A:20:MET:HE3	2:D:51:LYS:HE3	1.99	0.44
2:D:97:GLU:HB3	2:F:58:LYS:HE3	2.00	0.43
4:A:1322:NAG:C8	4:A:1322:NAG:C3	2.96	0.43
1:C:121:SER:OG	1:C:161:ARG:NH2	2.41	0.43
1:E:35:LYS:N	5:E:1328:PO4:O2	2.41	0.43
1:A:280:PRO:HD3	1:A:296:LEU:O	2.19	0.42
1:C:5:ILE:HD11	2:D:122:VAL:HG21	2.01	0.42
2:B:119:TYR:OH	2:B:132:GLU:HG3	2.19	0.42
1:E:90:CYS:HB2	1:E:133:ALA:O	2.19	0.42
1:A:279:THR:HG22	1:A:297:THR:HG22	2.02	0.41
4:A:1322:NAG:H83	4:A:1322:NAG:C3	2.49	0.41
1:A:19:ILE:HD12	2:B:101:LEU:HB3	2.01	0.41
2:D:98:LEU:HG	2:D:102:MET:HE2	2.03	0.41
2:B:102:MET:CE	2:F:102:MET:HE1	2.50	0.41
1:E:83:ILE:HD12	1:E:267:GLU:OE2	2.20	0.41
1:A:20:MET:HE1	2:D:110:PHE:CE2	2.57	0.40
1:A:4:CYS:O	2:B:24:TYR:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/327 (98%)	310 (97%)	10 (3%)	0	100	100
1	C	318/327 (97%)	307 (96%)	11 (4%)	0	100	100
1	E	320/327 (98%)	311 (97%)	9 (3%)	0	100	100
2	B	161/166 (97%)	159 (99%)	2 (1%)	0	100	100
2	D	161/166 (97%)	157 (98%)	4 (2%)	0	100	100
2	F	161/166 (97%)	157 (98%)	4 (2%)	0	100	100
All	All	1441/1479 (97%)	1401 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/294 (98%)	283 (98%)	6 (2%)	47	66
1	C	287/294 (98%)	284 (99%)	3 (1%)	68	82
1	E	288/294 (98%)	281 (98%)	7 (2%)	43	62
2	B	139/141 (99%)	137 (99%)	2 (1%)	59	76
2	D	134/141 (95%)	131 (98%)	3 (2%)	45	65
2	F	136/141 (96%)	132 (97%)	4 (3%)	37	55
All	All	1273/1305 (98%)	1248 (98%)	25 (2%)	48	67

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ILE
1	A	89	LEU
1	A	107	ARG
1	A	144	ARG
1	A	273	CYS
1	A	279	THR
2	B	29	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	30	GLN
1	C	4	CYS
1	C	19	ILE
1	C	316	LEU
2	D	30	GLN
2	D	61	THR
2	D	137	CYS
1	E	19	ILE
1	E	174	LEU
1	E	278	GLN
1	E	279	THR
1	E	281	ILE
1	E	296	LEU
1	E	316	LEU
2	F	60	ASN
2	F	89	LEU
2	F	133	LEU
2	F	156	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	94	ASN
1	A	96	ASN
1	A	221	GLN
2	B	25	HIS
2	B	42	GLN
2	B	60	ASN
2	B	154	ASN
1	C	2	GLN
1	C	72	ASN
1	C	110	HIS
1	C	137	GLN
1	C	191	GLN
2	D	42	GLN
1	E	72	ASN
1	E	103	HIS
1	E	191	GLN
1	E	221	GLN
1	E	235	ASN
2	F	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	62	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	G	1	3	15,15,15	0.40	0	21,21,21	1.34	3 (14%)
3	GAL	G	2	3	11,11,12	0.41	0	15,15,17	0.86	1 (6%)
3	SIA	G	3	3	20,20,21	0.59	0	21,28,31	1.22	3 (14%)
3	NAG	H	1	3	15,15,15	0.51	0	21,21,21	1.41	5 (23%)
3	GAL	H	2	3	11,11,12	0.41	0	15,15,17	0.87	1 (6%)
3	SIA	H	3	3	20,20,21	0.62	0	21,28,31	1.25	2 (9%)
3	NAG	I	1	3	15,15,15	0.65	0	21,21,21	1.97	6 (28%)
3	GAL	I	2	3	11,11,12	0.46	0	15,15,17	0.67	0
3	SIA	I	3	3	20,20,21	0.56	0	21,28,31	1.26	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3	-	0/6/26/26	0/1/1/1
3	GAL	G	2	3	-	0/2/19/22	0/1/1/1
3	SIA	G	3	3	-	0/18/34/38	0/1/1/1
3	NAG	H	1	3	-	4/6/26/26	0/1/1/1
3	GAL	H	2	3	-	0/2/19/22	0/1/1/1
3	SIA	H	3	3	-	0/18/34/38	0/1/1/1
3	NAG	I	1	3	-	6/6/26/26	0/1/1/1
3	GAL	I	2	3	-	0/2/19/22	0/1/1/1
3	SIA	I	3	3	-	0/18/34/38	0/1/1/1

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	1	NAG	C1-C2-N2	-4.55	105.46	110.73
3	H	1	NAG	C1-C2-N2	-4.07	106.02	110.73
3	I	1	NAG	C2-N2-C7	3.99	132.44	123.11
3	I	1	NAG	C4-C3-C2	3.76	115.87	110.40
3	I	1	NAG	C8-C7-N2	3.53	121.96	116.12
3	H	3	SIA	C4-C5-N5	-3.49	103.56	110.44
3	I	1	NAG	O5-C1-C2	-3.27	106.23	109.52
3	I	1	NAG	C3-C4-C5	3.27	116.15	110.23
3	I	3	SIA	O1B-C1-C2	2.78	119.93	112.71
3	G	3	SIA	C4-C5-N5	-2.61	105.29	110.44
3	G	3	SIA	O1B-C1-C2	2.58	119.41	112.71
3	I	1	NAG	O5-C5-C4	2.55	114.29	109.70
3	H	3	SIA	O1B-C1-C2	2.44	119.05	112.71
3	G	1	NAG	C3-C2-N2	-2.31	106.37	110.62
3	I	3	SIA	C4-C5-N5	-2.30	105.90	110.44
3	G	2	GAL	C1-C2-C3	2.28	112.96	109.64
3	H	2	GAL	C1-C2-C3	2.26	112.93	109.64
3	H	1	NAG	C3-C2-N2	-2.26	106.46	110.62
3	H	1	NAG	C1-C2-C3	2.20	113.55	110.54
3	I	3	SIA	O1A-C1-C2	-2.17	118.16	122.85
3	H	1	NAG	C4-C3-C2	2.14	113.52	110.40
3	G	3	SIA	O6-C2-C1	2.12	111.73	107.72
3	H	1	NAG	O5-C5-C6	2.04	111.48	106.44
3	G	1	NAG	C1-C2-C3	2.01	113.28	110.54

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	H	1	NAG	C1-C2-N2-C7
3	H	1	NAG	O5-C5-C6-O6
3	H	1	NAG	C4-C5-C6-O6
3	I	1	NAG	C8-C7-N2-C2
3	I	1	NAG	O7-C7-N2-C2
3	I	1	NAG	O5-C5-C6-O6
3	I	1	NAG	C1-C2-N2-C7
3	I	1	NAG	C4-C5-C6-O6
3	I	1	NAG	C3-C2-N2-C7
3	H	1	NAG	C3-C2-N2-C7

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



Oligosaccharide Chain H



Oligosaccharide Chain I



5.6 Ligand geometry

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PO4	A	1327	-	4,4,4	0.86	0	6,6,6	0.66	0
4	NAG	C	1322	1	14,14,15	0.50	0	17,19,21	1.28	2 (11%)
5	PO4	C	1326	-	4,4,4	0.77	0	6,6,6	0.51	0
5	PO4	C	1327	-	4,4,4	1.02	0	6,6,6	0.65	0
4	NAG	E	1321	1	14,14,15	0.57	0	17,19,21	1.29	2 (11%)
4	NAG	E	1324	1	14,14,15	0.46	0	17,19,21	1.53	3 (17%)
5	PO4	A	1328	-	4,4,4	0.93	0	6,6,6	0.55	0
4	NAG	C	1321	1	14,14,15	0.80	0	17,19,21	1.46	4 (23%)
4	NAG	A	1322	1	14,14,15	0.92	1 (7%)	17,19,21	2.49	6 (35%)
4	NAG	A	1321	1	14,14,15	0.47	0	17,19,21	1.24	3 (17%)
4	NAG	E	1323	1	14,14,15	0.57	0	17,19,21	1.18	2 (11%)
4	NAG	E	1322	1	14,14,15	0.56	0	17,19,21	1.70	5 (29%)
5	PO4	E	1328	-	4,4,4	0.98	0	6,6,6	0.52	0
4	NAG	A	1323	1	14,14,15	0.34	0	17,19,21	2.13	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1322	1	-	0/6/23/26	0/1/1/1
4	NAG	E	1321	1	-	2/6/23/26	0/1/1/1
4	NAG	E	1324	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1321	1	-	3/6/23/26	0/1/1/1
4	NAG	A	1322	1	1/1/5/7	5/6/23/26	0/1/1/1
4	NAG	A	1321	1	-	4/6/23/26	0/1/1/1
4	NAG	E	1323	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	1322	1	1/1/5/7	2/6/23/26	0/1/1/1
4	NAG	A	1323	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1322	NAG	C1-C2	2.64	1.55	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1323	NAG	C1-O5-C5	7.53	122.28	112.19
4	A	1322	NAG	C2-N2-C7	5.79	130.65	122.90
4	A	1322	NAG	C8-C7-N2	4.75	124.00	116.12
4	E	1324	NAG	C1-O5-C5	4.09	117.67	112.19
4	A	1322	NAG	C1-C2-N2	3.59	116.09	110.43
4	C	1322	NAG	C1-O5-C5	3.44	116.79	112.19
4	C	1321	NAG	C2-N2-C7	3.35	127.39	122.90
4	A	1322	NAG	O7-C7-C8	-3.35	116.09	122.05
4	E	1321	NAG	C1-O5-C5	3.28	116.58	112.19
4	E	1322	NAG	O5-C1-C2	-3.27	106.23	111.29
4	E	1324	NAG	O5-C1-C2	3.03	115.98	111.29
4	A	1322	NAG	C1-O5-C5	2.97	116.17	112.19
4	E	1322	NAG	C4-C3-C2	-2.70	107.06	111.02
4	C	1321	NAG	O7-C7-C8	-2.64	117.36	122.05
4	E	1322	NAG	C8-C7-N2	2.54	120.33	116.12
4	C	1321	NAG	C4-C3-C2	2.52	114.71	111.02
4	E	1322	NAG	C1-C2-N2	2.50	114.37	110.43
4	A	1322	NAG	O5-C1-C2	-2.46	107.48	111.29
4	E	1323	NAG	C1-O5-C5	2.44	115.45	112.19
4	C	1322	NAG	C3-C4-C5	-2.35	105.97	110.23
4	A	1321	NAG	O7-C7-C8	-2.27	118.02	122.05
4	E	1323	NAG	O5-C1-C2	-2.26	107.80	111.29
4	A	1323	NAG	O5-C5-C4	2.22	116.23	110.83
4	A	1323	NAG	C6-C5-C4	-2.19	107.65	113.02
4	A	1321	NAG	O5-C1-C2	-2.15	107.97	111.29
4	A	1321	NAG	C8-C7-N2	2.14	119.67	116.12
4	E	1321	NAG	O5-C5-C6	2.13	111.80	107.66
4	E	1324	NAG	C3-C4-C5	-2.11	106.41	110.23
4	E	1322	NAG	C1-O5-C5	2.09	114.98	112.19
4	C	1321	NAG	O7-C7-N2	2.00	125.52	121.98

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	1322	NAG	C1
4	E	1322	NAG	C1

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1322	NAG	C3-C2-N2-C7
4	C	1321	NAG	C3-C2-N2-C7
4	A	1322	NAG	O5-C5-C6-O6
4	C	1321	NAG	O5-C5-C6-O6
4	A	1322	NAG	C4-C5-C6-O6
4	C	1321	NAG	C4-C5-C6-O6
4	E	1321	NAG	O5-C5-C6-O6
4	A	1321	NAG	C8-C7-N2-C2
4	A	1321	NAG	O7-C7-N2-C2
4	A	1322	NAG	C8-C7-N2-C2
4	A	1322	NAG	O7-C7-N2-C2
4	E	1322	NAG	C8-C7-N2-C2
4	E	1322	NAG	O7-C7-N2-C2
4	A	1321	NAG	C4-C5-C6-O6
4	E	1321	NAG	C4-C5-C6-O6
4	A	1321	NAG	O5-C5-C6-O6
4	E	1323	NAG	C4-C5-C6-O6
4	E	1323	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1322	NAG	4	0
5	E	1328	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/327 (98%)	-0.14	3 (0%) 81 82	28, 47, 65, 82	0
1	C	320/327 (97%)	-0.21	3 (0%) 81 82	29, 43, 63, 112	0
1	E	322/327 (98%)	-0.29	4 (1%) 76 77	27, 39, 55, 102	0
2	B	163/166 (98%)	-0.07	1 (0%) 85 86	25, 44, 71, 83	0
2	D	163/166 (98%)	0.69	22 (13%) 7 8	28, 64, 149, 167	0
2	F	163/166 (98%)	0.54	22 (13%) 7 8	25, 57, 156, 176	0
All	All	1453/1479 (98%)	-0.01	55 (3%) 44 46	25, 44, 109, 176	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	-1	ASP	4.7
2	F	143	ARG	4.6
2	D	29	GLU	4.0
2	D	163	SER	3.8
2	F	163	SER	3.5
1	E	273	CYS	3.3
2	F	138	PHE	3.2
2	D	61	THR	3.1
1	A	320	PRO	3.1
2	F	153	ARG	3.0
2	D	141	TYR	2.8
2	F	137	CYS	2.8
2	D	144	CYS	2.8
2	D	134	GLY	2.8
2	D	152	VAL	2.8
2	F	18	VAL	2.8
1	A	273	CYS	2.7
1	E	320	PRO	2.7
1	E	0	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	138	PHE	2.5
2	F	27	SER	2.5
2	D	124	LEU	2.5
2	D	156	THR	2.5
2	F	61	THR	2.4
2	F	124	LEU	2.4
1	C	1	ASP	2.4
2	F	29	GLU	2.4
2	F	156	THR	2.4
2	D	27	SER	2.4
2	F	32	SER	2.4
2	F	160	PRO	2.3
1	A	214	THR	2.3
2	D	33	GLY	2.3
2	B	163	SER	2.3
2	D	133	LEU	2.3
2	D	38	LYS	2.2
2	D	137	CYS	2.2
2	D	143	ARG	2.1
2	D	159	TYR	2.1
1	C	3	ILE	2.1
2	F	152	VAL	2.1
2	D	31	GLY	2.1
2	F	134	GLY	2.1
2	F	141	TYR	2.1
1	C	320	PRO	2.1
2	D	140	PHE	2.1
2	D	160	PRO	2.1
2	F	140	PHE	2.1
2	F	130	ALA	2.1
2	D	157	TYR	2.1
2	D	158	ASP	2.1
2	F	60	ASN	2.1
2	F	31	GLY	2.1
2	F	155	GLY	2.1
2	F	35	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	I	1	15/15	0.60	0.19	99,107,111,111	0
3	NAG	G	1	15/15	0.69	0.17	100,109,117,118	0
3	NAG	H	1	15/15	0.70	0.17	88,98,108,108	0
3	GAL	G	2	11/12	0.82	0.12	58,81,91,92	0
3	GAL	H	2	11/12	0.86	0.12	52,75,77,79	0
3	GAL	I	2	11/12	0.89	0.10	52,76,84,88	0
3	SIA	G	3	20/21	0.95	0.07	44,48,53,53	0
3	SIA	I	3	20/21	0.95	0.07	35,39,48,50	0
3	SIA	H	3	20/21	0.97	0.06	34,37,47,48	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PO4	C	1326	5/5	0.67	0.16	85,90,91,92	0
4	NAG	E	1324	14/15	0.72	0.17	77,91,94,97	0
4	NAG	A	1322	14/15	0.72	0.13	63,76,85,88	0
5	PO4	E	1328	5/5	0.76	0.16	97,100,104,112	0
5	PO4	A	1327	5/5	0.78	0.19	108,112,116,122	0
4	NAG	C	1321	14/15	0.79	0.12	65,75,80,81	0
4	NAG	E	1322	14/15	0.80	0.12	64,77,87,89	0
4	NAG	A	1323	14/15	0.84	0.10	46,51,56,57	0
5	PO4	A	1328	5/5	0.85	0.14	101,102,104,107	0
4	NAG	C	1322	14/15	0.88	0.09	44,51,54,54	0
5	PO4	C	1327	5/5	0.90	0.12	80,81,85,93	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	E	1323	14/15	0.90	0.09	41,47,52,61	0
4	NAG	A	1321	14/15	0.91	0.09	39,48,54,61	0
4	NAG	E	1321	14/15	0.91	0.10	50,56,58,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.