



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:24 AM UTC

PDB ID : 6CQD / pdb_00006cqd
Title : Crystal structure of HPK1 in complex with ATP analogue (AMPPNP)
Authors : Wu, P.; Lehoux, I.; Franke, Y.; Mortara, K.; Wang, W.
Deposited on : 2018-03-14
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	16% 86% 11% .
1	B	297	23% 83% 13% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase kinase kinase kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total	C	N	O	S	0	0	0
			2277	1463	394	410	10			
1	B	291	Total	C	N	O	S	0	0	0
			2295	1475	397	412	11			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q92918
A	1	SER	-	expression tag	UNP Q92918
A	165	GLU	THR	conflict	UNP Q92918
A	171	GLU	SER	conflict	UNP Q92918
A	294	GLY	-	expression tag	UNP Q92918
A	295	ASN	-	expression tag	UNP Q92918
A	296	SER	-	expression tag	UNP Q92918
B	0	GLY	-	expression tag	UNP Q92918
B	1	SER	-	expression tag	UNP Q92918
B	165	GLU	THR	conflict	UNP Q92918
B	171	GLU	SER	conflict	UNP Q92918
B	294	GLY	-	expression tag	UNP Q92918
B	295	ASN	-	expression tag	UNP Q92918
B	296	SER	-	expression tag	UNP Q92918

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
2	B	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Mg 2 2	0	0
3	B	1	Total Mg 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	125	Total O 125 125	0	0
4	B	92	Total O 92 92	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase kinase kinase 1



- Molecule 1: Mitogen-activated protein kinase kinase kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	146.89Å 51.88Å 107.50Å 90.00° 131.03° 90.00°	Depositor
Resolution (Å)	46.99 – 2.12 46.99 – 2.12	Depositor EDS
% Data completeness (in resolution range)	95.8 (46.99-2.12) 99.9 (46.99-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.203 , 0.250 0.210 , 0.251	Depositor DCC
R_{free} test set	1768 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtriage
Anisotropy	0.640	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4854	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2324	0.74	0/3142
1	B	0.40	1/2344 (0.0%)	0.76	4/3169 (0.1%)
All	All	0.37	1/4668 (0.0%)	0.75	4/6311 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	233	GLN	CD-OE1	5.14	1.33	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	219	HIS	CA-C-N	5.48	124.94	119.24
1	B	219	HIS	C-N-CA	5.48	124.94	119.24
1	B	233	GLN	CA-C-N	5.01	125.55	120.38
1	B	233	GLN	C-N-CA	5.01	125.55	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2277	0	2325	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2295	0	2348	35	0
2	A	31	0	13	4	0
2	B	31	0	13	1	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	125	0	0	2	0
4	B	92	0	0	2	0
All	All	4854	0	4699	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:302:ANP:H5'1	2:B:302:ANP:H8	1.66	0.78
1:B:134:ILE:HD12	1:B:189:LYS:HE3	1.72	0.72
2:A:301:ANP:O2B	2:A:301:ANP:O3G	2.12	0.67
1:B:189:LYS:NZ	1:B:192:TYR:HE1	1.92	0.67
1:B:189:LYS:NZ	1:B:192:TYR:CE1	2.63	0.66
2:A:301:ANP:H8	2:A:301:ANP:H5'1	1.76	0.66
1:B:219:HIS:HB3	1:B:222:ARG:HG3	1.83	0.60
1:B:189:LYS:O	1:B:191:GLY:N	2.36	0.58
1:B:166:LEU:HD13	1:B:189:LYS:HD3	1.88	0.55
1:B:4:VAL:HA	1:B:60:GLN:HG2	1.89	0.54
1:B:173:ILE:HG13	1:B:179:MET:HE1	1.90	0.54
1:A:185:ALA:HA	1:B:228:THR:HG22	1.90	0.53
1:B:237:LEU:HD22	1:B:243:TRP:CD1	2.44	0.52
1:A:39:SER:OG	1:A:41:ASP:OD2	2.26	0.52
1:A:235:PRO:HD2	1:A:255:LEU:HD13	1.92	0.52
1:B:234:PRO:HG3	1:B:257:LYS:HG3	1.92	0.52
1:A:155:ASP:CG	2:A:301:ANP:O2A	2.55	0.49
1:A:225:PHE:O	1:A:229:LYS:HG2	2.12	0.49
1:A:46:LYS:HE2	1:A:48:VAL:HG12	1.94	0.49
1:B:81:TYR:HB2	1:B:88:TRP:HB2	1.95	0.49
1:B:253:VAL:HG22	1:B:263:PRO:HG2	1.96	0.47
1:A:53:ASP:N	4:A:414:HOH:O	2.48	0.47
1:A:223:VAL:HG12	1:A:227:MET:HE2	1.96	0.47
1:B:21:GLN:OE1	1:B:33:LYS:HD3	2.14	0.47
1:B:123:GLN:NE2	4:B:401:HOH:O	2.34	0.46
1:A:223:VAL:HG21	1:B:177:TYR:HE1	1.80	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:189:LYS:C	1:B:191:GLY:H	2.24	0.45
1:B:22:ARG:HE	1:B:30:GLU:HG2	1.81	0.45
1:A:121:VAL:HG11	1:A:201:LEU:HD13	1.99	0.45
1:B:258:SER:C	1:B:260:LYS:H	2.24	0.44
1:A:83:TRP:O	1:A:86:LYS:HG3	2.18	0.44
1:A:188:LEU:HB2	1:B:224:LEU:HD13	1.99	0.44
1:A:5:ASP:O	1:A:8:ILE:HG22	2.18	0.43
1:B:234:PRO:HD3	1:B:257:LYS:HD2	2.00	0.43
1:B:56:VAL:O	1:B:60:GLN:HG3	2.17	0.43
1:B:292:LYS:NZ	4:B:410:HOH:O	2.52	0.42
1:A:22:ARG:HD3	1:A:30:GLU:HG2	2.01	0.42
1:A:92:GLU:OE2	4:A:401:HOH:O	2.22	0.42
1:B:223:VAL:HG12	1:B:227:MET:HE2	2.01	0.42
1:B:59:LEU:H	1:B:59:LEU:HD12	1.84	0.42
1:B:234:PRO:HA	1:B:235:PRO:HD3	1.81	0.42
1:A:86:LYS:HD2	1:A:88:TRP:CZ2	2.55	0.41
1:B:166:LEU:CD1	1:B:189:LYS:HD3	2.50	0.41
1:B:63:ILE:HD11	1:B:87:LEU:HD11	2.02	0.41
1:A:100:GLN:HA	1:A:103:TYR:CE2	2.56	0.41
1:A:181:PRO:HD3	1:B:199:TRP:CE2	2.55	0.41
1:B:224:LEU:O	1:B:228:THR:HG23	2.19	0.41
1:A:31:VAL:HG21	2:A:301:ANP:H5'1	2.02	0.40
1:B:232:TYR:O	1:B:257:LYS:HD2	2.21	0.40
1:A:223:VAL:HG21	1:B:177:TYR:CE1	2.56	0.40
1:B:134:ILE:HD12	1:B:189:LYS:CE	2.47	0.40
1:B:173:ILE:HD12	1:B:173:ILE:HA	1.77	0.40
1:B:188:LEU:HD12	1:B:188:LEU:HA	1.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/297 (96%)	274 (96%)	11 (4%)	0	100	100
1	B	289/297 (97%)	276 (96%)	10 (4%)	3 (1%)	12	8
All	All	574/594 (97%)	550 (96%)	21 (4%)	3 (0%)	24	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	26	GLY
1	B	189	LYS
1	B	190	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/252 (97%)	239 (98%)	6 (2%)	43	48
1	B	247/252 (98%)	239 (97%)	8 (3%)	34	37
All	All	492/504 (98%)	478 (97%)	14 (3%)	38	42

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	VAL
1	A	11	ARG
1	A	59	LEU
1	A	136	ARG
1	A	182	GLU
1	A	226	LEU
1	B	4	VAL
1	B	30	GLU
1	B	136	ARG
1	B	173	ILE
1	B	188	LEU
1	B	189	LYS
1	B	224	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	269	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	10	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ANP	B	302	3	33,33,33	2.06	5 (15%)	45,52,52	0.62	0
2	ANP	A	301	3	33,33,33	2.02	6 (18%)	45,52,52	0.75	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	B	302	3	-	5/18/38/38	0/3/3/3
2	ANP	A	301	3	-	3/18/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	ANP	PG-O1G	8.32	1.58	1.46
2	A	301	ANP	PG-O1G	7.78	1.58	1.46
2	B	302	ANP	PB-O1B	6.51	1.56	1.46
2	A	301	ANP	PB-O1B	6.42	1.55	1.46
2	A	301	ANP	PB-O2B	-2.99	1.48	1.56
2	B	302	ANP	PB-O2B	-2.62	1.49	1.56
2	B	302	ANP	PG-N3B	2.56	1.70	1.63
2	B	302	ANP	PG-O3G	-2.51	1.50	1.56
2	A	301	ANP	PG-N3B	2.46	1.69	1.63
2	A	301	ANP	PG-O2G	-2.37	1.50	1.56
2	A	301	ANP	PB-O3A	-2.16	1.56	1.59

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	ANP	O1G-PG-N3B	-2.48	108.12	111.77
2	A	301	ANP	O2G-PG-O1G	-2.03	108.37	113.45

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	ANP	PB-N3B-PG-O1G
2	A	301	ANP	PG-N3B-PB-O1B
2	B	302	ANP	PB-N3B-PG-O1G
2	B	302	ANP	C5'-O5'-PA-O1A
2	B	302	ANP	C5'-O5'-PA-O2A
2	B	302	ANP	C5'-O5'-PA-O3A
2	B	302	ANP	PB-O3A-PA-O5'
2	A	301	ANP	O4'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	ANP	1	0
2	A	301	ANP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/297 (97%)	0.74	48 (16%) 4 5	22, 37, 82, 134	0
1	B	291/297 (97%)	1.21	68 (23%) 2 2	26, 49, 97, 162	0
All	All	580/594 (97%)	0.98	116 (20%) 3 3	22, 44, 95, 162	0

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	172	PHE	6.7
1	B	231	GLY	5.3
1	B	173	ILE	5.3
1	A	173	ILE	5.2
1	B	4	VAL	5.1
1	B	28	TYR	5.1
1	B	230	SER	4.8
1	B	260	LYS	4.7
1	B	172	PHE	4.4
1	B	239	GLU	4.3
1	A	48	VAL	4.2
1	A	28	TYR	4.2
1	A	170	LEU	4.2
1	B	55	ASP	4.1
1	B	166	LEU	4.0
1	A	56	VAL	4.0
1	A	29	GLY	4.0
1	A	174	GLY	3.9
1	B	25	GLY	3.9
1	A	168	ARG	3.9
1	A	25	GLY	3.9
1	A	54	ASP	3.8
1	A	166	LEU	3.8
1	A	26	GLY	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	PHE	3.8
1	B	170	LEU	3.7
1	A	53	ASP	3.7
1	B	229	LYS	3.7
1	B	238	LYS	3.7
1	B	294	GLY	3.6
1	A	85	GLN	3.6
1	B	52	PRO	3.6
1	B	237	LEU	3.6
1	B	240	LYS	3.6
1	B	190	GLY	3.5
1	B	70	ARG	3.5
1	B	27	THR	3.5
1	B	168	ARG	3.5
1	B	59	LEU	3.4
1	A	86	LYS	3.4
1	B	164	ALA	3.4
1	B	165	GLU	3.3
1	A	55	ASP	3.2
1	B	53	ASP	3.2
1	B	26	GLY	3.2
1	B	261	LYS	3.2
1	B	189	LYS	3.2
1	B	64	LEU	3.1
1	B	188	LEU	3.1
1	A	177	TYR	3.1
1	A	216	PHE	3.0
1	A	277	PRO	3.0
1	A	3	VAL	3.0
1	A	169	ARG	3.0
1	B	293	ASN	3.0
1	A	162	ILE	2.9
1	B	266	THR	2.9
1	B	6	PRO	2.9
1	B	253	VAL	2.8
1	A	27	THR	2.8
1	A	292	LYS	2.8
1	B	232	TYR	2.8
1	B	56	VAL	2.8
1	B	259	PRO	2.8
1	B	130	SER	2.8
1	A	294	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	226	LEU	2.8
1	B	193	ASN	2.7
1	B	256	THR	2.7
1	B	9	PHE	2.7
1	B	50	MET	2.6
1	B	162	ILE	2.6
1	A	24	GLY	2.6
1	B	24	GLY	2.6
1	A	163	GLY	2.6
1	B	292	LYS	2.6
1	A	175	THR	2.6
1	A	181	PRO	2.6
1	A	161	GLN	2.5
1	B	171	GLU	2.5
1	B	134	ILE	2.5
1	A	190	GLY	2.5
1	B	167	ALA	2.5
1	A	47	MET	2.5
1	A	184	ALA	2.5
1	B	236	ARG	2.5
1	A	293	ASN	2.5
1	A	105	VAL	2.4
1	A	186	VAL	2.4
1	B	127	TYR	2.4
1	A	57	SER	2.4
1	B	269	LEU	2.4
1	B	258	SER	2.4
1	A	187	ALA	2.4
1	B	216	PHE	2.4
1	B	225	PHE	2.4
1	B	49	LYS	2.4
1	B	131	GLN	2.3
1	B	191	GLY	2.3
1	B	257	LYS	2.3
1	A	221	LEU	2.3
1	B	5	ASP	2.3
1	B	169	ARG	2.3
1	A	164	ALA	2.3
1	B	163	GLY	2.3
1	A	229	LYS	2.3
1	B	29	GLY	2.3
1	B	23	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	63	ILE	2.2
1	A	278	GLY	2.1
1	B	245	ALA	2.1
1	A	4	VAL	2.1
1	A	11	ARG	2.1
1	A	83	TRP	2.0
1	B	187	ALA	2.0
1	B	262	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ANP	B	302	31/31	0.83	0.13	35,49,151,168	0
3	MG	A	303	1/1	0.85	0.09	49,49,49,49	0
3	MG	B	301	1/1	0.87	0.10	64,64,64,64	0
2	ANP	A	301	31/31	0.94	0.09	25,32,88,133	0
3	MG	A	302	1/1	0.95	0.09	39,39,39,39	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.