



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 10:47 AM UTC

PDB ID : 4CR7 / pdb_00004cr7
Title : Crystal structure of the N-acetyl-D-mannosamine dehydrogenase with n-acetylmannosamine
Authors : Gil-Ortiz, F.; Sola-Carvajal, A.; Garcia-Carmona, F.; Sanchez-Ferrer, A.; Rubio, V.
Deposited on : 2014-02-25
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	4% 85% 11% .
1	B	271	3% 86% 10% .
1	C	271	3% 86% 9% 5%
1	D	271	3% 86% 8% 5%
1	E	271	2% 88% 7% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	271	
1	G	271	
1	H	271	
1	I	271	
1	J	271	
1	K	271	
1	L	271	
1	M	271	
1	N	271	
1	O	271	
1	P	271	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BM3	D	1272	X	-	-	-
3	BM3	G	1273	X	-	-	-
3	BM3	I	1273	X	-	-	-
3	BM3	L	1272	X	-	-	-
3	BM3	M	1272	X	-	-	-
4	MAN	M	1273	-	-	X	-

2 Entry composition


There are 5 unique types of molecules in this entry. The entry contains 30349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-ACYLMANNOSAMINE 1-DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			1823	1130	327	354	12			
1	B	260	Total	C	N	O	S	0	0	0
			1819	1130	327	350	12			
1	C	257	Total	C	N	O	S	0	0	0
			1806	1123	323	348	12			
1	D	257	Total	C	N	O	S	0	1	0
			1818	1129	327	350	12			
1	E	259	Total	C	N	O	S	0	0	0
			1821	1130	326	353	12			
1	F	259	Total	C	N	O	S	0	1	0
			1832	1137	330	353	12			
1	G	258	Total	C	N	O	S	0	0	0
			1813	1127	325	349	12			
1	H	257	Total	C	N	O	S	0	0	0
			1807	1123	324	348	12			
1	I	261	Total	C	N	O	S	0	1	0
			1827	1136	325	353	13			
1	J	260	Total	C	N	O	S	0	0	0
			1826	1133	327	354	12			
1	K	260	Total	C	N	O	S	0	3	0
			1828	1138	327	350	13			
1	L	259	Total	C	N	O	S	0	0	0
			1819	1130	326	351	12			
1	M	260	Total	C	N	O	S	0	0	0
			1829	1135	329	353	12			
1	N	260	Total	C	N	O	S	0	0	0
			1826	1133	327	354	12			
1	O	260	Total	C	N	O	S	0	0	0
			1826	1133	327	354	12			
1	P	260	Total	C	N	O	S	0	0	0
			1826	1133	327	354	12			

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			13	8	5		
2	F	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			7	4	3		
2	G	1	Total	C	O	0	0
			7	4	3		
2	H	1	Total	C	O	0	0
			7	4	3		
2	I	1	Total	C	O	0	0
			7	4	3		
2	J	1	Total	C	O	0	0
			7	4	3		
2	K	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	P	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is 2-acetamido-2-deoxy- α -D-mannopyranose (CCD ID: BM3) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	8	1	6		
3	B	1	Total	C	N	O	0	0
			15	8	1	6		
3	C	1	Total	C	N	O	0	0
			15	8	1	6		
3	D	1	Total	C	N	O	0	0
			15	8	1	6		
3	E	1	Total	C	N	O	0	0
			15	8	1	6		
3	F	1	Total	C	N	O	0	0
			15	8	1	6		
3	G	1	Total	C	N	O	0	0
			15	8	1	6		
3	H	1	Total	C	N	O	0	0
			15	8	1	6		
3	I	1	Total	C	N	O	0	0
			15	8	1	6		
3	J	1	Total	C	N	O	0	0
			15	8	1	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	K	1	Total	C	N	O	0	0
			15	8	1	6		
3	L	1	Total	C	N	O	0	0
			15	8	1	6		
3	M	1	Total	C	N	O	0	0
			15	8	1	6		
3	N	1	Total	C	N	O	0	0
			15	8	1	6		
3	O	1	Total	C	N	O	0	0
			15	8	1	6		
3	P	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 4 is alpha-D-mannopyranose (CCD ID: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			12	6	6		
4	M	1	Total	C	O	0	0
			12	6	6		
4	N	1	Total	C	O	0	0
			12	6	6		
4	P	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	52	Total 52	O 52	0	0
5	B	59	Total 59	O 59	0	0
5	C	51	Total 51	O 51	0	0
5	D	62	Total 62	O 62	0	0
5	E	55	Total 55	O 55	0	0
5	F	53	Total 53	O 53	0	0
5	G	55	Total 55	O 55	0	0
5	H	45	Total 45	O 45	0	0
5	I	61	Total 61	O 61	0	0
5	J	53	Total 53	O 53	0	0
5	K	53	Total 53	O 53	0	0
5	L	52	Total 52	O 52	0	0
5	M	40	Total 40	O 40	0	0
5	N	41	Total 41	O 41	0	0
5	O	33	Total 33	O 33	0	0
5	P	46	Total 46	O 46	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE




- Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE



- Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE




- Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE







• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE





• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE





• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE




• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE




• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE





• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE










• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE





• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE





• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE





• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE



• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE



• Molecule 1: N-ACYLMANNOSAMINE 1-DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	99.39Å 100.12Å 111.60Å 67.43° 89.75° 72.46°	Depositor
Resolution (Å)	15.00 – 2.15 15.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.7 (15.00-2.15) 97.5 (15.00-2.15)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.201 , 0.231 0.203 , 0.232	Depositor DCC
R_{free} test set	10008 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	30349	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 78.03 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.5268e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BM3, PG4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/1844	0.85	1/2498 (0.0%)
1	B	0.84	0/1840	0.83	0/2493
1	C	0.85	0/1826	0.84	0/2472
1	D	0.87	0/1840	0.84	0/2487
1	E	0.85	0/1842	0.84	0/2495
1	F	0.84	0/1853	0.83	0/2510
1	G	0.89	0/1833	0.83	0/2483
1	H	0.85	0/1827	0.84	0/2474
1	I	0.82	0/1851	0.82	0/2508
1	J	0.84	0/1847	0.84	0/2502
1	K	0.84	0/1857	0.82	0/2515
1	L	0.88	0/1839	0.84	0/2490
1	M	0.87	0/1850	0.85	0/2505
1	N	0.87	0/1847	0.84	0/2502
1	O	0.85	0/1847	0.83	0/2502
1	P	0.84	0/1847	0.83	0/2502
All	All	0.85	0/29490	0.84	1/39938 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	VAL	CG1-CB-CG2	7.90	128.17	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1823	0	1813	22	0
1	B	1819	0	1816	20	0
1	C	1806	0	1808	15	2
1	D	1818	0	1825	18	0
1	E	1821	0	1820	13	0
1	F	1832	0	1834	20	0
1	G	1813	0	1814	13	0
1	H	1807	0	1813	13	0
1	I	1827	0	1820	15	0
1	J	1826	0	1822	17	0
1	K	1828	0	1832	24	0
1	L	1819	0	1816	20	0
1	M	1829	0	1829	20	0
1	N	1826	0	1822	20	0
1	O	1826	0	1822	12	0
1	P	1826	0	1822	20	1
2	A	7	0	9	1	0
2	B	7	0	9	1	0
2	E	7	0	9	0	0
2	F	41	0	54	2	0
2	G	7	0	9	0	0
2	H	7	0	9	0	0
2	I	7	0	9	0	0
2	J	7	0	9	0	0
2	K	7	0	9	0	0
2	P	7	0	9	0	0
3	A	15	0	15	3	0
3	B	15	0	15	1	0
3	C	15	0	15	0	0
3	D	15	0	15	0	0
3	E	15	0	15	1	0
3	F	15	0	15	1	0
3	G	15	0	14	1	0
3	H	15	0	15	1	0
3	I	15	0	15	1	0
3	J	15	0	15	2	0
3	K	15	0	15	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	15	0	14	3	0
3	M	15	0	14	1	0
3	N	15	0	15	2	0
3	O	15	0	15	0	0
3	P	15	0	15	0	0
4	D	12	0	12	4	0
4	M	12	0	12	6	0
4	N	12	0	12	4	0
4	P	12	0	12	3	1
5	A	52	0	0	3	0
5	B	59	0	0	3	0
5	C	51	0	0	2	2
5	D	62	0	0	2	0
5	E	55	0	0	1	0
5	F	53	0	0	2	0
5	G	55	0	0	0	0
5	H	45	0	0	1	0
5	I	61	0	0	1	0
5	J	53	0	0	0	0
5	K	53	0	0	1	0
5	L	52	0	0	0	0
5	M	40	0	0	0	0
5	N	41	0	0	1	0
5	O	33	0	0	0	0
5	P	46	0	0	1	2
All	All	30349	0	29548	281	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 281 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:258:LEU:O	4:M:1273:MAN:O4	1.70	1.07
1:D:258:LEU:O	4:D:1273:MAN:O4	1.75	1.03
1:N:258:LEU:O	4:N:1273:MAN:O4	1.75	1.03
1:C:66:CYS:SG	1:C:67:ASP:N	2.52	0.82
1:B:50:ARG:O	1:B:54:THR:HG23	1.81	0.81

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:2032:HOH:O	5:P:2037:HOH:O[1_545]	1.22	0.98
1:C:219:ASP:OD2	5:P:2037:HOH:O[1_545]	1.99	0.21
1:C:258:LEU:O	4:P:1274:MAN:O2[1_545]	2.14	0.06
1:P:219:ASP:OD2	5:C:2032:HOH:O[1_565]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/271 (94%)	249 (97%)	7 (3%)	0	100	100
1	B	256/271 (94%)	244 (95%)	12 (5%)	0	100	100
1	C	251/271 (93%)	243 (97%)	8 (3%)	0	100	100
1	D	251/271 (93%)	243 (97%)	8 (3%)	0	100	100
1	E	255/271 (94%)	248 (97%)	7 (3%)	0	100	100
1	F	256/271 (94%)	249 (97%)	7 (3%)	0	100	100
1	G	252/271 (93%)	242 (96%)	10 (4%)	0	100	100
1	H	251/271 (93%)	241 (96%)	10 (4%)	0	100	100
1	I	258/271 (95%)	248 (96%)	10 (4%)	0	100	100
1	J	256/271 (94%)	247 (96%)	9 (4%)	0	100	100
1	K	258/271 (95%)	251 (97%)	7 (3%)	0	100	100
1	L	253/271 (93%)	244 (96%)	9 (4%)	0	100	100
1	M	256/271 (94%)	248 (97%)	8 (3%)	0	100	100
1	N	256/271 (94%)	247 (96%)	9 (4%)	0	100	100
1	O	256/271 (94%)	247 (96%)	9 (4%)	0	100	100
1	P	256/271 (94%)	248 (97%)	8 (3%)	0	100	100
All	All	4077/4336 (94%)	3939 (97%)	138 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/183 (93%)	167 (98%)	3 (2%)	51	58
1	B	169/183 (92%)	165 (98%)	4 (2%)	43	47
1	C	169/183 (92%)	166 (98%)	3 (2%)	51	58
1	D	171/183 (93%)	165 (96%)	6 (4%)	32	32
1	E	171/183 (93%)	165 (96%)	6 (4%)	32	32
1	F	172/183 (94%)	164 (95%)	8 (5%)	23	21
1	G	170/183 (93%)	166 (98%)	4 (2%)	43	47
1	H	170/183 (93%)	166 (98%)	4 (2%)	43	47
1	I	170/183 (93%)	166 (98%)	4 (2%)	43	47
1	J	171/183 (93%)	167 (98%)	4 (2%)	44	49
1	K	171/183 (93%)	166 (97%)	5 (3%)	37	39
1	L	170/183 (93%)	166 (98%)	4 (2%)	43	47
1	M	171/183 (93%)	167 (98%)	4 (2%)	44	49
1	N	171/183 (93%)	167 (98%)	4 (2%)	44	49
1	O	171/183 (93%)	167 (98%)	4 (2%)	44	49
1	P	171/183 (93%)	166 (97%)	5 (3%)	37	39
All	All	2728/2928 (93%)	2656 (97%)	72 (3%)	40	43

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	201	VAL
1	P	204	ASN
1	N	68	LEU
1	O	201	VAL
1	F	91	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	204	ASN
1	O	204	ASN
1	I	204	ASN
1	M	204	ASN
1	I	187	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

34 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PG4	A	1272	-	6,6,12	0.75	0	5,5,11	0.35	0
3	BM3	G	1273	-	15,15,15	2.69	4 (26%)	21,21,21	2.15	3 (14%)
3	BM3	D	1272	-	15,15,15	2.36	4 (26%)	21,21,21	2.01	4 (19%)
2	PG4	J	1272	-	6,6,12	0.41	0	5,5,11	0.48	0
3	BM3	O	1272	-	15,15,15	2.92	4 (26%)	21,21,21	2.24	7 (33%)
2	PG4	F	1277	-	6,6,12	0.78	0	5,5,11	0.24	0
3	BM3	K	1273	-	15,15,15	2.89	4 (26%)	21,21,21	2.04	5 (23%)
3	BM3	M	1272	-	15,15,15	2.98	3 (20%)	21,21,21	2.28	4 (19%)
2	PG4	H	1272	-	6,6,12	0.39	0	5,5,11	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BM3	F	1275	-	15,15,15	2.95	3 (20%)	21,21,21	2.02	5 (23%)
2	PG4	F	1273	-	6,6,12	0.90	0	5,5,11	0.75	0
3	BM3	A	1273	-	15,15,15	2.70	5 (33%)	21,21,21	2.16	5 (23%)
2	PG4	G	1272	-	6,6,12	0.43	0	5,5,11	0.69	0
3	BM3	C	1272	-	15,15,15	2.50	3 (20%)	21,21,21	2.22	5 (23%)
4	MAN	N	1273	-	12,12,12	1.42	2 (16%)	17,17,17	2.85	11 (64%)
2	PG4	P	1272	-	6,6,12	0.82	0	5,5,11	0.45	0
3	BM3	P	1273	-	15,15,15	2.70	2 (13%)	21,21,21	2.19	4 (19%)
2	PG4	F	1276	-	6,6,12	0.71	0	5,5,11	0.11	0
2	PG4	F	1272	-	12,12,12	0.73	0	11,11,11	0.43	0
2	PG4	I	1272	-	6,6,12	0.74	0	5,5,11	0.39	0
3	BM3	I	1273	-	15,15,15	3.06	2 (13%)	21,21,21	2.13	4 (19%)
2	PG4	F	1274	-	6,6,12	0.44	0	5,5,11	0.47	0
2	PG4	K	1272	-	6,6,12	0.63	0	5,5,11	0.55	0
4	MAN	D	1273	-	12,12,12	1.60	3 (25%)	17,17,17	2.58	9 (52%)
2	PG4	E	1272	-	6,6,12	0.58	0	5,5,11	0.21	0
4	MAN	P	1274	-	12,12,12	1.91	3 (25%)	17,17,17	2.69	9 (52%)
3	BM3	H	1273	-	15,15,15	2.75	2 (13%)	21,21,21	1.89	3 (14%)
4	MAN	M	1273	-	12,12,12	1.47	2 (16%)	17,17,17	2.82	8 (47%)
3	BM3	N	1272	-	15,15,15	2.71	3 (20%)	21,21,21	2.09	4 (19%)
3	BM3	J	1273	-	15,15,15	2.74	4 (26%)	21,21,21	1.91	5 (23%)
3	BM3	L	1272	-	15,15,15	2.59	5 (33%)	21,21,21	2.06	4 (19%)
3	BM3	B	1273	-	15,15,15	2.72	2 (13%)	21,21,21	2.30	7 (33%)
2	PG4	B	1272	-	6,6,12	0.69	0	5,5,11	0.13	0
3	BM3	E	1273	-	15,15,15	3.04	2 (13%)	21,21,21	1.67	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	A	1272	-	-	3/4/4/10	-
3	BM3	G	1273	-	1/1/6/7	2/6/26/26	0/1/1/1
3	BM3	D	1272	-	1/1/6/7	0/6/26/26	0/1/1/1
2	PG4	J	1272	-	-	3/4/4/10	-
3	BM3	O	1272	-	-	2/6/26/26	0/1/1/1
2	PG4	F	1277	-	-	2/4/4/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BM3	K	1273	-	-	2/6/26/26	0/1/1/1
3	BM3	M	1272	-	1/1/6/7	2/6/26/26	0/1/1/1
2	PG4	H	1272	-	-	3/4/4/10	-
3	BM3	F	1275	-	-	2/6/26/26	0/1/1/1
2	PG4	F	1273	-	-	4/4/4/10	-
3	BM3	A	1273	-	-	2/6/26/26	0/1/1/1
2	PG4	G	1272	-	-	2/4/4/10	-
3	BM3	C	1272	-	-	0/6/26/26	0/1/1/1
4	MAN	N	1273	-	-	2/2/22/22	0/1/1/1
2	PG4	P	1272	-	-	2/4/4/10	-
3	BM3	P	1273	-	-	4/6/26/26	0/1/1/1
2	PG4	F	1276	-	-	2/4/4/10	-
2	PG4	F	1272	-	-	5/10/10/10	-
2	PG4	I	1272	-	-	1/4/4/10	-
3	BM3	I	1273	-	1/1/6/7	0/6/26/26	0/1/1/1
2	PG4	F	1274	-	-	3/4/4/10	-
2	PG4	K	1272	-	-	1/4/4/10	-
4	MAN	D	1273	-	-	2/2/22/22	0/1/1/1
2	PG4	E	1272	-	-	3/4/4/10	-
4	MAN	P	1274	-	-	2/2/22/22	0/1/1/1
3	BM3	H	1273	-	-	2/6/26/26	0/1/1/1
4	MAN	M	1273	-	-	2/2/22/22	0/1/1/1
3	BM3	N	1272	-	-	1/6/26/26	0/1/1/1
3	BM3	J	1273	-	-	0/6/26/26	0/1/1/1
3	BM3	L	1272	-	1/1/6/7	0/6/26/26	0/1/1/1
3	BM3	B	1273	-	-	0/6/26/26	0/1/1/1
2	PG4	B	1272	-	-	1/4/4/10	-
3	BM3	E	1273	-	-	4/6/26/26	0/1/1/1

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	1273	BM3	C1-C2	-11.09	1.39	1.52
3	E	1273	BM3	C1-C2	-10.74	1.40	1.52
3	M	1272	BM3	C1-C2	-10.24	1.40	1.52
3	F	1275	BM3	C1-C2	-10.20	1.40	1.52
3	K	1273	BM3	C1-C2	-10.10	1.40	1.52

The worst 5 of 109 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	1272	BM3	O1-C1-C2	6.43	122.58	109.22
3	P	1273	BM3	O5-C1-C2	6.42	115.97	109.52
4	M	1273	MAN	C1-O5-C5	-6.37	101.33	113.65
4	P	1274	MAN	C1-O5-C5	-6.01	102.03	113.65
4	N	1273	MAN	C1-O5-C5	-5.94	102.15	113.65

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	1272	BM3	C1
3	G	1273	BM3	C1
3	I	1273	BM3	C1
3	L	1272	BM3	C1
3	M	1272	BM3	C1

5 of 66 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	1275	BM3	O7-C7-N2-C2
3	F	1275	BM3	C8-C7-N2-C2
3	G	1273	BM3	C4-C5-C6-O6
3	E	1273	BM3	C4-C5-C6-O6
3	P	1273	BM3	O5-C5-C6-O6

There are no ring outliers.

19 monomers are involved in 39 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1272	PG4	1	0
3	G	1273	BM3	1	0
3	K	1273	BM3	1	0
3	M	1272	BM3	1	0
3	F	1275	BM3	1	0
3	A	1273	BM3	3	0
4	N	1273	MAN	4	0
2	F	1272	PG4	2	0
3	I	1273	BM3	1	0
4	D	1273	MAN	4	0
4	P	1274	MAN	3	1
3	H	1273	BM3	1	0
4	M	1273	MAN	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	N	1272	BM3	2	0
3	J	1273	BM3	2	0
3	L	1272	BM3	3	0
3	B	1273	BM3	1	0
2	B	1272	PG4	1	0
3	E	1273	BM3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	260/271 (95%)	0.20	11 (4%)	40	45	12, 26, 55, 68	0
1	B	260/271 (95%)	0.12	8 (3%)	51	55	11, 25, 51, 67	0
1	C	257/271 (94%)	0.08	7 (2%)	56	60	11, 23, 47, 67	0
1	D	257/271 (94%)	-0.01	9 (3%)	47	51	10, 21, 47, 64	1 (0%)
1	E	259/271 (95%)	-0.00	5 (1%)	66	70	10, 22, 48, 64	0
1	F	259/271 (95%)	-0.08	5 (1%)	66	70	10, 21, 45, 57	1 (0%)
1	G	258/271 (95%)	0.10	12 (4%)	36	41	11, 23, 49, 65	0
1	H	257/271 (94%)	0.16	4 (1%)	70	74	11, 25, 53, 64	0
1	I	261/271 (96%)	0.21	10 (3%)	44	49	12, 24, 51, 65	1 (0%)
1	J	260/271 (95%)	0.22	8 (3%)	51	55	12, 25, 52, 66	0
1	K	260/271 (95%)	0.29	14 (5%)	31	35	12, 26, 52, 68	3 (1%)
1	L	259/271 (95%)	0.22	6 (2%)	61	64	12, 24, 53, 72	0
1	M	260/271 (95%)	0.18	7 (2%)	56	60	12, 24, 47, 62	0
1	N	260/271 (95%)	0.21	6 (2%)	61	64	13, 25, 50, 66	0
1	O	260/271 (95%)	0.14	9 (3%)	47	51	12, 24, 49, 73	0
1	P	260/271 (95%)	0.24	11 (4%)	40	45	12, 25, 54, 68	0
All	All	4147/4336 (95%)	0.14	132 (3%)	50	54	10, 24, 51, 73	6 (0%)

The worst 5 of 132 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	70	ASP	7.8
1	J	70	ASP	6.2
1	K	70	ASP	5.1
1	O	68	LEU	4.7
1	P	70	ASP	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BM3	F	1275	15/15	0.61	0.16	41,43,47,47	15
3	BM3	L	1272	15/15	0.65	0.14	36,40,44,44	15
3	BM3	M	1272	15/15	0.70	0.14	44,50,52,53	0
3	BM3	I	1273	15/15	0.71	0.14	43,47,49,52	15
3	BM3	K	1273	15/15	0.73	0.14	37,51,54,54	15
3	BM3	A	1273	15/15	0.73	0.12	42,49,52,52	0
3	BM3	D	1272	15/15	0.73	0.12	37,47,50,50	0
2	PG4	P	1272	7/13	0.74	0.18	35,38,40,40	7
2	PG4	F	1276	7/13	0.74	0.17	50,52,55,57	7
3	BM3	P	1273	15/15	0.74	0.12	34,40,45,45	15
3	BM3	O	1272	15/15	0.75	0.13	38,39,41,41	15
2	PG4	F	1277	7/13	0.75	0.21	55,59,67,67	7
2	PG4	F	1273	7/13	0.76	0.18	29,32,36,42	7
3	BM3	E	1273	15/15	0.76	0.13	32,35,39,43	15
3	BM3	B	1273	15/15	0.76	0.14	36,39,41,41	15
4	MAN	N	1273	12/12	0.76	0.16	20,31,32,32	12
4	MAN	D	1273	12/12	0.77	0.18	21,29,31,34	12
2	PG4	A	1272	7/13	0.78	0.11	25,29,32,33	7
3	BM3	G	1273	15/15	0.78	0.11	43,49,52,53	0
3	BM3	C	1272	15/15	0.79	0.10	39,46,47,47	0
2	PG4	F	1272	13/13	0.80	0.12	38,41,42,43	13
4	MAN	M	1273	12/12	0.80	0.14	18,27,29,30	12
2	PG4	G	1272	7/13	0.80	0.11	37,37,41,42	7
4	MAN	P	1274	12/12	0.80	0.15	20,29,31,31	12
2	PG4	F	1274	7/13	0.81	0.12	32,35,38,41	7
2	PG4	K	1272	7/13	0.81	0.12	32,33,34,34	7
3	BM3	N	1272	15/15	0.81	0.12	42,47,52,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PG4	B	1272	7/13	0.84	0.13	35,36,37,40	7
2	PG4	J	1272	7/13	0.84	0.10	34,35,36,38	7
2	PG4	E	1272	7/13	0.85	0.10	34,35,37,38	7
3	BM3	J	1273	15/15	0.85	0.09	47,49,53,54	0
2	PG4	H	1272	7/13	0.85	0.10	37,37,39,40	7
3	BM3	H	1273	15/15	0.85	0.12	36,38,41,43	15
2	PG4	I	1272	7/13	0.89	0.11	31,32,36,36	7

6.5 Other polymers [i](#)

There are no such residues in this entry.