



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:22 AM UTC

PDB ID : 8CR6 / pdb_00008cr6
Title : mouse Interleukin-12
Authors : Merceron, R.; Felix, J.; Lambert, E.; Bloch, Y.; Savvides, S.N.
Deposited on : 2023-03-07
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	75% 13% 13%
2	B	253	58% 7% 35%
3	C	3	100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3639 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-12 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2324	1463	389	455	17			

- Molecule 2 is a protein called Interleukin-12 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	164	Total	C	N	O	S	0	0	0
			1260	780	219	246	15			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	216	GLY	-	expression tag	UNP P43431
B	217	THR	-	expression tag	UNP P43431
B	218	SER	-	expression tag	UNP P43431
B	219	ASP	-	expression tag	UNP P43431
B	220	GLU	-	expression tag	UNP P43431
B	221	VAL	-	expression tag	UNP P43431
B	222	ASP	-	expression tag	UNP P43431
B	223	GLY	-	expression tag	UNP P43431
B	224	GLY	-	expression tag	UNP P43431
B	225	SER	-	expression tag	UNP P43431
B	226	GLY	-	expression tag	UNP P43431
B	227	GLY	-	expression tag	UNP P43431
B	228	SER	-	expression tag	UNP P43431
B	229	GLY	-	expression tag	UNP P43431
B	230	LEU	-	expression tag	UNP P43431
B	231	ASN	-	expression tag	UNP P43431
B	232	ASP	-	expression tag	UNP P43431
B	233	ILE	-	expression tag	UNP P43431
B	234	PHE	-	expression tag	UNP P43431
B	235	GLU	-	expression tag	UNP P43431

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	236	ALA	-	expression tag	UNP P43431
B	237	GLN	-	expression tag	UNP P43431
B	238	LYS	-	expression tag	UNP P43431
B	239	ILE	-	expression tag	UNP P43431
B	240	GLU	-	expression tag	UNP P43431
B	241	TRP	-	expression tag	UNP P43431
B	242	HIS	-	expression tag	UNP P43431
B	243	GLU	-	expression tag	UNP P43431
B	244	GLY	-	expression tag	UNP P43431
B	245	ARG	-	expression tag	UNP P43431
B	246	THR	-	expression tag	UNP P43431
B	247	LYS	-	expression tag	UNP P43431
B	248	HIS	-	expression tag	UNP P43431
B	249	HIS	-	expression tag	UNP P43431
B	250	HIS	-	expression tag	UNP P43431
B	251	HIS	-	expression tag	UNP P43431
B	252	HIS	-	expression tag	UNP P43431
B	253	HIS	-	expression tag	UNP P43431

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-12 subunit beta



- Molecule 2: Interleukin-12 subunit alpha



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.11Å 70.77Å 127.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.34 – 2.85 47.34 – 2.85	Depositor EDS
% Data completeness (in resolution range)	96.2 (47.34-2.85) 96.2 (47.34-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.86Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.255 , 0.296 (Not available) , 0.299	Depositor DCC
R_{free} test set	627 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	97.9	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 71.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3639	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/2378	0.28	0/3227
2	B	0.11	0/1273	0.25	0/1717
All	All	0.11	0/3651	0.27	0/4944

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2324	0	2204	25	0
2	B	1260	0	1231	11	0
3	C	39	0	34	0	0
4	B	14	0	13	0	0
5	A	2	0	0	0	0
All	All	3639	0	3482	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:MET:HE1	1:A:332:ARG:H	1.47	0.78
2:B:48:VAL:HG12	2:B:195:LEU:HB3	1.81	0.62
2:B:47:MET:HE2	2:B:151:LEU:HD23	1.83	0.61
1:A:136:ARG:HG3	1:A:192:GLN:HG2	1.84	0.60
1:A:90:CYS:SG	1:A:98:SER:OG	2.60	0.60
1:A:26:LEU:HD23	1:A:27:GLU:HG3	1.86	0.57
1:A:269:LYS:HB2	1:A:295:LEU:HD11	1.86	0.56
1:A:50:CYS:SG	1:A:98:SER:OG	2.63	0.54
2:B:151:LEU:HD21	2:B:191:LEU:HD11	1.90	0.53
1:A:275:GLN:NE2	1:A:311:CYS:SG	2.77	0.51
1:A:45:THR:HG22	1:A:78:THR:HG22	1.93	0.49
1:A:297:GLU:O	1:A:317:ARG:NH2	2.45	0.49
2:B:116:MET:HE2	2:B:212:LEU:HD21	1.94	0.49
1:A:227:ILE:HA	1:A:230:ILE:HG12	1.95	0.48
1:A:239:LEU:HD13	1:A:253:TRP:HB3	1.96	0.47
1:A:243:PRO:HA	1:A:249:VAL:HG12	1.96	0.47
2:B:160:ASP:HA	2:B:163:MET:HE3	1.96	0.47
2:B:137:ILE:HG23	2:B:187:VAL:HG22	1.97	0.47
1:A:122:ASN:HD22	1:A:122:ASN:C	2.24	0.46
1:A:309:ASN:HA	1:A:330:PRO:HA	1.97	0.46
1:A:296:VAL:HG12	1:A:298:LYS:H	1.80	0.46
2:B:104:GLY:HA2	2:B:168:HIS:HB2	1.97	0.45
1:A:129:GLU:HA	1:A:230:ILE:HB	1.99	0.44
1:A:272:VAL:HG22	1:A:312:VAL:HG22	2.00	0.44
1:A:32:VAL:HG22	1:A:103:LEU:HB2	2.00	0.43
1:A:169:MET:HE2	1:A:169:MET:HB3	1.97	0.42
1:A:244:LEU:HG	1:A:245:LYS:H	1.84	0.42
2:B:45:ASP:O	2:B:48:VAL:HG22	2.19	0.42
1:A:147:MET:HE3	1:A:147:MET:HB2	1.85	0.42
1:A:195:VAL:O	1:A:195:VAL:HG13	2.19	0.42
1:A:201:GLU:HB2	2:B:78:LEU:HD22	2.02	0.42
2:B:48:VAL:O	2:B:52:ARG:HG2	2.20	0.41
1:A:27:GLU:HB2	1:A:30:VAL:HB	2.03	0.41
1:A:59:THR:OG1	1:A:91:HIS:HB2	2.21	0.41
2:B:141:LEU:HD13	2:B:149:ILE:HG23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/335 (85%)	275 (96%)	10 (4%)	0	100	100
2	B	156/253 (62%)	150 (96%)	6 (4%)	0	100	100
All	All	441/588 (75%)	425 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/307 (86%)	262 (100%)	1 (0%)	84	92
2	B	141/224 (63%)	141 (100%)	0	100	100
All	All	404/531 (76%)	403 (100%)	1 (0%)	87	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	331	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	ASN
1	A	152	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	36	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,1	14,14,15	0.33	0	17,19,21	0.60	0
3	NAG	C	2	3	14,14,15	0.25	0	17,19,21	0.43	0
3	BMA	C	3	3	11,11,12	0.62	0	15,15,17	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	1/6/23/26	0/1/1/1
3	BMA	C	3	3	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	3	BMA	O5-C5-C6-O6
3	C	1	NAG	C1-C2-N2-C7
3	C	2	NAG	C1-C2-N2-C7
3	C	1	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	301	2	14,14,15	0.29	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	301	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/335 (87%)	-0.08	2 (0%) 84 80	58, 104, 187, 264	0
2	B	164/253 (64%)	0.25	3 (1%) 67 61	57, 99, 180, 236	0
All	All	457/588 (77%)	0.04	5 (1%) 78 73	57, 101, 187, 264	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	61	THR	2.3
1	A	45	THR	2.2
2	B	164	GLN	2.1
2	B	130	TYR	2.1
1	A	122	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	C	3	11/12	0.49	0.12	111,123,130,131	0
3	NAG	C	2	14/15	0.89	0.07	89,103,112,118	0
3	NAG	C	1	14/15	0.90	0.09	70,90,94,95	0

The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	301	14/15	0.57	0.11	162,169,176,182	0

6.5 Other polymers [i](#)

There are no such residues in this entry.