



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:20 PM UTC

PDB ID : 3CSN / pdb\_00003csn  
Title : Structure of the *Serratia marcescens* hemophore receptor HasR in complex with its hemophore HasA  
Authors : Krieg, S.; Diederichs, K.  
Deposited on : 2008-04-10  
Resolution : 3.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	865	 3% 58% 26% 13%
1	B	865	 3% 56% 28% 13%
2	C	206	 2% 60% 18% 21%
2	D	206	 2% 56% 22% 21%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HasR protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	753	5889	3674	1043	1159	13	0	0	0
1	B	753	5889	3674	1043	1159	13	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	645	ALA	GLY	SEE REMARK 999	UNP Q79AD2
B	645	ALA	GLY	SEE REMARK 999	UNP Q79AD2

- Molecule 2 is a protein called Hemophore HasA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	163	1184	743	189	251	1	0	0	0
2	D	163	1184	743	189	251	1	0	0	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-17	MET	-	expression tag	UNP Q54450
C	-16	ARG	-	expression tag	UNP Q54450
C	-15	GLY	-	expression tag	UNP Q54450
C	-14	SER	-	expression tag	UNP Q54450
C	-13	HIS	-	expression tag	UNP Q54450
C	-12	HIS	-	expression tag	UNP Q54450
C	-11	HIS	-	expression tag	UNP Q54450
C	-10	HIS	-	expression tag	UNP Q54450
C	-9	HIS	-	expression tag	UNP Q54450
C	-8	HIS	-	expression tag	UNP Q54450

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-7	GLY	-	expression tag	UNP Q54450
C	-6	ILE	-	expression tag	UNP Q54450
C	-5	ARG	-	expression tag	UNP Q54450
C	-4	MET	-	expression tag	UNP Q54450
C	-3	ARG	-	expression tag	UNP Q54450
C	-2	ALA	-	expression tag	UNP Q54450
C	-1	ARG	-	expression tag	UNP Q54450
C	0	TYR	-	expression tag	UNP Q54450
C	1	PRO	-	expression tag	UNP Q54450
D	-17	MET	-	expression tag	UNP Q54450
D	-16	ARG	-	expression tag	UNP Q54450
D	-15	GLY	-	expression tag	UNP Q54450
D	-14	SER	-	expression tag	UNP Q54450
D	-13	HIS	-	expression tag	UNP Q54450
D	-12	HIS	-	expression tag	UNP Q54450
D	-11	HIS	-	expression tag	UNP Q54450
D	-10	HIS	-	expression tag	UNP Q54450
D	-9	HIS	-	expression tag	UNP Q54450
D	-8	HIS	-	expression tag	UNP Q54450
D	-7	GLY	-	expression tag	UNP Q54450
D	-6	ILE	-	expression tag	UNP Q54450
D	-5	ARG	-	expression tag	UNP Q54450
D	-4	MET	-	expression tag	UNP Q54450
D	-3	ARG	-	expression tag	UNP Q54450
D	-2	ALA	-	expression tag	UNP Q54450
D	-1	ARG	-	expression tag	UNP Q54450
D	0	TYR	-	expression tag	UNP Q54450
D	1	PRO	-	expression tag	UNP Q54450

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	10	Total O 10 10	0	0
4	B	7	Total O 7 7	0	0
4	C	1	Total O 1 1	0	0
4	D	1	Total O 1 1	0	0







## 4 Data and refinement statistics i

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.65Å 164.74Å 597.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.43 – 3.00 49.43 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.43-3.00) 99.3 (49.43-3.00)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 3.01Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.217 , 0.244 0.215 , 0.242	Depositor DCC
$R_{free}$ test set	3890 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.4	Xtrriage
Anisotropy	0.543	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 66.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.039 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	14195	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.63	0/6027	0.93	12/8188 (0.1%)
1	B	0.63	0/6027	0.93	16/8188 (0.2%)
2	C	0.53	1/1211 (0.1%)	0.84	2/1649 (0.1%)
2	D	0.54	1/1211 (0.1%)	0.88	4/1649 (0.2%)
All	All	0.62	2/14476 (0.0%)	0.92	34/19674 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	42	SER	C-N	6.89	1.41	1.33
2	D	42	SER	C-N	5.76	1.40	1.33

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	42	SER	CA-C-N	-11.60	112.11	122.43
2	D	42	SER	C-N-CA	-11.60	112.11	122.43
2	C	42	SER	CA-C-N	-9.09	113.24	122.69
2	C	42	SER	C-N-CA	-9.09	113.24	122.69
1	A	564	SER	CA-C-N	8.64	128.58	119.85

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5889	0	5597	204	0
1	B	5889	0	5597	209	0
2	C	1184	0	1078	31	0
2	D	1184	0	1078	34	0
3	A	6	0	8	2	0
3	B	24	0	32	1	0
4	A	10	0	0	0	0
4	B	7	0	0	1	0
4	C	1	0	0	0	0
4	D	1	0	0	1	0
All	All	14195	0	13390	468	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 468 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:779:LYS:HZ3	1:A:820:LYS:HG3	1.26	0.99
1:B:779:LYS:HZ3	1:B:820:LYS:HG3	1.33	0.94
1:A:575:VAL:HG22	1:A:577:TRP:H	1.35	0.91
1:B:720:TRP:CD1	1:B:721:LEU:HD13	2.08	0.88
1:B:575:VAL:HG22	1:B:577:TRP:H	1.37	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	751/865 (87%)	657 (88%)	84 (11%)	10 (1%)	<b>9</b> <b>38</b>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	751/865 (87%)	652 (87%)	89 (12%)	10 (1%)	9	38
2	C	159/206 (77%)	147 (92%)	9 (6%)	3 (2%)	6	30
2	D	159/206 (77%)	146 (92%)	10 (6%)	3 (2%)	6	30
All	All	1820/2142 (85%)	1602 (88%)	192 (10%)	26 (1%)	9	36

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	480	PRO
1	B	301	ASP
1	B	480	PRO
1	A	573	PRO
1	A	832	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	616/695 (89%)	587 (95%)	29 (5%)	23	58
1	B	616/695 (89%)	585 (95%)	31 (5%)	22	56
2	C	124/158 (78%)	123 (99%)	1 (1%)	73	86
2	D	124/158 (78%)	123 (99%)	1 (1%)	73	86
All	All	1480/1706 (87%)	1418 (96%)	62 (4%)	26	61

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	780	LEU
1	B	720	TRP
1	B	318	PRO
1	B	712	ASN
1	B	832	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	119	ASN
2	C	41	ASN
1	B	257	HIS
1	B	833	ASN
1	B	248	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	B	868	-	5,5,5	0.42	0	5,5,5	0.84	0
3	GOL	B	866	-	5,5,5	0.54	0	5,5,5	0.42	0
3	GOL	B	869	-	5,5,5	0.43	0	5,5,5	0.45	0
3	GOL	A	866	-	5,5,5	0.44	0	5,5,5	0.64	0
3	GOL	B	867	-	5,5,5	0.48	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	868	-	-	2/4/4/4	-
3	GOL	B	866	-	-	2/4/4/4	-
3	GOL	B	869	-	-	0/4/4/4	-
3	GOL	A	866	-	-	2/4/4/4	-
3	GOL	B	867	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	866	GOL	O1-C1-C2-O2
3	A	866	GOL	O1-C1-C2-C3
3	B	866	GOL	O1-C1-C2-C3
3	B	867	GOL	O1-C1-C2-C3
3	B	868	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	866	GOL	1	0
3	A	866	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	753/865 (87%)	0.12	25 (3%)	49 28	44, 76, 123, 215	0
1	B	753/865 (87%)	0.10	25 (3%)	49 28	43, 75, 122, 206	0
2	C	163/206 (79%)	0.12	5 (3%)	51 30	36, 95, 119, 140	0
2	D	163/206 (79%)	0.11	4 (2%)	58 35	36, 94, 118, 140	0
All	All	1832/2142 (85%)	0.11	59 (3%)	50 29	36, 79, 122, 215	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	605	SER	5.8
2	D	1	PRO	5.2
2	C	28	GLY	4.9
1	A	603	HIS	4.9
1	B	603	HIS	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	866	6/6	0.86	0.23	59,103,141,173	0
3	GOL	B	866	6/6	0.87	0.17	58,76,87,88	0
3	GOL	B	867	6/6	0.90	0.17	50,93,107,116	0
3	GOL	B	869	6/6	0.91	0.15	41,95,107,110	0
3	GOL	B	868	6/6	0.94	0.15	49,73,97,98	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.