



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 09:48 PM UTC

PDB ID : 9CSG / pdb_00009csg
Title : Human Serum Albumin Bound to Cerastecin Compound 5e
Authors : Hruza, A.; Klein, D.J.; Ishchenko, A.
Deposited on : 2024-07-23
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	582	11% 97% .

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9746 atoms, of which 4723 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Albumin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	582	Total	C	H	N	O	S	4551	0	0
			9181	2923	4551	783	883	41			

- Molecule 2 is MYRISTIC ACID (CCD ID: MYR) (formula: $C_{14}H_{28}O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	27	0
			43	14	27	2		
2	A	1	Total	C	H	O	18	0
			31	11	18	2		
2	A	1	Total	C	H	O	19	0
			33	12	19	2		
2	A	1	Total	C	H	O	18	0
			31	11	18	2		
2	A	1	Total	C	H	O	27	0
			43	14	27	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	27	0
			43	14	27	2		

- Molecule 3 is 2-(4-butylbenzene-1-sulfonamido)-5-(4-{3-carboxy-4-[4-(2-methoxyethyl)benzene-1-sulfonamido]anilino}-4-oxobutanamido)benzoic acid (CCD ID: A1AZR) (formula: $C_{37}H_{40}N_4O_{11}S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	S	36	0
			90	37	36	4	11	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	251	Total	O	0	0
			251	251		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Albumin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	172.65Å 38.62Å 97.78Å 90.00° 104.89° 90.00°	Depositor
Resolution (Å)	94.50 – 1.91 94.50 – 1.91	Depositor EDS
% Data completeness (in resolution range)	62.1 (94.50-1.91) 62.1 (94.50-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 1.91Å)	Xtriage
Refinement program	BUSTER 2.11.8 (23-JAN-2024)	Depositor
R, R_{free}	0.230 , 0.262 0.219 , 0.252	Depositor DCC
R_{free} test set	1523 reflections (3.12%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9746	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1AZR, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/4720	0.97	0/6366

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4630	4551	4552	10	0
2	A	88	136	137	2	0
3	A	54	36	0	2	0
4	A	251	0	0	0	0
All	All	5023	4723	4689	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:CYS:SG	1:A:105:HIS:HD2	2.32	0.53
1:A:90:CYS:SG	1:A:105:HIS:CD2	3.08	0.47
1:A:186:ARG:HB2	3:A:907:A1AZR:N3	2.31	0.46
1:A:459:GLN:HG3	3:A:907:A1AZR:O1	2.16	0.46
1:A:411:TYR:CZ	2:A:904:MYR:H72	2.52	0.45
1:A:67:HIS:CE1	1:A:251:LEU:HD12	2.52	0.45
1:A:214:TRP:CD1	1:A:343:VAL:HG11	2.54	0.43
1:A:25:ILE:HD11	1:A:139:LEU:HD22	2.01	0.42
1:A:507:PHE:CD1	2:A:905:MYR:H101	2.55	0.41
1:A:25:ILE:O	1:A:29:GLN:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	580/582 (100%)	568 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	509/509 (100%)	507 (100%)	2 (0%)	84	83

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	GLU
1	A	66	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	HIS
1	A	33	GLN
1	A	67	HIS
1	A	105	HIS
1	A	503	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MYR	A	905	-	15,15,15	0.58	0	15,15,15	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MYR	A	906	-	15,15,15	0.57	0	15,15,15	0.53	0
3	A1AZR	A	907	-	57,57,57	0.68	1 (1%)	80,80,80	0.87	3 (3%)
2	MYR	A	904	-	12,12,15	0.71	0	12,12,15	0.56	0
2	MYR	A	902	-	12,12,15	0.66	0	12,12,15	0.54	0
2	MYR	A	903	-	13,13,15	0.62	0	13,13,15	0.58	0
2	MYR	A	901	-	15,15,15	0.52	0	15,15,15	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MYR	A	905	-	-	3/13/13/13	-
2	MYR	A	906	-	-	7/13/13/13	-
3	A1AZR	A	907	-	-	4/51/51/51	0/4/4/4
2	MYR	A	904	-	-	4/10/10/13	-
2	MYR	A	902	-	-	0/10/10/13	-
2	MYR	A	903	-	-	3/11/11/13	-
2	MYR	A	901	-	-	5/13/13/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	907	A1AZR	S1-N3	-2.25	1.60	1.63

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	907	A1AZR	O8-S1-O9	-2.94	115.95	119.52
3	A	907	A1AZR	C7-S-N	2.71	110.24	106.88
3	A	907	A1AZR	O1-S-O	-2.48	116.51	119.52

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	906	MYR	C1-C2-C3-C4
2	A	906	MYR	C11-C10-C9-C8
2	A	904	MYR	C5-C6-C7-C8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	904	MYR	C4-C5-C6-C7
2	A	904	MYR	C2-C3-C4-C5
2	A	906	MYR	C11-C12-C13-C14
3	A	907	A1AZR	C10-C15-C16-O2
2	A	905	MYR	C7-C8-C9-C10
2	A	901	MYR	C3-C4-C5-C6
2	A	905	MYR	C9-C10-C11-C12
2	A	904	MYR	C3-C4-C5-C6
3	A	907	A1AZR	C10-C15-C16-O3
2	A	903	MYR	C5-C6-C7-C8
3	A	907	A1AZR	C14-C15-C16-O3
2	A	906	MYR	O2-C1-C2-C3
2	A	903	MYR	O1-C1-C2-C3
2	A	906	MYR	O1-C1-C2-C3
2	A	903	MYR	O2-C1-C2-C3
3	A	907	A1AZR	C14-C15-C16-O2
2	A	901	MYR	C11-C10-C9-C8
2	A	901	MYR	O2-C1-C2-C3
2	A	906	MYR	C3-C4-C5-C6
2	A	901	MYR	C2-C3-C4-C5
2	A	901	MYR	O1-C1-C2-C3
2	A	906	MYR	C5-C6-C7-C8
2	A	905	MYR	C10-C11-C12-C13


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	905	MYR	1	0
3	A	907	A1AZR	2	0
2	A	904	MYR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	582/582 (100%)	0.90	63 (10%)	11 14	11, 23, 40, 56	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	467	THR	4.0
1	A	466	LYS	3.6
1	A	362	ALA	3.5
1	A	301	ASP	3.5
1	A	361	CYS	3.3
1	A	370	TYR	3.3
1	A	366	PRO	3.3
1	A	373	VAL	3.2
1	A	584	GLY	3.2
1	A	367	HIS	3.1
1	A	277	GLU	3.1
1	A	284	LEU	3.1
1	A	302	LEU	3.1
1	A	156	PHE	3.1
1	A	300	ALA	3.1
1	A	481	LEU	3.0
1	A	363	ALA	3.0
1	A	283	LEU	2.9
1	A	364	ALA	2.9
1	A	315	VAL	2.8
1	A	214	TRP	2.8
1	A	509	PHE	2.7
1	A	371	ALA	2.7
1	A	493	VAL	2.7
1	A	497	TYR	2.5
1	A	198	LEU	2.5
1	A	463	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	ALA	2.5
1	A	305	LEU	2.5
1	A	496	THR	2.5
1	A	299	PRO	2.5
1	A	554	PHE	2.5
1	A	513	ILE	2.4
1	A	360	CYS	2.4
1	A	108	ASP	2.4
1	A	59	ALA	2.4
1	A	516	LEU	2.3
1	A	544	LEU	2.3
1	A	372	LYS	2.3
1	A	495	GLU	2.3
1	A	374	PHE	2.3
1	A	561	ALA	2.3
1	A	356	THR	2.3
1	A	546	ALA	2.2
1	A	436	LYS	2.2
1	A	322	ALA	2.2
1	A	310	VAL	2.2
1	A	468	PRO	2.2
1	A	569	ALA	2.2
1	A	507	PHE	2.2
1	A	3	HIS	2.2
1	A	498	VAL	2.2
1	A	49	PHE	2.1
1	A	483	ASN	2.1
1	A	551	PHE	2.1
1	A	519	LYS	2.1
1	A	365	ASP	2.1
1	A	557	LYS	2.1
1	A	109	ASN	2.1
1	A	66	LEU	2.1
1	A	206	PHE	2.0
1	A	252	GLU	2.0
1	A	83	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MYR	A	901	16/16	0.79	0.18	43,47,50,51	27
2	MYR	A	902	13/16	0.80	0.21	60,61,62,62	18
2	MYR	A	906	16/16	0.80	0.22	68,70,71,72	27
2	MYR	A	904	13/16	0.82	0.21	39,41,44,45	18
3	A1AZR	A	907	54/54	0.86	0.15	48,60,68,68	36
2	MYR	A	905	16/16	0.89	0.16	57,60,64,65	27
2	MYR	A	903	14/16	0.90	0.14	38,42,46,47	19

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1AZR A 907:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.