



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:51 PM UTC

PDB ID : 3CTS / pdb_00003cts
Title : CRYSTALLOGRAPHIC REFINEMENT AND ATOMIC MODELS OF
TWO DIFFERENT FORMS OF CITRATE SYNTHASE AT 2.7 AND 1.7
ANGSTROMS RESOLUTION
Authors : Remington, S.; Wiegand, G.; Huber, R.
Deposited on : 1984-01-27
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	437	 92% 7% •

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3469 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CITRATE SYNTHASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	432	3309	2115	574	603	17	70	0	3

- Molecule 2 is COENZYME A (CCD ID: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	46	20	7	16	3	0	0

- Molecule 3 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	101	Total	O	0	0
			101	101		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: CITRATE SYNTHASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.14Å 78.25Å 58.40Å 90.00° 78.50° 90.00°	Depositor
Resolution (Å)	5.00 – 1.70	Depositor
% Data completeness (in resolution range)	(Not available) (5.00-1.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	EREF	Depositor
R, R_{free}	0.192 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3469	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	15

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	145	UNK	Mainchain
1	A	170	UNK	Mainchain
1	A	222	UNK	Mainchain
1	A	239	UNK	Mainchain
1	A	242	UNK	Mainchain
1	A	278	UNK	Mainchain
1	A	286	UNK	Mainchain
1	A	291	UNK	Mainchain
1	A	413	UNK	Mainchain
1	A	431	UNK	Mainchain
1	A	49	UNK	Mainchain
1	A	51	UNK	Mainchain
1	A	66	UNK	Mainchain
1	A	67	UNK	Mainchain
1	A	81	UNK	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3309	0	1215	8	1
2	A	46	0	27	0	0
3	A	13	0	5	0	0
4	A	101	0	0	0	1
All	All	3469	0	1247	8	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:UNK:HG1	1:A:230:UNK:HB2	1.39	1.04
1:A:346:UNK:HB1	1:A:380:UNK:HG2	1.87	0.56
1:A:285:UNK:HB1	1:A:346:UNK:CE	2.36	0.55
1:A:187:UNK:O	1:A:191:UNK:HB2	2.09	0.52
1:A:108:UNK:O	1:A:112:UNK:HG3	2.15	0.47
1:A:86:UNK:HG3	1:A:87:UNK:CD	2.45	0.46
1:A:283:UNK:O	1:A:287:UNK:HB2	2.16	0.44
1:A:103:UNK:N	1:A:106:UNK:HG2	2.36	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:UNK:CB	4:A:523:HOH:O[1_554]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	CIT	A	439	-	12,12,12	1.51	2 (16%)	17,17,17	1.53	3 (17%)
2	COA	A	438	-	46,48,50	2.13	10 (21%)	68,73,75	1.95	18 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIT	A	439	-	-	2/16/16/16	-
2	COA	A	438	-	-	7/46/62/64	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	438	COA	P3B-O3B	6.47	1.70	1.59
2	A	438	COA	C5A-N7A	-5.56	1.28	1.39
2	A	438	COA	C8A-N7A	4.94	1.41	1.31
2	A	438	COA	P2A-O3A	4.91	1.64	1.59
2	A	438	COA	C4A-N9A	-4.25	1.28	1.37
2	A	438	COA	C3P-N4P	3.23	1.51	1.45
2	A	438	COA	P3B-O7A	3.07	1.60	1.50
2	A	438	COA	P1A-O3A	3.05	1.62	1.59
3	A	439	CIT	C2-C3	2.79	1.57	1.54
2	A	438	COA	C4A-N3A	-2.63	1.29	1.34
3	A	439	CIT	C3-C6	-2.55	1.50	1.53
2	A	438	COA	O9P-C9P	2.08	1.27	1.23

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	438	COA	N9A-C8A-N7A	-6.42	104.83	113.94
2	A	438	COA	C5A-C4A-N3A	-5.36	119.34	126.72
2	A	438	COA	O4B-C1B-C2B	-4.71	96.54	106.62
2	A	438	COA	C6A-C5A-C4A	4.12	122.81	117.18
2	A	438	COA	C4A-N9A-C8A	3.82	109.75	105.74
2	A	438	COA	C5A-N7A-C8A	3.78	109.39	103.45
2	A	438	COA	N6A-C6A-N1A	3.34	125.81	118.38
2	A	438	COA	C4A-N9A-C1B	-3.09	119.41	126.63
2	A	438	COA	N3A-C4A-N9A	3.03	132.31	127.17
2	A	438	COA	C3B-C2B-C1B	2.66	105.75	99.89
2	A	438	COA	C4A-C5A-N7A	-2.60	107.61	110.58
2	A	438	COA	O5A-P2A-O3A	2.42	113.83	107.27
3	A	439	CIT	O7-C3-C2	2.39	114.83	109.38
3	A	439	CIT	C3-C4-C5	-2.35	107.50	113.92
2	A	438	COA	C5A-C4A-N9A	2.30	108.32	105.81
2	A	438	COA	CDP-CBP-CCP	2.29	112.00	108.22
2	A	438	COA	CEP-CBP-CCP	-2.27	104.48	108.22
2	A	438	COA	C2B-C3B-C4B	-2.27	99.27	103.24
2	A	438	COA	C7P-C6P-C5P	-2.24	108.66	112.39
3	A	439	CIT	C4-C3-C6	-2.24	105.09	110.03
2	A	438	COA	C4B-O4B-C1B	2.23	114.38	109.47

There are no chirality outliers.


All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	438	COA	CDP-CBP-CCP-O6A
2	A	438	COA	CEP-CBP-CCP-O6A
3	A	439	CIT	C2-C3-C6-O5
3	A	439	CIT	O7-C3-C6-O5
2	A	438	COA	P2A-O3A-P1A-O1A
2	A	438	COA	C3B-O3B-P3B-O8A
2	A	438	COA	C3B-O3B-P3B-O9A
2	A	438	COA	C3B-O3B-P3B-O7A
2	A	438	COA	N8P-C9P-CAP-OAP

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.