



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:29 AM UTC

PDB ID : 1CU6 / pdb_00001cu6
Title : T4 LYSOZYME MUTANT L91A
Authors : Gassner, N.C.; Baase, W.A.; Lindstrom, J.D.; Lu, J.; Matthews, B.W.
Deposited on : 1999-08-20
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	4% 36% 45% 15% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1289	811	235	238	5			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	THR	CYS	engineered mutation	UNP P00720
A	97	ALA	CYS	engineered mutation	UNP P00720
A	91	ALA	LEU	engineered mutation	UNP P00720

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cl	0	0
			2	2		

- Molecule 3 is 2-HYDROXYETHYL DISULFIDE (CCD ID: HED) (formula: C₄H₁₀O₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	S	
			8	4	2	2	
							0
							0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	110	Total	O		
			110	110		
					0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	60.90Å 60.90Å 97.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.10 40.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.0 (40.00-2.10) 97.2 (40.00-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.58 (at 1.66Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.174 , (Not available) 0.192 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 106.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1409	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, HED

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.13	35/1309 (2.7%)	2.32	73/1763 (4.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	18

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	87	VAL	C-O	-7.99	1.15	1.24
1	A	44	SER	CA-CB	7.72	1.65	1.53
1	A	106	MET	CA-C	-7.39	1.43	1.52
1	A	142	THR	CA-CB	7.07	1.61	1.53
1	A	4	PHE	N-CA	-6.95	1.38	1.46
1	A	154	ARG	C-N	-6.84	1.23	1.33
1	A	108	GLU	CD-OE2	6.70	1.38	1.25
1	A	94	VAL	N-CA	-6.30	1.38	1.46
1	A	78	ILE	CA-C	6.23	1.60	1.52
1	A	67	PHE	CA-C	6.16	1.60	1.52
1	A	13	LEU	CA-C	-6.00	1.45	1.52
1	A	80	ARG	NE-CZ	5.94	1.39	1.33
1	A	27	ILE	CA-C	5.91	1.59	1.52
1	A	76	ARG	CA-CB	5.90	1.62	1.53
1	A	115	THR	C-O	-5.74	1.17	1.24
1	A	83	LYS	C-O	-5.72	1.16	1.24
1	A	144	ASN	C-O	-5.69	1.17	1.24
1	A	43	LYS	C-N	-5.67	1.26	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	59	THR	N-CA	-5.62	1.39	1.45
1	A	61	ASP	CG-OD2	5.56	1.35	1.25
1	A	140	ASN	C-N	5.55	1.42	1.33
1	A	47	ASP	CA-CB	5.49	1.61	1.53
1	A	130	ALA	C-N	-5.45	1.27	1.33
1	A	106	MET	C-O	-5.44	1.16	1.24
1	A	40	ASN	N-CA	-5.33	1.40	1.46
1	A	104	PHE	CA-C	5.27	1.59	1.52
1	A	103	VAL	C-N	-5.22	1.26	1.33
1	A	115	THR	C-N	-5.22	1.26	1.33
1	A	149	VAL	N-CA	-5.19	1.40	1.46
1	A	79	LEU	C-N	-5.18	1.25	1.33
1	A	129	ALA	CA-CB	5.17	1.61	1.53
1	A	150	ILE	N-CA	-5.14	1.40	1.46
1	A	127	ASP	CG-OD2	5.11	1.35	1.25
1	A	61	ASP	N-CA	-5.10	1.40	1.46
1	A	161	TYR	N-CA	-5.06	1.39	1.46

All (73) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ARG	CD-NE-CZ	-10.93	109.10	124.40
1	A	145	ARG	CD-NE-CZ	-10.08	110.28	124.40
1	A	143	PRO	O-C-N	-8.46	112.51	122.24
1	A	82	ALA	CA-C-O	8.00	129.17	119.49
1	A	109	THR	O-C-N	7.47	129.76	122.07
1	A	148	ARG	O-C-N	7.43	129.73	122.07
1	A	2	ASN	CA-CB-CG	-7.42	105.18	112.60
1	A	76	ARG	CA-C-O	7.41	128.40	120.55
1	A	76	ARG	CB-CG-CD	7.36	128.22	111.30
1	A	120	MET	N-CA-CB	6.93	120.06	110.01
1	A	37	PRO	CB-CA-C	-6.90	100.20	111.31
1	A	109	THR	CA-CB-OG1	-6.75	99.48	109.60
1	A	104	PHE	N-CA-CB	-6.55	100.57	110.07
1	A	19	LYS	CA-C-O	-6.17	113.65	120.38
1	A	9	ILE	CB-CG1-CD1	-6.12	100.94	113.80
1	A	135	LYS	CB-CA-C	-6.08	102.55	112.03
1	A	81	ASN	OD1-CG-ND2	-6.07	116.53	122.60
1	A	47	ASP	CA-CB-CG	6.02	118.62	112.60
1	A	132	ASN	N-CA-C	-5.87	104.80	111.14
1	A	15	LEU	N-CA-C	5.86	120.27	113.18
1	A	17	ILE	CA-C-O	-5.84	114.45	121.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	ASN	CA-CB-CG	5.83	118.42	112.60
1	A	151	THR	CA-CB-OG1	-5.82	100.86	109.60
1	A	145	ARG	NE-CZ-NH2	5.82	124.44	119.20
1	A	4	PHE	CA-CB-CG	-5.81	107.99	113.80
1	A	106	MET	CA-CB-CG	5.78	125.66	114.10
1	A	18	TYR	CA-C-O	-5.76	115.06	121.51
1	A	39	LEU	CA-C-O	5.75	126.59	119.97
1	A	66	LEU	CB-CA-C	-5.74	100.02	110.63
1	A	160	ALA	CA-C-N	-5.72	113.10	122.54
1	A	160	ALA	C-N-CA	-5.72	113.10	122.54
1	A	24	TYR	N-CA-CB	5.71	118.69	110.06
1	A	73	ALA	O-C-N	-5.61	115.75	122.15
1	A	20	ASP	CB-CG-OD2	-5.60	105.51	118.40
1	A	50	ILE	CA-C-O	5.58	125.77	119.92
1	A	36	SER	CA-C-N	5.57	125.27	119.64
1	A	36	SER	C-N-CA	5.57	125.27	119.64
1	A	38	SER	N-CA-C	5.55	118.29	109.96
1	A	21	THR	OG1-CB-CG2	-5.55	98.21	109.30
1	A	133	LEU	CA-C-O	5.49	126.35	120.20
1	A	127	ASP	CA-C-N	-5.45	112.02	120.31
1	A	127	ASP	C-N-CA	-5.45	112.02	120.31
1	A	93	ALA	CB-CA-C	5.44	119.82	110.79
1	A	15	LEU	CD1-CG-CD2	-5.44	98.84	110.80
1	A	145	ARG	CA-C-O	5.43	126.53	120.82
1	A	22	GLU	CA-C-O	5.41	125.62	119.18
1	A	26	THR	CA-CB-OG1	-5.38	101.54	109.60
1	A	102	MET	N-CA-CB	5.35	118.08	110.16
1	A	120	MET	CG-SD-CE	5.32	112.61	100.90
1	A	128	GLU	CG-CD-OE1	5.29	130.57	118.40
1	A	106	MET	CA-C-O	-5.28	112.95	119.18
1	A	115	THR	CA-CB-OG1	-5.26	101.70	109.60
1	A	98	ALA	O-C-N	-5.24	116.57	122.12
1	A	95	ARG	NE-CZ-NH2	-5.22	114.50	119.20
1	A	19	LYS	CB-CA-C	-5.21	100.70	109.72
1	A	19	LYS	CA-C-N	5.20	130.33	121.29
1	A	19	LYS	C-N-CA	5.20	130.33	121.29
1	A	56	GLY	CA-C-N	-5.20	115.72	122.94
1	A	56	GLY	C-N-CA	-5.20	115.72	122.94
1	A	65	LYS	O-C-N	-5.19	116.23	122.15
1	A	53	ASN	CA-CB-CG	-5.18	107.42	112.60
1	A	109	THR	N-CA-C	5.17	116.60	111.07
1	A	20	ASP	CA-C-O	-5.15	116.60	122.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	THR	CA-CB-CG2	-5.15	101.74	110.50
1	A	106	MET	N-CA-CB	5.11	118.42	110.80
1	A	119	ARG	N-CA-C	-5.11	105.80	111.36
1	A	132	ASN	CA-C-N	-5.10	112.94	120.38
1	A	132	ASN	C-N-CA	-5.10	112.94	120.38
1	A	144	ASN	OD1-CG-ND2	-5.09	117.51	122.60
1	A	140	ASN	CA-C-O	-5.06	113.15	119.38
1	A	69	GLN	CG-CD-NE2	5.06	123.98	116.40
1	A	9	ILE	CA-C-O	5.05	126.66	121.05
1	A	160	ALA	N-CA-C	-5.00	106.44	112.54

There are no chirality outliers.

All (18) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	103	VAL	Mainchain
1	A	104	PHE	Sidechain
1	A	106	MET	Mainchain
1	A	12	GLY	Mainchain
1	A	125	ARG	Sidechain
1	A	134	ALA	Mainchain
1	A	155	THR	Mainchain
1	A	160	ALA	Mainchain
1	A	18	TYR	Sidechain
1	A	20	ASP	Mainchain
1	A	22	GLU	Sidechain
1	A	30	GLY	Mainchain
1	A	38	SER	Mainchain
1	A	72	ASP	Sidechain
1	A	79	LEU	Mainchain
1	A	84	LEU	Mainchain
1	A	88	TYR	Sidechain
1	A	89	ASP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1289	0	1313	32	1
2	A	2	0	0	0	0
3	A	8	0	10	0	1
4	A	110	0	0	4	1
All	All	1409	0	1323	32	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:HG2	1:A:158:TRP:CD2	2.25	0.72
1:A:16:LYS:HE3	1:A:57:VAL:HG23	1.78	0.64
1:A:1:MET:HG2	1:A:158:TRP:CE3	2.37	0.60
1:A:53:ASN:ND2	4:A:278:HOH:O	2.38	0.57
1:A:106:MET:HE1	1:A:138:TRP:CD1	2.42	0.55
1:A:14:ARG:O	1:A:28:GLY:N	2.38	0.54
1:A:106:MET:HE1	1:A:138:TRP:CG	2.44	0.52
1:A:16:LYS:HD2	1:A:57:VAL:CG2	2.41	0.50
1:A:46:LEU:HD23	1:A:56:GLY:HA2	1.94	0.50
1:A:19:LYS:HA	1:A:24:TYR:O	2.10	0.50
1:A:24:TYR:CE2	1:A:35:LYS:HB3	2.46	0.49
1:A:151:THR:O	1:A:155:THR:HG23	2.12	0.49
1:A:76:ARG:CB	1:A:76:ARG:NH1	2.76	0.48
1:A:11:GLU:OE2	4:A:204:HOH:O	2.20	0.47
1:A:16:LYS:HE2	1:A:17:ILE:H	1.79	0.47
1:A:106:MET:HE1	1:A:138:TRP:CD2	2.51	0.45
1:A:106:MET:CE	1:A:138:TRP:CD1	3.00	0.45
1:A:47:ASP:OD1	1:A:54:THR:N	2.33	0.44
1:A:50:ILE:HD13	1:A:50:ILE:HA	1.71	0.44
1:A:143:PRO:O	1:A:147:LYS:HG3	2.18	0.44
1:A:84:LEU:C	1:A:86:PRO:CD	2.92	0.43
1:A:114:PHE:O	1:A:115:THR:C	2.61	0.42
1:A:16:LYS:O	1:A:17:ILE:C	2.58	0.42
1:A:141:GLN:HB3	4:A:336:HOH:O	2.18	0.42
1:A:7:LEU:HD23	1:A:7:LEU:HA	1.89	0.42
1:A:100:ILE:HD13	1:A:100:ILE:HG21	1.87	0.41
1:A:67:PHE:O	1:A:68:ASN:C	2.61	0.41
1:A:76:ARG:NH1	1:A:76:ARG:CG	2.77	0.41
1:A:128:GLU:HG3	4:A:311:HOH:O	2.21	0.41
1:A:24:TYR:CD2	1:A:35:LYS:HA	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:PHE:HD2	1:A:67:PHE:CE1	2.40	0.40
1:A:106:MET:HE1	1:A:138:TRP:CE2	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:N	4:A:240:HOH:O[5_555]	1.98	0.22
3:A:170:HED:O1	3:A:170:HED:O1[5_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/164 (98%)	155 (97%)	5 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/136 (98%)	125 (93%)	9 (7%)	15	12

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	16	LYS
1	A	48	LYS
1	A	53	ASN
1	A	76	ARG
1	A	83	LYS
1	A	131	VAL
1	A	154	ARG
1	A	162	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HED	A	170	-	7,7,7	0.38	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HED	A	170	-	-	3/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	170	HED	C6-C5-S4-S3
3	A	170	HED	S4-C5-C6-O6
3	A	170	HED	C1-C2-S3-S4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	170	HED	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/164 (98%)	-0.07	6 (3%) 45 47	9, 19, 43, 62	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	76	ARG	3.5
1	A	80	ARG	2.6
1	A	55	ASN	2.3
1	A	16	LYS	2.3
1	A	162	LYS	2.2
1	A	40	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	HED	A	170	8/8	0.87	0.17	51,70,100,100	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	178	1/1	0.91	0.11	32,32,32,32	1
2	CL	A	173	1/1	0.98	0.06	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.