



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 01:27 AM UTC

PDB ID : 3CU1 / pdb_00003cu1
Title : Crystal Structure of 2:2:2 FGFR2D2:FGF1:SOS complex
Authors : Guo, F.; Dakshinamurthy, R.; Thallapuram, S.K.K.; Sakon, J.
Deposited on : 2008-04-15
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	
1	C	100	
2	B	131	
2	D	131	
3	E	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GU4	F	1	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4183 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fibroblast growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	100	832	524	157	145	6	0	3	0
1	C	100	825	520	154	145	6	0	2	0

- Molecule 2 is a protein called Heparin-binding growth factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	128	1022	646	179	193	4	0	0	0
2	D	131	1051	665	184	198	4	0	0	0

- Molecule 3 is an oligosaccharide called 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	O	S			
3	E	2	55	12	35	8	0	0	0
3	F	2	55	12	35	8	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	80	Total	O	0	0
			80	80		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	68	Total O 68 68	0	0
4	C	98	Total O 98 98	0	0
4	D	97	Total O 97 97	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fibroblast growth factor receptor 2




- Molecule 1: Fibroblast growth factor receptor 2





- Molecule 2: Heparin-binding growth factor 1



- Molecule 2: Heparin-binding growth factor 1





- Molecule 3: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain E:  100%



- Molecule 3: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain F:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.93Å 110.38Å 74.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.95 – 2.60 42.95 – 2.60	Depositor EDS
% Data completeness (in resolution range)	90.0 (42.95-2.60) 90.0 (42.95-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.24 (at 2.61Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.180 , 0.277 0.261 , 0.317	Depositor DCC
R_{free} test set	1044 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	52.0	Xtrriage
Anisotropy	0.186	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4183	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GU4, YYJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.07	105/868 (12.1%)	1.80	21/1173 (1.8%)
1	C	3.13	87/857 (10.2%)	1.83	23/1159 (2.0%)
2	B	2.55	66/1045 (6.3%)	1.74	26/1410 (1.8%)
2	D	2.86	93/1075 (8.7%)	1.88	37/1450 (2.6%)
All	All	2.89	351/3845 (9.1%)	1.81	107/5192 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	2	0

The worst 5 of 351 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	171	ALA	CA-CB	-12.61	1.32	1.53
1	C	158	ASN	CA-CB	-12.01	1.44	1.53
1	A	247	ASP	CA-C	11.71	1.66	1.52
2	B	91	GLU	C-O	-10.70	1.10	1.24
1	A	242	HIS	CA-C	-10.64	1.39	1.52

The worst 5 of 107 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	91	GLU	N-CA-C	15.05	131.90	112.34
2	B	92	ASN	N-CA-C	13.81	130.84	113.43
2	D	91	GLU	CA-C-N	12.23	143.71	121.70
2	D	91	GLU	C-N-CA	12.23	143.71	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	186	MET	CA-C-N	-10.53	110.10	120.52

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	D	91	GLU	CA
2	D	92	ASN	CA

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	832	0	810	64	0
1	C	825	0	801	89	0
2	B	1022	0	1007	93	0
2	D	1051	0	1035	138	0
3	E	55	0	6	4	0
3	F	55	0	6	16	0
4	A	80	0	0	26	0
4	B	68	0	0	32	0
4	C	98	0	0	33	0
4	D	97	0	0	47	0
All	All	4183	0	3665	390	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 52.

The worst 5 of 390 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (\AA)	Clash overlap (\AA)
1:C:197:GLU:CG	1:C:197:GLU:CB	1.74	1.63
2:D:26:LEU:CD2	2:D:26:LEU:CG	1.82	1.57
2:D:128:LYS:CE	2:D:128:LYS:NZ	1.68	1.57
2:D:128:LYS:CE	2:D:128:LYS:CD	1.77	1.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:10:LYS:CG	2:D:11:PRO:HD3	1.07	1.51

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	101/100 (101%)	98 (97%)	3 (3%)	0	100	100
1	C	100/100 (100%)	93 (93%)	7 (7%)	0	100	100
2	B	126/131 (96%)	113 (90%)	11 (9%)	2 (2%)	7	16
2	D	129/131 (98%)	114 (88%)	9 (7%)	6 (5%)	2	2
All	All	456/462 (99%)	418 (92%)	30 (7%)	8 (2%)	6	14

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	10	LYS
2	D	92	ASN
2	B	52	GLY
2	D	50	SER
2	D	80	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/88 (103%)	82 (90%)	9 (10%)	7	16
1	C	90/88 (102%)	85 (94%)	5 (6%)	19	40
2	B	112/115 (97%)	98 (88%)	14 (12%)	4	9
2	D	115/115 (100%)	96 (84%)	19 (16%)	2	4
All	All	408/406 (100%)	361 (88%)	47 (12%)	5	11

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	9	LYS
2	D	49	GLU
2	D	10	LYS
2	D	32	ASP
2	D	54	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	92	ASN
2	D	80	ASN
1	C	173	ASN
2	D	45	GLN
1	C	167	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GU4	E	1	3	27,27,28	1.56	6 (22%)	32,43,45	2.19	10 (31%)
3	YYJ	E	2	3	27,28,28	2.79	8 (29%)	33,46,46	2.89	12 (36%)
3	GU4	F	1	3	27,27,28	2.37	9 (33%)	32,43,45	3.28	11 (34%)
3	YYJ	F	2	3	27,28,28	2.23	7 (25%)	33,46,46	2.20	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GU4	E	1	3	-	11/21/38/41	0/1/1/1
3	YYJ	E	2	3	-	10/23/42/42	0/1/1/1
3	GU4	F	1	3	-	16/21/38/41	0/1/1/1
3	YYJ	F	2	3	-	13/23/42/42	0/1/1/1

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	YYJ	O3-S3	6.43	1.76	1.57
3	E	2	YYJ	O6-S6	6.20	1.73	1.56
3	E	2	YYJ	O1-S1	6.18	1.73	1.56
3	F	1	GU4	O2-S2	5.75	1.74	1.57
3	F	2	YYJ	O6-S6	5.42	1.71	1.56

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	GU4	O2-C2-C3	14.70	122.93	106.65
3	E	2	YYJ	C3-O3-S3	12.38	133.00	117.69
3	F	2	YYJ	C3-O3-S3	8.55	128.26	117.69
3	E	1	GU4	O2-C2-C3	7.81	115.29	106.65
3	E	2	YYJ	C4-O4-S4	4.65	130.19	119.04

There are no chirality outliers.

5 of 50 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	1	GU4	C5-C4-O4-S4
3	E	1	GU4	C1-C2-O2-S2
3	E	1	GU4	C2-O2-S2-O12
3	E	2	YYJ	O1-C1-C2-C3
3	E	2	YYJ	O1-C1-C2-O2


There are no ring outliers.

4 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	1	GU4	11	0
3	E	1	GU4	3	0
3	F	2	YYJ	5	0
3	E	2	YYJ	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/100 (100%)	0.19	4 (4%) 42 37	4, 9, 15, 29	3 (3%)
1	C	100/100 (100%)	0.33	4 (4%) 42 37	3, 9, 15, 23	2 (2%)
2	B	128/131 (97%)	0.21	3 (2%) 61 55	2, 10, 17, 21	1 (0%)
2	D	131/131 (100%)	0.25	6 (4%) 37 32	5, 11, 17, 22	1 (0%)
All	All	459/462 (99%)	0.24	17 (3%) 45 39	2, 10, 17, 29	7 (1%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	9	LYS	4.1
1	A	212[A]	GLN	3.6
1	C	201	GLU	3.5
2	B	49	GLU	3.5
2	D	7	ASN	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	YYJ	F	2	28/28	0.74	0.17	101,113,114,115	0
3	YYJ	E	2	28/28	0.79	0.17	102,113,115,115	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GU4	F	1	27/28	0.86	0.21	71,83,88,92	0
3	GU4	E	1	27/28	0.90	0.18	73,85,90,94	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.