



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 09:15 AM EDT

PDB ID : 5CUY / pdb_00005cuy
Title : Crystal structure of Trypanosoma brucei Vacuolar Soluble Pyrophosphatases
in apo form
Authors : Yang, Y.Y.; Ko, T.P.; Liu, W.D.; Zheng, Y.Y.; Chen, C.C.; Guo, R.T.
Deposited on : 2015-07-25
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	
1	B	414	
1	C	414	
1	D	414	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acidocalcisomal pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	392	3157	2014	537	583	23	0	0	0
1	B	388	3124	1990	534	577	23	0	0	0
1	C	392	3161	2016	541	581	23	0	0	0
1	D	396	3190	2033	546	588	23	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	406	ARG	LEU	engineered mutation	UNP Q384W3
B	406	ARG	LEU	engineered mutation	UNP Q384W3
C	406	ARG	LEU	engineered mutation	UNP Q384W3
D	406	ARG	LEU	engineered mutation	UNP Q384W3

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 13 6 7	0	0
2	B	1	Total C O 13 6 7	0	0
2	C	1	Total C O 13 6 7	0	0
2	D	1	Total C O 13 6 7	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	164	Total O 164 164	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	183	Total 183	O 183	0	0
4	C	171	Total 171	O 171	0	0
4	D	168	Total 168	O 168	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acidocalcisomal pyrophosphatase






● Molecule 1: Acidocalcisomal pyrophosphatase



● Molecule 1: Acidocalcisomal pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	199.92Å 70.14Å 141.76Å 90.00° 106.38° 90.00°	Depositor
Resolution (Å)	25.00 – 2.50 25.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	94.2 (25.00-2.50) 94.1 (25.00-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.16 (at 2.50Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.194 , 0.252 0.189 , 0.243	Depositor DCC
R_{free} test set	3117 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtrriage
Anisotropy	0.866	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13374	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.86	2/3245 (0.1%)	1.24	29/4395 (0.7%)
1	B	0.93	2/3210 (0.1%)	1.26	30/4344 (0.7%)
1	C	0.86	1/3249 (0.0%)	1.21	26/4398 (0.6%)
1	D	0.90	3/3278 (0.1%)	1.24	29/4437 (0.7%)
All	All	0.89	8/12982 (0.1%)	1.24	114/17574 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	302	MET	SD-CE	7.19	1.97	1.79
1	A	29	VAL	CA-CB	6.03	1.62	1.54
1	D	73	THR	CA-CB	-5.56	1.44	1.53
1	A	394	VAL	CA-CB	5.42	1.61	1.54
1	B	331	VAL	CA-CB	-5.20	1.47	1.54

The worst 5 of 114 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	19	LEU	N-CA-C	-11.45	96.53	112.45
1	A	300	ILE	N-CA-C	10.66	122.32	112.29
1	C	300	ILE	N-CA-C	10.00	121.69	112.29
1	B	367	VAL	N-CA-C	9.90	122.71	113.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	157	ALA	N-CA-C	-9.69	98.03	110.53

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	185	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3157	0	3038	145	0
1	B	3124	0	3006	158	0
1	C	3161	0	3049	143	0
1	D	3190	0	3075	145	0
2	A	13	0	5	1	0
2	B	13	0	5	3	0
2	C	13	0	5	1	0
2	D	13	0	5	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	164	0	0	3	0
4	B	183	0	0	11	0
4	C	171	0	0	4	0
4	D	168	0	0	8	0
All	All	13374	0	12188	533	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 533 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:279:SER:H	1:D:401:MET:HE2	1.06	1.17
1:B:322:ILE:HG12	1:B:327:MET:HE1	1.19	1.12
1:C:279:SER:H	1:C:401:MET:HE2	1.06	1.12
1:A:340:ILE:HG21	1:D:302:MET:HE1	1.34	1.08
1:B:289:THR:HG22	1:B:290:GLY:H	1.12	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/414 (94%)	361 (93%)	24 (6%)	5 (1%)	9	18
1	B	384/414 (93%)	351 (91%)	27 (7%)	6 (2%)	7	14
1	C	390/414 (94%)	358 (92%)	22 (6%)	10 (3%)	4	7
1	D	394/414 (95%)	365 (93%)	23 (6%)	6 (2%)	8	16
All	All	1558/1656 (94%)	1435 (92%)	96 (6%)	27 (2%)	7	13

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	ARG
1	B	148	ARG
1	C	92	HIS
1	C	148	ARG
1	C	149	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/361 (94%)	313 (92%)	28 (8%)	10	23
1	B	337/361 (93%)	308 (91%)	29 (9%)	10	21
1	C	341/361 (94%)	312 (92%)	29 (8%)	10	22
1	D	344/361 (95%)	315 (92%)	29 (8%)	10	22
All	All	1363/1444 (94%)	1248 (92%)	115 (8%)	10	22

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	24	VAL
1	D	295	VAL
1	C	148	ARG
1	D	289	THR
1	D	146	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	166	GLN
1	D	269	ASN
1	D	404	ASN
1	B	166	GLN
1	B	146	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	D	901	-	12,12,12	0.89	0	17,17,17	1.47	2 (11%)
2	CIT	B	901	-	12,12,12	0.85	0	17,17,17	1.41	1 (5%)
2	CIT	A	901	-	12,12,12	0.98	0	17,17,17	1.40	1 (5%)
2	CIT	C	901	-	12,12,12	1.17	0	17,17,17	1.44	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	D	901	-	-	0/16/16/16	-
2	CIT	B	901	-	-	0/16/16/16	-
2	CIT	A	901	-	-	0/16/16/16	-
2	CIT	C	901	-	-	0/16/16/16	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	CIT	O6-C6-C3	3.90	120.62	113.14
2	C	901	CIT	O6-C6-C3	3.72	120.28	113.14
2	A	901	CIT	O6-C6-C3	3.71	120.25	113.14
2	D	901	CIT	O6-C6-C3	3.62	120.09	113.14
2	D	901	CIT	O7-C3-C6	2.16	112.02	108.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	901	CIT	2	0
2	B	901	CIT	3	0
2	A	901	CIT	1	0
2	C	901	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/414 (94%)	-0.25	6 (1%) 72 68	25, 48, 86, 106	0
1	B	388/414 (93%)	-0.36	9 (2%) 61 57	27, 42, 86, 111	0
1	C	392/414 (94%)	-0.27	4 (1%) 79 76	26, 47, 95, 119	0
1	D	396/414 (95%)	-0.30	5 (1%) 75 71	26, 44, 87, 106	0
All	All	1568/1656 (94%)	-0.29	24 (1%) 72 68	25, 45, 90, 119	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	149	THR	4.8
1	A	147	SER	3.9
1	A	149	THR	3.8
1	C	145	VAL	3.4
1	B	152	PHE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CIT	C	901	13/13	0.77	0.17	108,108,110,111	0
2	CIT	D	901	13/13	0.81	0.17	104,106,106,106	0
2	CIT	A	901	13/13	0.84	0.16	107,108,111,111	0
3	MG	D	902	1/1	0.85	0.13	49,49,49,49	0
2	CIT	B	901	13/13	0.88	0.14	87,90,93,93	0
3	MG	C	902	1/1	0.93	0.15	49,49,49,49	0
3	MG	B	902	1/1	0.93	0.04	59,59,59,59	0
3	MG	A	902	1/1	0.94	0.13	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.