



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:26 AM UTC

PDB ID : 6CU0 / pdb\_00006cu0  
Title : Crystal structure of 4-1BBL/4-1BB (C121S) complex in P21 space group  
Authors : Aruna, B.; Zajonc, D.M.; Doukov, T.  
Deposited on : 2018-03-23  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	<div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 1%, green 1%, green 79%, yellow 79%, yellow 88%, orange 88%, orange 91%, red 91%, red 95%, grey 95%);"></div> <div style="margin-left: 5px;">%</div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>79%</span> <span>12%</span> <span>•• 8%</span> </div>
1	B	165	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, green 1%, green 80%, yellow 80%, yellow 88%, orange 88%, orange 91%, red 91%, red 95%, grey 95%);"></div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>80%</span> <span>8%</span> <span>• 11%</span> </div>
1	C	165	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, green 1%, green 79%, yellow 79%, yellow 88%, orange 88%, orange 91%, red 91%, red 95%, grey 95%);"></div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>79%</span> <span>8%</span> <span>• 11%</span> </div>
1	D	165	<div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 1%, green 1%, green 78%, yellow 78%, yellow 88%, orange 88%, orange 91%, red 91%, red 95%, grey 95%);"></div> <div style="margin-left: 5px;">%</div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>78%</span> <span>11%</span> <span>• 11%</span> </div>
1	E	165	<div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 1%, green 1%, green 78%, yellow 78%, yellow 88%, orange 88%, orange 91%, red 91%, red 95%, grey 95%);"></div> <div style="margin-left: 5px;">%</div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>78%</span> <span>8%</span> <span>• 13%</span> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	165	<p>% 77% 14% 8%</p>
2	G	138	<p>88% 7% 5%</p>
2	H	138	<p>% 90% 6%</p>
2	I	138	<p>89% 6% 5%</p>
2	J	138	<p>% 87% 7% 7%</p>
2	K	138	<p>% 91%</p>
2	L	138	<p>2% 86% 7% 7%</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	152	1072	692	185	195	0	0	0	
1	B	147	1033	666	179	187	1	0	0	
1	C	147	999	644	169	186	0	0	0	
1	D	147	1023	656	177	189	1	0	0	
1	E	144	987	631	171	184	1	0	0	
1	F	151	1046	672	182	191	1	0	0	

- Molecule 2 is a protein called Tumor necrosis factor receptor superfamily member 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	G	131	880	517	158	184	21	0	0	0
2	H	132	899	527	161	190	21	0	0	0
2	J	129	865	507	154	183	21	0	0	0
2	K	132	868	507	158	182	21	0	0	0
2	L	129	863	506	153	184	20	0	0	0
2	I	131	869	511	151	186	21	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	121	SER	CYS	engineered mutation	UNP Q07011

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	121	SER	CYS	engineered mutation	UNP Q07011
J	121	SER	CYS	engineered mutation	UNP Q07011
K	121	SER	CYS	engineered mutation	UNP Q07011
L	121	SER	CYS	engineered mutation	UNP Q07011
I	121	SER	CYS	engineered mutation	UNP Q07011

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	G	1	Total	C	N	O	0	0
			14	8	1	5		
3	H	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	2	Total	O	0	0
			2	2		
4	C	1	Total	O	0	0
			1	1		
4	E	1	Total	O	0	0
			1	1		

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	F	1	Total O 1 1	0	0
4	G	7	Total O 7 7	0	0
4	H	3	Total O 3 3	0	0
4	J	5	Total O 5 5	0	0
4	K	13	Total O 13 13	0	0
4	L	2	Total O 2 2	0	0
4	I	4	Total O 4 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor ligand superfamily member 9




- Molecule 1: Tumor necrosis factor ligand superfamily member 9




- Molecule 1: Tumor necrosis factor ligand superfamily member 9



- Molecule 1: Tumor necrosis factor ligand superfamily member 9




- Molecule 1: Tumor necrosis factor ligand superfamily member 9




GLU  
TLE

- Molecule 1: Tumor necrosis factor ligand superfamily member 9

Chain F: 


GLU  
TLE

- Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain G: 





- Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain H: 




- Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain J: 




- Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain K: 




- Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain L: 



- Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain I:  89% 6% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.96Å 114.75Å 126.04Å 90.00° 101.24° 90.00°	Depositor
Resolution (Å)	50.00 – 3.20 50.00 – 3.20	Depositor EDS
% Data completeness (in resolution range)	94.6 (50.00-3.20) 94.6 (50.00-3.20)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, $R_{free}$	0.247 , 0.296 0.258 , 0.302	Depositor DCC
$R_{free}$ test set	1668 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	126.3	Xtrriage
Anisotropy	0.329	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 164.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	11476	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	160.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/1095	0.91	2/1501 (0.1%)
1	B	0.59	0/1054	0.78	0/1443
1	C	0.59	0/1017	0.78	1/1398 (0.1%)
1	D	0.64	0/1043	0.77	0/1428
1	E	0.63	0/1006	0.80	0/1377
1	F	0.57	0/1068	0.81	1/1464 (0.1%)
2	G	0.68	0/894	0.80	0/1214
2	H	0.66	0/913	0.82	0/1239
2	I	0.69	0/883	0.82	0/1204
2	J	0.61	0/879	0.83	0/1193
2	K	0.62	0/880	0.78	0/1195
2	L	0.71	0/875	0.85	1/1187 (0.1%)
All	All	0.64	0/11607	0.81	5/15843 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	PRO	CA-C-O	-12.78	111.85	121.31
1	F	192	ALA	N-CA-C	6.43	120.29	111.54
1	C	190	SER	N-CA-C	-6.18	104.54	111.28
1	A	192	ALA	N-CA-C	5.10	121.67	110.80
2	L	149	ASN	N-CA-C	5.06	116.80	111.28

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1029	11	0
1	B	1033	0	990	7	0
1	C	999	0	943	6	0
1	D	1023	0	956	10	0
1	E	987	0	909	7	0
1	F	1046	0	986	12	0
2	G	880	0	701	5	0
2	H	899	0	717	3	0
2	I	869	0	669	4	0
2	J	865	0	675	3	0
2	K	868	0	677	5	0
2	L	863	0	676	5	0
3	G	14	0	13	0	0
3	H	14	0	13	0	0
4	A	5	0	0	1	0
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	7	0	0	1	0
4	H	3	0	0	1	0
4	I	4	0	0	0	0
4	J	5	0	0	1	0
4	K	13	0	0	3	0
4	L	2	0	0	0	0
All	All	11476	0	9954	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 64 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:149:ASN:OD1	2:J:150:GLY:N	1.94	1.00
2:L:146:VAL:HA	2:L:158:CYS:SG	2.01	1.00

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:SER:O	1:A:192:ALA:HB2	1.87	0.73
1:D:139:GLY:HA3	1:D:241:THR:HG21	1.69	0.73
1:D:151:ARG:HE	1:D:194:ASN:HD22	1.37	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/165 (91%)	142 (95%)	7 (5%)	1 (1%)	18	52
1	B	143/165 (87%)	139 (97%)	4 (3%)	0	100	100
1	C	143/165 (87%)	136 (95%)	7 (5%)	0	100	100
1	D	143/165 (87%)	136 (95%)	6 (4%)	1 (1%)	18	52
1	E	138/165 (84%)	133 (96%)	5 (4%)	0	100	100
1	F	147/165 (89%)	143 (97%)	4 (3%)	0	100	100
2	G	127/138 (92%)	116 (91%)	11 (9%)	0	100	100
2	H	128/138 (93%)	117 (91%)	11 (9%)	0	100	100
2	I	127/138 (92%)	116 (91%)	11 (9%)	0	100	100
2	J	125/138 (91%)	116 (93%)	9 (7%)	0	100	100
2	K	126/138 (91%)	115 (91%)	11 (9%)	0	100	100
2	L	125/138 (91%)	114 (91%)	11 (9%)	0	100	100
All	All	1622/1818 (89%)	1523 (94%)	97 (6%)	2 (0%)	48	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	191	GLU

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	192	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/125 (78%)	91 (93%)	7 (7%)	13	44
1	B	94/125 (75%)	88 (94%)	6 (6%)	16	48
1	C	89/125 (71%)	81 (91%)	8 (9%)	9	34
1	D	91/125 (73%)	85 (93%)	6 (7%)	15	47
1	E	86/125 (69%)	81 (94%)	5 (6%)	18	51
1	F	94/125 (75%)	87 (93%)	7 (7%)	13	43
2	G	91/120 (76%)	87 (96%)	4 (4%)	25	59
2	H	94/120 (78%)	90 (96%)	4 (4%)	26	59
2	I	89/120 (74%)	86 (97%)	3 (3%)	32	64
2	J	88/120 (73%)	84 (96%)	4 (4%)	24	58
2	K	87/120 (72%)	86 (99%)	1 (1%)	65	79
2	L	87/120 (72%)	85 (98%)	2 (2%)	44	70
All	All	1088/1470 (74%)	1031 (95%)	57 (5%)	21	54

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	119	SER
2	I	153	GLU
1	F	162	SER
2	I	108	GLN
2	J	153	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	210	GLN
2	L	59	GLN
2	G	59	GLN
2	J	138	ASN
2	I	67	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	H	501	2	14,14,15	0.82	1 (7%)	17,19,21	1.30	3 (17%)
3	NAG	G	500	2	14,14,15	0.39	0	17,19,21	1.44	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	H	501	2	-	0/6/23/26	0/1/1/1
3	NAG	G	500	2	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	501	NAG	C1-C2	2.66	1.56	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	500	NAG	C1-O5-C5	3.33	116.65	112.19
3	H	501	NAG	C1-C2-N2	3.07	115.27	110.43
3	H	501	NAG	C1-O5-C5	2.76	115.89	112.19
3	G	500	NAG	C4-C3-C2	-2.45	107.42	111.02
3	G	500	NAG	O5-C1-C2	-2.39	107.59	111.29

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	500	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	152/165 (92%)	-0.46	1 (0%) 84 69	88, 121, 179, 235	0
1	B	147/165 (89%)	-0.38	0 100 100	98, 144, 191, 224	0
1	C	147/165 (89%)	-0.37	0 100 100	109, 139, 179, 189	0
1	D	147/165 (89%)	-0.38	1 (0%) 84 69	100, 132, 175, 211	0
1	E	144/165 (87%)	-0.26	2 (1%) 73 54	107, 143, 180, 205	0
1	F	151/165 (91%)	-0.43	1 (0%) 84 69	112, 156, 206, 244	0
2	G	131/138 (94%)	-0.32	0 100 100	104, 149, 199, 226	0
2	H	132/138 (95%)	-0.34	1 (0%) 82 67	30, 156, 238, 249	0
2	I	131/138 (94%)	-0.38	0 100 100	120, 156, 268, 313	0
2	J	129/138 (93%)	-0.22	2 (1%) 70 51	30, 198, 234, 249	0
2	K	132/138 (95%)	-0.15	1 (0%) 82 67	183, 231, 275, 298	0
2	L	129/138 (93%)	-0.19	3 (2%) 61 41	30, 167, 224, 235	0
All	All	1672/1818 (91%)	-0.33	12 (0%) 84 69	30, 154, 237, 313	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	149	ASN	4.6
1	E	165	LEU	4.5
2	H	59	GLN	4.4
1	E	147	LEU	3.6
2	J	146	VAL	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	H	501	14/15	0.81	0.07	186,195,202,204	0
3	NAG	G	500	14/15	0.90	0.08	190,198,201,201	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.