



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:51 PM UTC

PDB ID : 6CUT / pdb_00006cut
Title : Engineered Holo TrpB from *Pyrococcus furiosus*, PfTrpB7E6 with (2S,3S)-iso propylserine bound as the external aldimine
Authors : Boville, C.E.; Scheele, R.A.; Buller, A.R.; Arnold, F.H.
Deposited on : 2018-03-26
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	
1	B	388	
1	C	388	
1	D	388	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan synthase beta chain 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	2860	1822	492	534	12	0	0	0
1	B	385	2974	1897	517	546	14	0	12	0
1	C	377	2768	1772	476	509	11	0	0	0
1	D	385	2877	1833	496	536	12	0	0	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	16	VAL	ILE	engineered mutation	UNP Q8U093
A	17	GLY	GLU	engineered mutation	UNP Q8U093
A	91	PRO	LEU	engineered mutation	UNP Q8U093
A	95	LEU	PHE	engineered mutation	UNP Q8U093
A	161	ALA	LEU	engineered mutation	UNP Q8U093
A	173	GLU	VAL	engineered mutation	UNP Q8U093
A	274	LEU	PHE	engineered mutation	UNP Q8U093
A	292	SER	THR	engineered mutation	UNP Q8U093
A	384	ALA	VAL	engineered mutation	UNP Q8U093
B	16	VAL	ILE	engineered mutation	UNP Q8U093
B	17	GLY	GLU	engineered mutation	UNP Q8U093
B	91	PRO	LEU	engineered mutation	UNP Q8U093
B	95	LEU	PHE	engineered mutation	UNP Q8U093
B	161	ALA	LEU	engineered mutation	UNP Q8U093
B	173	GLU	VAL	engineered mutation	UNP Q8U093
B	274	LEU	PHE	engineered mutation	UNP Q8U093
B	292	SER	THR	engineered mutation	UNP Q8U093
B	384	ALA	VAL	engineered mutation	UNP Q8U093
C	16	VAL	ILE	engineered mutation	UNP Q8U093
C	17	GLY	GLU	engineered mutation	UNP Q8U093
C	91	PRO	LEU	engineered mutation	UNP Q8U093

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	95	LEU	PHE	engineered mutation	UNP Q8U093
C	161	ALA	LEU	engineered mutation	UNP Q8U093
C	173	GLU	VAL	engineered mutation	UNP Q8U093
C	274	LEU	PHE	engineered mutation	UNP Q8U093
C	292	SER	THR	engineered mutation	UNP Q8U093
C	384	ALA	VAL	engineered mutation	UNP Q8U093
D	16	VAL	ILE	engineered mutation	UNP Q8U093
D	17	GLY	GLU	engineered mutation	UNP Q8U093
D	91	PRO	LEU	engineered mutation	UNP Q8U093
D	95	LEU	PHE	engineered mutation	UNP Q8U093
D	161	ALA	LEU	engineered mutation	UNP Q8U093
D	173	GLU	VAL	engineered mutation	UNP Q8U093
D	274	LEU	PHE	engineered mutation	UNP Q8U093
D	292	SER	THR	engineered mutation	UNP Q8U093
D	384	ALA	VAL	engineered mutation	UNP Q8U093

- Molecule 2 is (2S,3S)-3-hydroxy-2-[(E)-({3-hydroxy-2-methyl-5-[(phosphonoxy)methyl]pyridin-4-yl}methylidene)amino]-4-methylpentanoic acid (non-preferred name) (CCD ID: FEJ) (formula: C₁₄H₂₁N₂O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			25	14	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			25	14	2	8	1		
2	C	1	Total	C	N	O	P	0	0
			25	14	2	8	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	P	0	0
			25	14	2	8	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	B	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	D	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	85	Total	O	0	0
			85	85		
4	B	112	Total	O	0	0
			112	112		
4	C	85	Total	O	0	0
			85	85		
4	D	107	Total	O	0	0
			107	107		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tryptophan synthase beta chain 1

Chain A: 




- Molecule 1: Tryptophan synthase beta chain 1

Chain B: 



- Molecule 1: Tryptophan synthase beta chain 1

Chain C: 



- Molecule 1: Tryptophan synthase beta chain 1

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.22Å 107.43Å 159.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.77 40.00 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.5 (40.00-1.77) 99.5 (40.00-1.77)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 1.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0218	Depositor
R, R_{free}	0.193 , 0.225 (Not available) , 0.226	Depositor DCC
R_{free} test set	6812 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtrriage
Anisotropy	0.255	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11972	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, FEJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2920	0.70	0/3964
1	B	0.44	0/3035	0.69	0/4108
1	C	0.45	0/2827	0.73	0/3842
1	D	0.45	0/2937	0.69	0/3983
All	All	0.44	0/11719	0.70	0/15897

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2860	0	2778	5	0
1	B	2974	0	2931	13	0
1	C	2768	0	2671	3	0
1	D	2877	0	2810	2	0
2	A	25	0	0	0	0
2	B	25	0	0	0	0
2	C	25	0	0	0	0
2	D	25	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	85	0	0	0	0
4	B	112	0	0	0	0
4	C	85	0	0	0	0
4	D	107	0	0	0	0
All	All	11972	0	11190	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137[B]:GLN:HB3	1:B:140:ASN:HD22	1.59	0.66
1:B:136[B]:ARG:O	1:B:137[B]:GLN:HG2	2.02	0.59
1:B:136[B]:ARG:C	1:B:137[B]:GLN:HG2	2.28	0.59
1:D:15:LEU:HD21	1:D:173:GLU:HA	1.90	0.54
1:B:131[B]:ALA:O	1:B:134[B]:VAL:HB	2.09	0.53
1:A:211:ILE:HG21	1:A:219:PRO:HD3	1.92	0.52
1:B:211:ILE:HG21	1:B:219:PRO:HD3	1.92	0.51
1:B:63:ILE:HD11	1:B:354:MET:HG2	1.94	0.48
1:B:134[B]:VAL:O	1:B:138[B]:LYS:HA	2.14	0.47
1:A:119:ALA:O	1:B:58:ARG:NH2	2.46	0.47
1:C:188:GLY:HA2	1:C:275:HIS:O	2.16	0.46
1:C:18:PRO:HG3	1:C:176:PHE:CD1	2.52	0.45
1:A:188:GLY:HA2	1:A:275:HIS:O	2.18	0.44
1:D:211:ILE:HG21	1:D:219:PRO:HD3	1.98	0.44
1:B:274:LEU:CD2	1:B:289:ILE:HD12	2.49	0.43
1:A:90:ALA:HB3	1:A:91:PRO:HD3	2.00	0.42
1:B:154:VAL:HG13	1:B:167:GLU:HG3	2.02	0.41
1:C:106:GLY:O	1:C:133:ASP:HB3	2.19	0.41
1:B:274:LEU:HD22	1:B:289:ILE:HD12	2.03	0.41
1:A:63:ILE:HD11	1:A:354:MET:HG2	2.03	0.40
1:B:110:HIS:CE1	1:B:184:GLY:HA2	2.57	0.40
1:B:289:ILE:HD11	1:B:302:PRO:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/388 (99%)	377 (98%)	6 (2%)	0	100	100
1	B	395/388 (102%)	385 (98%)	9 (2%)	1 (0%)	36	22
1	C	373/388 (96%)	369 (99%)	3 (1%)	1 (0%)	36	22
1	D	383/388 (99%)	378 (99%)	4 (1%)	1 (0%)	36	22
All	All	1534/1552 (99%)	1509 (98%)	22 (1%)	3 (0%)	43	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	186	VAL
1	D	186	VAL
1	C	186	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/305 (92%)	278 (100%)	1 (0%)	84	78
1	B	291/305 (95%)	287 (99%)	4 (1%)	59	44
1	C	263/305 (86%)	261 (99%)	2 (1%)	73	63
1	D	282/305 (92%)	280 (99%)	2 (1%)	76	66
All	All	1115/1220 (91%)	1106 (99%)	9 (1%)	76	63

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	322	VAL
1	B	137[A]	GLN
1	B	137[B]	GLN
1	B	290	LYS
1	B	322	VAL
1	C	267	ASN
1	C	322	VAL
1	D	13	GLU
1	D	322	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	109	GLN
1	A	155	ASN
1	B	140	ASN
1	D	283	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FEJ	B	401	-	23,25,25	2.09	5 (21%)	31,36,36	1.47	5 (16%)
2	FEJ	C	401	-	23,25,25	2.11	5 (21%)	31,36,36	1.41	2 (6%)
2	FEJ	A	401	-	23,25,25	2.12	6 (26%)	31,36,36	1.38	2 (6%)
2	FEJ	D	401	-	23,25,25	2.19	6 (26%)	31,36,36	1.44	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FEJ	B	401	-	-	6/23/23/23	0/1/1/1
2	FEJ	C	401	-	-	6/23/23/23	0/1/1/1
2	FEJ	A	401	-	-	7/23/23/23	0/1/1/1
2	FEJ	D	401	-	-	5/23/23/23	0/1/1/1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FEJ	CAA-CAS	-6.45	1.40	1.50
2	D	401	FEJ	CAA-CAS	-6.28	1.40	1.50
2	C	401	FEJ	CAA-CAS	-6.25	1.40	1.50
2	B	401	FEJ	CAA-CAS	-6.17	1.40	1.50
2	B	401	FEJ	CAL-N	4.40	1.35	1.27
2	C	401	FEJ	CAL-N	4.12	1.35	1.27
2	A	401	FEJ	CAL-N	3.89	1.34	1.27
2	C	401	FEJ	CAB-CAM	-3.89	1.40	1.50
2	D	401	FEJ	CAB-CAM	-3.79	1.41	1.50
2	B	401	FEJ	CAB-CAM	-3.67	1.41	1.50
2	D	401	FEJ	CAL-N	3.65	1.34	1.27
2	A	401	FEJ	CAB-CAM	-3.59	1.41	1.50
2	D	401	FEJ	PAP-OAQ	3.23	1.60	1.50
2	B	401	FEJ	CAO-NAR	2.98	1.40	1.34
2	C	401	FEJ	CAO-NAR	2.96	1.40	1.34
2	A	401	FEJ	CAO-NAR	2.95	1.40	1.34
2	D	401	FEJ	CAO-NAR	2.94	1.40	1.34
2	D	401	FEJ	CAN-CAL	-2.75	1.40	1.46
2	C	401	FEJ	CAN-CAL	-2.55	1.41	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FEJ	CAN-CAL	-2.52	1.41	1.46
2	A	401	FEJ	CG-CB	2.29	1.56	1.53
2	B	401	FEJ	CAN-CAL	-2.26	1.41	1.46

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	FEJ	CAN-CAT-CAS	5.22	123.08	120.14
2	C	401	FEJ	CAN-CAT-CAS	5.08	123.00	120.14
2	A	401	FEJ	CAN-CAT-CAS	5.01	122.96	120.14
2	D	401	FEJ	CAN-CAT-CAS	4.98	122.94	120.14
2	D	401	FEJ	OAE-PAP-OAD	3.06	119.28	107.80
2	A	401	FEJ	OAE-PAP-OAQ	2.18	119.32	110.83
2	B	401	FEJ	CAT-CAN-CAM	-2.15	116.55	118.28
2	B	401	FEJ	CAA-CAS-CAT	2.15	123.31	120.80
2	B	401	FEJ	OAD-PAP-OAC	-2.13	101.11	106.67
2	D	401	FEJ	CB-CA-N	2.13	115.74	109.23
2	D	401	FEJ	CAA-CAS-CAT	2.12	123.27	120.80
2	C	401	FEJ	OAE-PAP-OAQ	2.09	118.97	110.83
2	B	401	FEJ	OAE-PAP-OAQ	2.03	118.75	110.83

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	FEJ	C-CA-N-CAL
2	A	401	FEJ	CB-CA-N-CAL
2	A	401	FEJ	OXT-C-CA-CB
2	A	401	FEJ	O-C-CA-CB
2	B	401	FEJ	C-CA-N-CAL
2	B	401	FEJ	CB-CA-N-CAL
2	B	401	FEJ	N-CA-CB-OG
2	B	401	FEJ	N-CA-CB-CG
2	C	401	FEJ	C-CA-N-CAL
2	C	401	FEJ	CB-CA-N-CAL
2	C	401	FEJ	OXT-C-CA-CB
2	C	401	FEJ	N-CA-CB-OG
2	C	401	FEJ	N-CA-CB-CG
2	D	401	FEJ	C-CA-N-CAL
2	D	401	FEJ	CB-CA-N-CAL
2	A	401	FEJ	CA-CB-CG-CD1
2	D	401	FEJ	OXT-C-CA-CB

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	401	FEJ	O-C-CA-CB
2	A	401	FEJ	OG-CB-CG-CD1
2	A	401	FEJ	CAB-OAC-PAP-OAQ
2	D	401	FEJ	N-CA-CB-CG
2	B	401	FEJ	OXT-C-CA-N
2	B	401	FEJ	C-CA-CB-CG
2	C	401	FEJ	C-CA-CB-CG


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/388 (99%)	0.32	11 (2%) 53 61	30, 44, 64, 80	0
1	B	385/388 (99%)	0.39	25 (6%) 25 30	23, 39, 65, 87	12 (3%)
1	C	377/388 (97%)	0.62	36 (9%) 14 15	27, 44, 73, 85	0
1	D	385/388 (99%)	0.24	6 (1%) 70 77	29, 42, 61, 85	0
All	All	1532/1552 (98%)	0.39	78 (5%) 33 39	23, 42, 67, 87	12 (0%)

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	142	PHE	4.8
1	C	134	VAL	4.7
1	C	153	PRO	4.5
1	B	135[A]	GLU	4.1
1	C	268	ALA	3.9
1	B	131[A]	ALA	3.7
1	C	174	ALA	3.7
1	C	152	ILE	3.6
1	D	142	PHE	3.5
1	B	138[A]	LYS	3.5
1	C	141	VAL	3.5
1	B	165	ILE	3.3
1	C	160	THR	3.3
1	C	172	TRP	3.2
1	B	139[A]	MET	3.1
1	B	164	ALA	3.1
1	C	139	MET	3.1
1	A	142	PHE	3.1
1	B	108	GLY	3.0
1	B	107	ALA	2.9
1	C	107	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	260	GLY	2.8
1	B	156	SER	2.8
1	C	161	ALA	2.8
1	C	178	TYR	2.7
1	C	140	ASN	2.7
1	C	131	ALA	2.7
1	B	281	PHE	2.6
1	C	176	PHE	2.6
1	C	129	MET	2.6
1	C	31	ASP	2.6
1	A	160	THR	2.5
1	C	169	LEU	2.4
1	B	300	ASP	2.4
1	C	269	GLY	2.4
1	B	161	ALA	2.4
1	A	267	ASN	2.4
1	C	289	ILE	2.4
1	C	130	GLY	2.4
1	D	382	LEU	2.4
1	C	301	TYR	2.4
1	B	158	SER	2.3
1	D	165	ILE	2.3
1	C	271	VAL	2.3
1	A	384	ALA	2.3
1	B	289	ILE	2.3
1	B	174	ALA	2.3
1	C	383	LYS	2.3
1	B	163	ASP	2.3
1	B	159	ARG	2.2
1	C	170	ARG	2.2
1	A	157	GLY	2.2
1	C	132	GLU	2.2
1	C	128	TYR	2.2
1	A	289	ILE	2.2
1	A	381	VAL	2.2
1	C	151	VAL	2.2
1	A	139	MET	2.2
1	A	285	GLU	2.2
1	B	154	VAL	2.1
1	B	106	GLY	2.1
1	C	11	VAL	2.1
1	C	96	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	385	SER	2.1
1	B	157	GLY	2.1
1	A	300	ASP	2.1
1	D	160	THR	2.1
1	A	291	PRO	2.1
1	C	162	LYS	2.1
1	B	137[A]	GLN	2.0
1	C	109	GLN	2.0
1	B	134[A]	VAL	2.0
1	C	133	ASP	2.0
1	C	381	VAL	2.0
1	B	153	PRO	2.0
1	C	314	ILE	2.0
1	D	281	PHE	2.0
1	D	134	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
2	FEJ	B	401	25/25	0.94	0.10	31,35,50,51	0
3	NA	A	402	1/1	0.95	0.13	45,45,45,45	0
2	FEJ	C	401	25/25	0.96	0.08	30,34,45,46	0
2	FEJ	A	401	25/25	0.96	0.08	31,34,42,42	0
3	NA	C	402	1/1	0.97	0.04	39,39,39,39	0
3	NA	D	402	1/1	0.97	0.08	35,35,35,35	0
2	FEJ	D	401	25/25	0.98	0.06	33,35,39,40	0
3	NA	B	402	1/1	0.99	0.11	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FEJ B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FEJ C 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FEJ A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.