



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 11:07 AM UTC

PDB ID : 9CU1 / pdb_00009cu1
EMDB ID : EMD-45925
Title : Azotobacter vinelandii filamentous 2:2:1 MoFeP:FeP:FeSII-Complex (termini; C1 symmetry)
Authors : Narehood, S.M.; Cook, B.D.; Srisantitham, S.; Eng, V.H.; Shiau, A.; Britt, R.D.; Herzik, M.A.; Tezcan, F.A.
Deposited on : 2024-07-25
Resolution : 2.83 Å (reported)
Based on initial model : 7UT7

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11847 (2.33 - 3.33)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	492	29% 78% 19% .
1	C	492	76% 68% 29% .
1	H	492	28% 76% 21% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	492	
2	B	523	
2	D	523	
2	I	523	
2	K	523	
3	E	290	
3	F	290	
3	L	290	
3	M	290	
4	G	122	
5	N	122	

2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 43507 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrogenase molybdenum-iron protein alpha chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	477	Total	C	N	O	S	1	0
			3790	2410	646	709	25		
1	C	477	Total	C	N	O	S	1	0
			3793	2412	646	709	26		
1	H	477	Total	C	N	O	S	1	0
			3790	2410	646	709	25		
1	J	477	Total	C	N	O	S	1	0
			3793	2412	646	709	26		

- Molecule 2 is a protein called Nitrogenase molybdenum-iron protein beta chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	522	Total	C	N	O	S	0	0
			4174	2666	705	775	28		
2	D	522	Total	C	N	O	S	0	0
			4174	2666	705	775	28		
2	I	522	Total	C	N	O	S	0	0
			4174	2666	705	775	28		
2	K	522	Total	C	N	O	S	0	0
			4174	2666	705	775	28		

- Molecule 3 is a protein called Nitrogenase iron protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	274	Total	C	N	O	S	0	0
			2077	1298	353	405	21		
3	F	283	Total	C	N	O	S	0	0
			2145	1340	363	421	21		
3	L	272	Total	C	N	O	S	0	0
			2060	1288	351	401	20		
3	M	280	Total	C	N	O	S	0	0
			2121	1323	359	418	21		

- Molecule 4 is a protein called Protein FeSII.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	121	Total	C	N	O	S	0	0
			918	574	161	177	6		


- Molecule 5 is a protein called Protein FeSII.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	N	121	Total	C	N	O	S	0	0
			920	574	163	177	6		

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
N	27	LYS	LEU	conflict	UNP Q44501
N	28	LYS	LEU	conflict	UNP Q44501

- Molecule 6 is 3-HYDROXY-3-CARBOXY-ADIPIC ACID (CCD ID: HCA) (formula: $C_7H_{10}O_7$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
6	A	1	Total	C	O	0
			14	7	7	
6	C	1	Total	C	O	0
			14	7	7	
6	H	1	Total	C	O	0
			14	7	7	
6	J	1	Total	C	O	0
			14	7	7	

- Molecule 7 is iron-sulfur-molybdenum cluster with interstitial carbon (CCD ID: ICS) (formula: CFe₇MoS₉) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
7	A	1	Total	C	Fe	Mo	S	0
			18	1	7	1	9	
7	C	1	Total	C	Fe	Mo	S	0
			18	1	7	1	9	
7	H	1	Total	C	Fe	Mo	S	0
			18	1	7	1	9	
7	J	1	Total	C	Fe	Mo	S	0
			18	1	7	1	9	

- Molecule 8 is FE(8)-S(7) CLUSTER (CCD ID: CLF) (formula: Fe₈S₇) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
8	A	1	Total	Fe	S	0
			15	8	7	
8	D	1	Total	Fe	S	0
			15	8	7	
8	H	1	Total	Fe	S	0
			15	8	7	
8	K	1	Total	Fe	S	0
			15	8	7	

- Molecule 9 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
9	B	1	Total	Fe	0
			1	1	
9	D	1	Total	Fe	0
			1	1	
9	I	1	Total	Fe	0
			1	1	
9	K	1	Total	Fe	0
			1	1	

- Molecule 10 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
10	E	1	Total 27	C 10	N 5	O 10	P 2	0
10	F	1	Total 27	C 10	N 5	O 10	P 2	0
10	L	1	Total 27	C 10	N 5	O 10	P 2	0
10	M	1	Total 27	C 10	N 5	O 10	P 2	0

- Molecule 11 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	AltConf
11	E	1	Total Mg 1 1	0
11	F	1	Total Mg 1 1	0
11	L	1	Total Mg 1 1	0
11	M	1	Total Mg 1 1	0

- Molecule 12 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
12	F	1	Total	Fe	S	0
			8	4	4	
12	L	1	Total	Fe	S	0
			8	4	4	

- Molecule 13 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
13	G	1	Total	Fe	S	0
			4	2	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
13	N	1	Total	Fe	S	0
			4	2	2	


- Molecule 14 is water.

Mol	Chain	Residues	Atoms		AltConf
14	A	73	Total	O	0
			73	73	
14	B	86	Total	O	0
			86	86	
14	C	97	Total	O	0
			97	97	
14	D	73	Total	O	0
			73	73	
14	E	78	Total	O	0
			78	78	
14	F	88	Total	O	0
			88	88	
14	G	26	Total	O	0
			26	26	
14	H	83	Total	O	0
			83	83	
14	I	108	Total	O	0
			108	108	
14	J	74	Total	O	0
			74	74	
14	K	100	Total	O	0
			100	100	
14	L	77	Total	O	0
			77	77	
14	M	94	Total	O	0
			94	94	
14	N	19	Total	O	0
			19	19	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitrogenase molybdenum-iron protein alpha chain





- Molecule 1: Nitrogenase molybdenum-iron protein alpha chain







• Molecule 1: Nitrogenase molybdenum-iron protein alpha chain





• Molecule 1: Nitrogenase molybdenum-iron protein alpha chain








• Molecule 2: Nitrogenase molybdenum-iron protein beta chain




• Molecule 2: Nitrogenase molybdenum-iron protein beta chain







• Molecule 2: Nitrogenase molybdenum-iron protein beta chain




• Molecule 3: Nitrogenase iron protein 1




- Molecule 3: Nitrogenase iron protein 1

Chain F:



- Molecule 3: Nitrogenase iron protein 1


Chain L:





- Molecule 3: Nitrogenase iron protein 1

Chain M:







• Molecule 4: Protein FeSII



• Molecule 5: Protein FeSII



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	30209	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	750	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	165000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.704	Depositor
Minimum map value	-0.536	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.027	Depositor
Recommended contour level	0.221	Depositor
Map size ($\text{\AA}$)	563.2, 563.2, 563.2	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.1, 1.1, 1.1	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ICS, MG, SF4, FE, FES, ADP, HCA, CLF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/3881	0.30	0/5234
1	C	0.16	0/3884	0.36	0/5237
1	H	0.27	0/3881	0.49	1/5234 (0.0%)
1	J	0.33	0/3884	0.54	2/5237 (0.0%)
2	B	0.13	0/4280	0.31	0/5786
2	D	0.11	0/4280	0.28	0/5786
2	I	0.13	0/4280	0.31	0/5786
2	K	0.13	0/4280	0.31	0/5786
3	E	0.14	0/2101	0.40	0/2829
3	F	0.21	0/2169	0.42	1/2921 (0.0%)
3	L	0.14	0/2084	0.36	0/2807
3	M	0.15	0/2145	0.42	0/2889
4	G	0.25	0/931	0.46	1/1256 (0.1%)
5	N	0.26	0/933	0.47	0/1256
All	All	0.19	0/43013	0.38	5/58044 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1
3	E	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	357	GLY	N-CA-C	6.82	120.92	112.73
4	G	27	LEU	N-CA-C	-5.41	105.49	111.71
3	F	119	ASP	N-CA-C	-5.39	106.28	112.92
1	J	429	PHE	CB-CA-C	5.18	119.39	110.79
1	H	360	PRO	CB-CA-C	-5.05	104.29	112.62

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	E	117	GLU	Peptide
1	H	357	GLY	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3790	0	3727	61	0
1	C	3793	0	3735	99	0
1	H	3790	0	3727	64	0
1	J	3793	0	3734	73	0
2	B	4174	0	4088	59	0
2	D	4174	0	4088	62	0
2	I	4174	0	4088	63	0
2	K	4174	0	4089	70	0
3	E	2077	0	2089	60	0
3	F	2145	0	2157	43	0
3	L	2060	0	2074	60	0
3	M	2121	0	2124	63	0
4	G	918	0	936	12	0
5	N	920	0	940	13	0
6	A	14	0	7	1	0
6	C	14	0	7	0	0
6	H	14	0	6	0	0
6	J	14	0	6	0	0
7	A	18	0	0	3	0
7	C	18	0	0	1	0
7	H	18	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	J	18	0	0	2	0
8	A	15	0	0	1	0
8	D	15	0	0	0	0
8	H	15	0	0	0	0
8	K	15	0	0	0	0
9	B	1	0	0	0	0
9	D	1	0	0	0	0
9	I	1	0	0	0	0
9	K	1	0	0	0	0
10	E	27	0	12	1	0
10	F	27	0	12	1	0
10	L	27	0	12	2	0
10	M	27	0	12	4	0
11	E	1	0	0	0	0
11	F	1	0	0	0	0
11	L	1	0	0	0	0
11	M	1	0	0	0	0
12	F	8	0	0	0	0
12	L	8	0	0	1	0
13	G	4	0	0	0	0
13	N	4	0	0	0	0
14	A	73	0	0	1	0
14	B	86	0	0	2	0
14	C	97	0	0	4	0
14	D	73	0	0	1	0
14	E	78	0	0	1	0
14	F	88	0	0	3	0
14	G	26	0	0	0	0
14	H	83	0	0	1	0
14	I	108	0	0	1	0
14	J	74	0	0	1	0
14	K	100	0	0	5	0
14	L	77	0	0	2	0
14	M	94	0	0	4	0
14	N	19	0	0	0	0
All	All	43507	0	41670	757	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 757 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:24:VAL:HG21	3:L:36:ILE:HD11	1.54	0.87
1:C:81:ILE:HG12	1:C:134:LEU:HD21	1.59	0.83
3:E:9:TYR:HB3	3:E:165:ILE:HD13	1.61	0.81
1:C:199:ASN:HD21	1:C:279:MET:HA	1.46	0.80
1:C:223:VAL:HG12	1:C:270:LEU:HB3	1.65	0.78

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/492 (32%)	151 (97%)	4 (3%)	0	100	100
1	H	476/492 (97%)	464 (98%)	12 (2%)	0	100	100
1	J	476/492 (97%)	464 (98%)	12 (2%)	0	100	100
2	I	520/523 (99%)	509 (98%)	11 (2%)	0	100	100
2	K	512/523 (98%)	500 (98%)	12 (2%)	0	100	100
3	E	260/290 (90%)	249 (96%)	11 (4%)	0	100	100
3	F	281/290 (97%)	274 (98%)	7 (2%)	0	100	100
3	L	270/290 (93%)	263 (97%)	7 (3%)	0	100	100
3	M	278/290 (96%)	269 (97%)	9 (3%)	0	100	100
4	G	119/122 (98%)	117 (98%)	2 (2%)	0	100	100
5	N	119/122 (98%)	118 (99%)	1 (1%)	0	100	100
All	All	3466/3926 (88%)	3378 (98%)	88 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	70/415 (17%)	66 (94%)	4 (6%)	18	39
1	H	407/415 (98%)	399 (98%)	8 (2%)	48	72
1	J	408/415 (98%)	393 (96%)	15 (4%)	30	55
2	D	454/455 (100%)	451 (99%)	3 (1%)	76	88
2	I	454/455 (100%)	445 (98%)	9 (2%)	48	72
2	K	454/455 (100%)	447 (98%)	7 (2%)	57	77
3	E	221/234 (94%)	216 (98%)	5 (2%)	44	68
3	F	229/234 (98%)	225 (98%)	4 (2%)	53	75
3	L	219/234 (94%)	213 (97%)	6 (3%)	39	63
3	M	226/234 (97%)	222 (98%)	4 (2%)	51	74
4	G	102/103 (99%)	100 (98%)	2 (2%)	48	72
5	N	102/103 (99%)	98 (96%)	4 (4%)	28	53
All	All	3346/3752 (89%)	3275 (98%)	71 (2%)	46	70

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	L	3	MET
3	L	52	SER
3	M	84	VAL
1	H	445	ASP
1	H	380	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	271	ASN
2	K	268	GLN
2	I	163	ASN
3	L	55	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	K	18	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 8 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	ADP	F	302	11	28,29,29	1.43	4 (14%)	43,45,45	1.84	9 (20%)
7	ICS	A	502	1	6,30,30	1.75	1 (16%)	-		
12	SF4	L	303	3	0,12,12	-	-	-		
8	CLF	A	503	1,2	0,24,24	-	-	-		
7	ICS	C	502	1	6,30,30	1.72	1 (16%)	-		
10	ADP	L	301	11	28,29,29	1.44	5 (17%)	43,45,45	1.83	8 (18%)
8	CLF	H	503	1,2	0,24,24	-	-	-		
8	CLF	D	602	1,2	0,24,24	-	-	-		
7	ICS	H	502	1	6,30,30	1.83	1 (16%)	-		
8	CLF	K	602	1,2	0,24,24	-	-	-		
10	ADP	M	301	11	28,29,29	1.41	4 (14%)	43,45,45	1.84	10 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	SF4	F	301	3	0,12,12	-	-	-		
6	HCA	C	501	-	13,13,13	1.28	1 (7%)	15,18,18	1.55	3 (20%)
13	FES	G	201	4	0,4,4	-	-	-		
7	ICS	J	502	1	6,30,30	1.83	1 (16%)	-		
10	ADP	E	301	11	28,29,29	1.41	4 (14%)	43,45,45	1.82	9 (20%)
6	HCA	H	501	-	13,13,13	1.16	1 (7%)	15,18,18	1.46	3 (20%)
6	HCA	J	501	-	13,13,13	1.15	1 (7%)	15,18,18	1.23	2 (13%)
6	HCA	A	501	-	13,13,13	1.29	1 (7%)	15,18,18	1.61	3 (20%)
13	FES	N	201	5	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	ADP	E	301	11	-	7/16/32/32	0/3/3/3
8	CLF	A	503	1,2	-	-	0/12/10/10
10	ADP	F	302	11	-	7/16/32/32	0/3/3/3
10	ADP	L	301	11	-	5/16/32/32	0/3/3/3
8	CLF	H	503	1,2	-	-	0/12/10/10
12	SF4	F	301	3	-	-	0/6/5/5
6	HCA	H	501	-	-	8/17/17/17	-
8	CLF	D	602	1,2	-	-	0/12/10/10
8	CLF	K	602	1,2	-	-	0/12/10/10
12	SF4	L	303	3	-	-	0/6/5/5
6	HCA	C	501	-	-	0/17/17/17	-
6	HCA	J	501	-	-	1/17/17/17	-
13	FES	G	201	4	-	-	0/1/1/1
6	HCA	A	501	-	-	1/17/17/17	-
10	ADP	M	301	11	-	5/16/32/32	0/3/3/3
13	FES	N	201	5	-	-	0/1/1/1

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	L	301	ADP	C5-C4	4.75	1.47	1.39
10	M	301	ADP	C5-C4	4.72	1.47	1.39
10	F	302	ADP	C5-C4	4.69	1.47	1.39
10	E	301	ADP	C5-C4	4.65	1.47	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	J	502	ICS	S2B-FE6	-3.68	2.16	2.24

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	L	301	ADP	C5-C4-N3	-5.96	118.51	126.72
10	M	301	ADP	C5-C4-N3	-5.87	118.63	126.72
10	F	302	ADP	C5-C4-N3	-5.85	118.67	126.72
10	E	301	ADP	C5-C4-N3	-5.75	118.80	126.72
10	L	301	ADP	N3-C4-N9	4.71	135.18	127.17

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	501	HCA	C2-C3-C4-C5
6	H	501	HCA	C2-C3-C4-C5
6	H	501	HCA	C7-C3-C4-C5
6	H	501	HCA	O7-C3-C4-C5
10	E	301	ADP	C5'-O5'-PA-O1A


There are no ring outliers.

11 monomers are involved in 19 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	F	302	ADP	1	0
7	A	502	ICS	3	0
12	L	303	SF4	1	0
8	A	503	CLF	1	0
7	C	502	ICS	1	0
10	L	301	ADP	2	0
7	H	502	ICS	2	0
10	M	301	ADP	4	0
7	J	502	ICS	2	0
10	E	301	ADP	1	0
6	A	501	HCA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is


within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand ICS A 502




Bond lengths




Bond angles




Torsions




Rings







Ligand ICS C 502




Bond lengths



Bond angles




Torsions




Rings


Ligand ADP L 301




Bond lengths




Bond angles




Torsions




Rings







Ligand ICS H 502




Bond lengths





Bond angles





Torsions






Rings























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-45925. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y



Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256




Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256




Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 242




Y Index: 261




Z Index: 313


6.3.2 Raw map



X Index: 242



Y Index: 260




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.221. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_45925_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 152 nm^3 ; this corresponds to an approximate mass of 137 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.353 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.353 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.83	-	-
Author-provided FSC curve	2.83	3.41	2.91
Unmasked-calculated*	7.25	14.37	7.76

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.25 differs from the reported value 2.83 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-45925 and PDB model 9CU1. Per-residue inclusion information can be found in section 3 on page 11.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion ⓘ



At the recommended contour level, 43% of all backbone atoms, 43% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.221) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4260	0.3330
A	0.5110	0.3420
B	0.4940	0.3450
C	0.2410	0.3160
D	0.4080	0.3320
E	0.1470	0.2150
F	0.2160	0.3020
G	0.5330	0.3600
H	0.5440	0.3440
I	0.5760	0.3670
J	0.5050	0.3680
K	0.5630	0.3800
L	0.2220	0.2530
M	0.2690	0.2980
N	0.5330	0.3500

1.0

0.0

<0.0