



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:09 PM UTC

PDB ID : 4CVM / pdb_00004cvm
Title : PaMurF in complex with AMP-PNP and UDP-MurNAc-tripeptide (mDAP)
Authors : Ruane, K.M.; Majce, V.
Deposited on : 2014-03-28
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	5% 87% 6% 7%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-N-ACETYLMURAMOYL-TRIPETIDE--D-ALANYL-D- ALANINE LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	1	0
			3159	1980	563	606	10			

There are 19 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	459	SER	-	expression tag	UNP Q9HVVZ7
A	460	SER	-	expression tag	UNP Q9HVVZ7
A	461	VAL	-	expression tag	UNP Q9HVVZ7
A	462	ASP	-	expression tag	UNP Q9HVVZ7
A	463	LYS	-	expression tag	UNP Q9HVVZ7
A	464	LEU	-	expression tag	UNP Q9HVVZ7
A	465	ALA	-	expression tag	UNP Q9HVVZ7
A	466	ALA	-	expression tag	UNP Q9HVVZ7
A	467	ALA	-	expression tag	UNP Q9HVVZ7
A	468	LEU	-	expression tag	UNP Q9HVVZ7
A	469	GLU	-	expression tag	UNP Q9HVVZ7
A	470	HIS	-	expression tag	UNP Q9HVVZ7
A	471	HIS	-	expression tag	UNP Q9HVVZ7
A	472	HIS	-	expression tag	UNP Q9HVVZ7
A	473	HIS	-	expression tag	UNP Q9HVVZ7
A	474	HIS	-	expression tag	UNP Q9HVVZ7
A	475	HIS	-	expression tag	UNP Q9HVVZ7
A	39	GLU	GLN	conflict	UNP Q9HVVZ7
A	428	ASP	GLN	conflict	UNP Q9HVVZ7

- Molecule 2 is ALANINE (CCD ID: ALA) (formula: $C_3H_7NO_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 3 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: C₉H₁₄N₂O₁₂P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			24	9	2	11	2		

- Molecule 4 is N-acetyl-alpha-muramic acid (CCD ID: MUB) (formula: C₁₁H₁₉NO₈).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			19	11	1	7		

- Molecule 5 is GAMMA-D-GLUTAMIC ACID (CCD ID: FGA) (formula: $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			6	3	1	2		

- Molecule 6 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $\text{C}_{10}\text{H}_{17}\text{N}_6\text{O}_{12}\text{P}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Mg	0	0
			2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	59	Total	O	0	0
			59	59		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.54Å 86.05Å 88.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.61 – 2.06 61.61 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.9 (61.61-2.06) 99.9 (61.61-2.06)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.182 , 0.224 0.194 , 0.233	Depositor DCC
R_{free} test set	1428 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.022 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3305	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FGA, UDP, ANP, MUB, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	3/3200 (0.1%)	1.02	5/4354 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	THR	C-O	8.73	1.34	1.24
1	A	147	VAL	CA-CB	-6.01	1.51	1.54
1	A	172	GLU	C-O	-5.08	1.18	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	409	ALA	N-CA-C	5.77	117.65	111.36
1	A	358	SER	N-CA-C	5.35	119.91	112.90
1	A	428	ASP	CA-C-N	5.33	125.27	119.78
1	A	428	ASP	C-N-CA	5.33	125.27	119.78
1	A	217	ALA	N-CA-C	5.26	117.76	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3159	0	3171	11	0
2	A	5	0	4	0	0
3	A	24	0	11	0	0
4	A	19	0	17	0	0
5	A	6	0	2	0	0
6	A	31	0	13	0	0
7	A	2	0	0	0	0
8	A	59	0	0	0	0
All	All	3305	0	3218	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ALA:HB2	1:A:173:ILE:CD1	2.23	0.69
1:A:104:LEU:H	1:A:183:HIS:HD2	1.45	0.64
1:A:167:ALA:HB2	1:A:173:ILE:HD11	1.84	0.58
1:A:186:ILE:HD11	1:A:299:ALA:HB2	1.90	0.53
1:A:117:LYS:HE3	1:A:118:GLU:OE2	2.14	0.47
1:A:348:LEU:HD22	1:A:385:TYR:CG	2.49	0.47
1:A:429:PRO:O	1:A:430:THR:OG1	2.27	0.46
1:A:107:MET:HE1	1:A:162:VAL:HG13	1.97	0.46
1:A:434:LEU:HD23	1:A:434:LEU:C	2.41	0.45
1:A:264:ARG:HH12	1:A:283:ASN:HD22	1.66	0.43
1:A:12:VAL:HG13	1:A:13:ALA:N	2.34	0.42

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	438/475 (92%)	431 (98%)	5 (1%)	2 (0%)	24 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	216	ALA
1	A	425	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	306/345 (89%)	305 (100%)	1 (0%)	86 89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	426	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	183	HIS
1	A	283	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	UDP	A	1452	4	21,25,26	1.19	2 (9%)	32,37,40	1.80	9 (28%)
6	ANP	A	1456	7	33,33,33	2.06	11 (33%)	45,52,52	2.34	15 (33%)
2	ALA	A	1454	4,5	3,4,5	0.62	0	2,4,6	0.87	0
5	FGA	A	1455	2	5,5,9	1.08	0	6,6,11	1.10	0
4	MUB	A	1453	3,2	18,19,20	1.14	1 (5%)	21,26,28	1.79	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UDP	A	1452	4	-	0/13/31/32	0/2/2/2
6	ANP	A	1456	7	-	4/18/38/38	0/3/3/3
2	ALA	A	1454	4,5	-	0/1/2/4	-
5	FGA	A	1455	2	-	4/4/4/9	-
4	MUB	A	1453	3,2	-	1/11/32/34	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1456	ANP	C5-C4	4.59	1.47	1.39
6	A	1456	ANP	PG-N3B	3.96	1.73	1.63
6	A	1456	ANP	PB-O3A	3.81	1.63	1.59
6	A	1456	ANP	PB-N3B	3.72	1.73	1.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1456	ANP	PG-O1G	3.53	1.51	1.46
6	A	1456	ANP	PG-O3G	-3.38	1.47	1.56
6	A	1456	ANP	C8-N7	3.09	1.37	1.31
6	A	1456	ANP	PA-O3A	2.97	1.62	1.59
6	A	1456	ANP	PB-O1B	2.95	1.50	1.46
4	A	1453	MUB	C1-C2	2.42	1.55	1.52
3	A	1452	UDP	C5-C4	-2.34	1.38	1.43
6	A	1456	ANP	C5-C6	2.29	1.47	1.41
6	A	1456	ANP	C5-N7	-2.10	1.35	1.39
3	A	1452	UDP	O4'-C4'	-2.10	1.40	1.45

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1456	ANP	O1G-PG-N3B	-5.26	104.03	111.77
6	A	1456	ANP	N3-C2-N1	-5.06	120.92	128.58
6	A	1456	ANP	C5-C4-N3	-5.06	119.75	126.72
3	A	1452	UDP	C4-N3-C2	-4.51	121.01	126.61
6	A	1456	ANP	N3-C4-N9	4.41	134.67	127.17
6	A	1456	ANP	C2-N3-C4	4.29	122.31	111.83
4	A	1453	MUB	C1-O5-C5	-4.21	105.51	113.65
6	A	1456	ANP	O2B-PB-O1B	4.13	118.74	109.87
4	A	1453	MUB	C1-C2-N2	-4.07	106.01	110.73
3	A	1452	UDP	N3-C2-N1	3.90	119.97	114.89
6	A	1456	ANP	O2G-PG-O3G	3.87	118.00	107.59
6	A	1456	ANP	C4-N9-C8	3.66	109.58	105.74
3	A	1452	UDP	C5-C4-N3	3.52	119.73	114.80
4	A	1453	MUB	O5-C1-C2	3.35	112.88	109.52
6	A	1456	ANP	O2G-PG-O1G	-3.26	105.29	113.45
6	A	1456	ANP	N9-C8-N7	-3.16	109.45	113.94
3	A	1452	UDP	O2A-PA-O3A	3.14	115.76	107.27
6	A	1456	ANP	O3G-PG-O1G	2.97	120.89	113.45
3	A	1452	UDP	O4'-C1'-C2'	-2.69	100.86	106.62
6	A	1456	ANP	C5-N7-C8	2.53	107.43	103.45
6	A	1456	ANP	C2-N1-C6	2.50	122.84	118.73
3	A	1452	UDP	O4-C4-C5	-2.46	120.92	125.16
3	A	1452	UDP	C1'-N1-C2	2.37	121.86	117.59
6	A	1456	ANP	C6-C5-N7	2.34	136.60	132.09
3	A	1452	UDP	C4'-O4'-C1'	-2.32	104.34	109.47
6	A	1456	ANP	C4-C5-N7	-2.29	107.96	110.58
4	A	1453	MUB	O6-C6-C5	-2.27	103.59	111.33
3	A	1452	UDP	O2-C2-N3	-2.15	117.53	121.49

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1453	MUB	O10-C10-C9-C11
6	A	1456	ANP	PB-N3B-PG-O1G
6	A	1456	ANP	PG-N3B-PB-O1B
6	A	1456	ANP	PA-O3A-PB-O2B
5	A	1455	FGA	OXT-C-CA-N
5	A	1455	FGA	O-C-CA-CB
5	A	1455	FGA	OXT-C-CA-CB
5	A	1455	FGA	O-C-CA-N
6	A	1456	ANP	PA-O3A-PB-O1B

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand UDP A 1452



Ligand ANP A 1456





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/475 (93%)	0.28	25 (5%) 30 30	16, 31, 47, 57	1 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	369	GLY	4.1
1	A	12	VAL	3.6
1	A	376	GLU	3.1
1	A	239	GLY	3.0
1	A	168	SER	2.7
1	A	378	ALA	2.7
1	A	402	HIS	2.6
1	A	425	ALA	2.6
1	A	191	GLY	2.5
1	A	426	THR	2.5
1	A	409	ALA	2.4
1	A	216	ALA	2.4
1	A	398	PRO	2.3
1	A	15	ASP	2.3
1	A	406	ALA	2.3
1	A	248	ASP	2.2
1	A	199	GLY	2.1
1	A	344	PRO	2.1
1	A	358	SER	2.1
1	A	422	GLY	2.1
1	A	238	SER	2.1
1	A	377	GLN	2.1
1	A	382	VAL	2.1
1	A	449	ALA	2.1
1	A	412	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	FGA	A	1455	6/10	0.73	0.16	34,41,48,61	0
2	ALA	A	1454	5/6	0.93	0.10	29,29,31,32	0
4	MUB	A	1453	19/20	0.94	0.08	30,33,39,40	0
7	MG	A	1457	1/1	0.94	0.09	39,39,39,39	0
6	ANP	A	1456	31/31	0.97	0.07	25,28,40,41	0
3	UDP	A	1452	24/25	0.98	0.05	22,28,33,34	0
7	MG	A	1458	1/1	0.98	0.07	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MUB A 1453:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ANP A 1456:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UDP A 1452:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.