



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:51 AM UTC

PDB ID : 3CW6 / pdb_00003cw6
Title : E. coli Initiator tRNA
Authors : Barraud, P.; Schmitt, E.; Mechulam, Y.; Dardel, F.; Tisne, C.
Deposited on : 2008-04-21
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)
RNA backbone	3983	1048 (3.60-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	77	21% 36% 44% 14% 5%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called Initiator tRNA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	77	1645	734	297	536	77	1	0	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Initiator tRNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.80Å 93.80Å 218.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.91 – 3.30 38.91 – 3.30	Depositor EDS
% Data completeness (in resolution range)	86.1 (38.91-3.30) 86.0 (38.91-3.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 3.32Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.231 , 0.274 0.233 , 0.253	Depositor DCC
R_{free} test set	779 reflections (8.49%)	wwPDB-VP
Wilson B-factor (Å ²)	114.9	Xtriage
Anisotropy	0.353	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.52 , 999.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1645	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, H2U, PSU, 5MU, 4SU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/1725	1.29	14/2687 (0.5%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	16	C	P-O3'-C3'	-7.06	109.61	120.20
1	A	37	A	O4'-C4'-C3'	-6.69	99.41	106.10
1	A	58	A	O3'-P-O5'	-6.67	93.99	104.00
1	A	9	G	O5'-C5'-C4'	6.36	121.23	111.70
1	A	9	G	P-O5'-C5'	6.00	129.90	120.90
1	A	24	U	P-O3'-C3'	-6.00	111.21	120.20
1	A	56	C	O4'-C4'-C3'	-5.80	98.20	104.00
1	A	47	U	C4'-C3'-C2'	-5.62	96.98	102.60
1	A	9	G	O4'-C4'-C3'	-5.51	100.59	106.10
1	A	64	G	C4'-C3'-C2'	-5.46	97.14	102.60
1	A	47	U	O4'-C4'-C3'	-5.36	98.64	104.00
1	A	3	C	C5'-C4'-C3'	-5.25	108.12	116.00
1	A	5	G	O4'-C4'-C3'	-5.22	98.78	104.00
1	A	47	U	C3'-C2'-O2'	5.20	118.50	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1645	0	842	27	0
All	All	1645	0	842	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:H2U:O2'	1:A:21:A:OP1	2.03	0.75
1:A:16:C:O2'	1:A:61:C:OP1	2.12	0.67
1:A:20:H2U:H4'	1:A:20:H2U:OP1	1.93	0.67
1:A:10:G:H2'	1:A:11:A:H8	1.66	0.61
1:A:17:C:H3'	1:A:17(A):U:H2'	1.84	0.58
1:A:27:U:H2'	1:A:28:C:H6	1.69	0.58
1:A:19:G:C5	1:A:57:A:C2	2.92	0.57
1:A:22:G:H8	1:A:22:G:H5''	1.70	0.56
1:A:27:U:H2'	1:A:28:C:C6	2.41	0.55
1:A:13:C:C2'	1:A:14:A:O5'	2.55	0.55
1:A:20:H2U:OP1	1:A:20:H2U:C4'	2.57	0.53
1:A:14:A:C5	1:A:22:G:C2	2.97	0.52
1:A:20:H2U:O2'	1:A:21:A:P	2.68	0.51
1:A:51:C:H2'	1:A:52:G:O4'	2.10	0.51
1:A:19:G:C4	1:A:57:A:C2	2.99	0.50
1:A:15:G:N2	1:A:59:A:C5	2.80	0.49
1:A:9:G:N3	1:A:45:G:H2'	2.27	0.49
1:A:14:A:C4	1:A:22:G:N2	2.82	0.48
1:A:56:C:O2	1:A:56:C:H2'	2.14	0.47
1:A:13:C:H2'	1:A:14:A:O5'	2.15	0.47
1:A:72:A:H2'	1:A:73:A:C8	2.51	0.46
1:A:22:G:H5''	1:A:22:G:C8	2.52	0.44
1:A:10:G:H2'	1:A:11:A:C8	2.50	0.43
1:A:45:G:O5'	1:A:45:G:H8	2.03	0.41
1:A:48:C:C2	1:A:59:A:O4'	2.74	0.41
1:A:23:C:H2'	1:A:24:U:H6	1.86	0.41
1:A:15:G:N2	1:A:59:A:C4	2.88	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	76/77 (98%)	24 (31%)	6 (7%)

All (24) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	2	G
1	A	4	G
1	A	7	G
1	A	8	4SU
1	A	9	G
1	A	10	G
1	A	13	C
1	A	14	A
1	A	16	C
1	A	18	G
1	A	20	H2U
1	A	21	A
1	A	31	G
1	A	34	C
1	A	37	A
1	A	38	A
1	A	42	G
1	A	47	U
1	A	48	C
1	A	49	G
1	A	56	C
1	A	59	A
1	A	60	U
1	A	76	A

All (6) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	9	G
1	A	13	C
1	A	19	G
1	A	37	A
1	A	46	A
1	A	47	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	5MU	A	54	1	19,22,23	1.62	6 (31%)	27,32,35	2.21	10 (37%)
1	4SU	A	8	1	18,21,22	2.19	5 (27%)	25,30,33	3.45	10 (40%)
1	H2U	A	20	1	18,21,22	1.24	2 (11%)	19,30,33	1.23	2 (10%)
1	OMC	A	32	1	19,22,23	0.44	0	25,31,34	0.77	1 (4%)
1	PSU	A	55	1	18,21,22	1.49	2 (11%)	21,30,33	2.07	7 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5MU	A	54	1	-	0/7/25/26	0/2/2/2
1	4SU	A	8	1	-	1/7/25/26	0/2/2/2
1	H2U	A	20	1	-	6/7/38/39	0/2/2/2
1	OMC	A	32	1	-	1/9/27/28	0/2/2/2
1	PSU	A	55	1	-	1/7/25/26	0/2/2/2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	8	4SU	C4-S4	-6.20	1.57	1.68
1	A	8	4SU	C4-N3	-3.73	1.33	1.37
1	A	54	5MU	C4-C5	-3.65	1.38	1.44
1	A	55	PSU	C6-C5	3.52	1.39	1.35
1	A	8	4SU	C2-N1	3.45	1.43	1.38
1	A	20	H2U	C4-N3	-3.04	1.32	1.37
1	A	20	H2U	C2-N3	-3.00	1.32	1.38
1	A	8	4SU	C5-C4	-2.88	1.39	1.42
1	A	54	5MU	C6-C5	2.64	1.38	1.34
1	A	54	5MU	C2-N1	2.53	1.42	1.38
1	A	55	PSU	C4-N3	-2.43	1.34	1.38
1	A	8	4SU	C2-N3	-2.38	1.33	1.38
1	A	54	5MU	C4-N3	-2.35	1.34	1.38
1	A	54	5MU	C6-N1	-2.14	1.34	1.38
1	A	54	5MU	C2-N3	-2.07	1.34	1.38

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	8	4SU	C5-C4-N3	9.35	123.45	114.75
1	A	8	4SU	C4-N3-C2	-9.00	118.68	127.31
1	A	8	4SU	C5-C4-S4	-5.91	117.55	124.31
1	A	55	PSU	N1-C2-N3	5.47	120.94	115.17
1	A	54	5MU	C5-C4-N3	5.26	119.89	115.32
1	A	54	5MU	O4-C4-C5	-4.38	119.91	124.92
1	A	54	5MU	C4-N3-C2	-4.34	121.65	127.34
1	A	8	4SU	N3-C2-N1	4.06	120.18	114.89
1	A	8	4SU	C2'-C3'-C4'	-3.88	95.11	102.61
1	A	8	4SU	C2'-C1'-N1	-3.85	102.53	113.25
1	A	55	PSU	C4-N3-C2	-3.66	121.33	126.37
1	A	54	5MU	N3-C2-N1	3.28	119.16	114.89
1	A	54	5MU	C5-C6-N1	-3.27	119.77	123.31
1	A	8	4SU	O3'-C3'-C4'	3.19	120.26	111.08
1	A	55	PSU	O4'-C4'-C3'	-3.09	99.01	105.15
1	A	54	5MU	C5M-C5-C6	-2.97	118.83	122.85
1	A	55	PSU	O2-C2-N1	-2.94	119.75	122.79
1	A	8	4SU	O3'-C3'-C2'	-2.67	103.26	111.82
1	A	54	5MU	C5M-C5-C4	2.51	121.47	118.78
1	A	8	4SU	C1'-N1-C2	2.44	121.98	117.59
1	A	55	PSU	C5-C6-N1	-2.27	118.99	122.14
1	A	8	4SU	O2-C2-N3	-2.26	117.31	121.49
1	A	32	OMC	CM2-O2'-C2'	2.18	120.08	114.47
1	A	20	H2U	C5-C6-N1	-2.13	105.07	111.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	5MU	C2'-C1'-N1	-2.13	107.33	113.25
1	A	55	PSU	O2'-C2'-C1'	-2.12	106.17	111.21
1	A	54	5MU	C1'-N1-C2	2.10	121.37	117.59
1	A	55	PSU	C6-C5-C4	2.08	119.58	118.17
1	A	54	5MU	C6-C5-C4	2.03	119.69	118.02
1	A	20	H2U	C3'-C2'-C1'	2.02	105.28	101.46

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	20	H2U	C4'-C5'-O5'-P
1	A	20	H2U	O4'-C1'-N1-C2
1	A	20	H2U	O4'-C1'-N1-C6
1	A	20	H2U	C3'-C4'-C5'-O5'
1	A	20	H2U	O4'-C4'-C5'-O5'
1	A	55	PSU	O4'-C4'-C5'-O5'
1	A	32	OMC	C3'-C2'-O2'-CM2
1	A	20	H2U	C2'-C1'-N1-C2
1	A	8	4SU	C2'-C1'-N1-C2

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	20	H2U	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	72/77 (93%)	1.38	16 (22%) 2 2	66, 82, 100, 120	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	C	7.1
1	A	2	G	6.5
1	A	47	U	6.4
1	A	75	C	4.2
1	A	59	A	4.2
1	A	74	C	4.2
1	A	26	G	3.5
1	A	18	G	3.2
1	A	76	A	2.7
1	A	16	C	2.7
1	A	3	C	2.3
1	A	4	G	2.3
1	A	5	G	2.2
1	A	73	A	2.2
1	A	48	C	2.1
1	A	56	C	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	OMC	A	32	21/22	0.85	0.17	83,84,84,84	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	4SU	A	8	20/21	0.86	0.25	77,83,88,90	0
1	H2U	A	20	20/21	0.88	0.15	72,77,85,86	0
1	PSU	A	55	20/21	0.89	0.13	66,70,72,72	0
1	5MU	A	54	21/22	0.92	0.13	68,70,74,75	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.