



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 03:17 AM UTC

PDB ID : 4CZX / pdb_00004czz
Title : Complex of *Neurospora crassa* PAN2 (WD40) with PAN3 (C-TERM)
Authors : Jonas, S.; Izaurralde, E.; Weichenrieder, O.
Deposited on : 2014-04-22
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	3% 90% 10%
2	B	147	2% 56% 13% • 30%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MES	A	400	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	8	0
			2576	1644	449	466	17			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PRO	-	expression tag	UNP P0C581
A	-1	HIS	-	expression tag	UNP P0C581
A	0	MET	-	expression tag	UNP P0C581

- Molecule 2 is a protein called PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	103	Total	C	N	O	S	0	3	0
			847	532	152	159	4			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	510	MET	-	expression tag	UNP Q7SDP4
B	511	GLY	-	expression tag	UNP Q7SDP4
B	512	SER	-	expression tag	UNP Q7SDP4
B	513	SER	-	expression tag	UNP Q7SDP4
B	514	HIS	-	expression tag	UNP Q7SDP4
B	515	HIS	-	expression tag	UNP Q7SDP4
B	516	HIS	-	expression tag	UNP Q7SDP4
B	517	HIS	-	expression tag	UNP Q7SDP4
B	518	HIS	-	expression tag	UNP Q7SDP4
B	519	HIS	-	expression tag	UNP Q7SDP4
B	520	SER	-	expression tag	UNP Q7SDP4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	521	SER	-	expression tag	UNP Q7SDP4
B	522	GLY	-	expression tag	UNP Q7SDP4
B	523	THR	-	expression tag	UNP Q7SDP4
B	524	GLY	-	expression tag	UNP Q7SDP4
B	525	SER	-	expression tag	UNP Q7SDP4
B	526	GLY	-	expression tag	UNP Q7SDP4
B	527	GLU	-	expression tag	UNP Q7SDP4
B	528	ASN	-	expression tag	UNP Q7SDP4
B	529	LEU	-	expression tag	UNP Q7SDP4
B	530	TYR	-	expression tag	UNP Q7SDP4
B	531	PHE	-	expression tag	UNP Q7SDP4
B	532	GLN	-	expression tag	UNP Q7SDP4
B	533	GLY	-	expression tag	UNP Q7SDP4
B	534	HIS	-	expression tag	UNP Q7SDP4
B	535	MET	-	expression tag	UNP Q7SDP4
B	536	LEU	-	expression tag	UNP Q7SDP4
B	537	GLU	-	expression tag	UNP Q7SDP4

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	238	Total 238	O 238	0	0
6	B	70	Total 70	O 70	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN2



• Molecule 2: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	55.92Å 160.65Å 124.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 1.85 49.21 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.21-1.85) 99.6 (49.21-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 1.86Å)	Xtriage
Refinement program	PHENIX PHENIX.REFINE 1.8.4_1496)	Depositor
R, R_{free}	0.168 , 0.197 0.173 , 0.199	Depositor DCC
R_{free} test set	2432 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.860	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3754	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MES, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2669	0.77	2/3627 (0.1%)
2	B	0.41	0/866	0.65	0/1162
All	All	0.42	0/3535	0.74	2/4789 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	272	ASN	CA-C-N	5.52	125.61	119.87
1	A	272	ASN	C-N-CA	5.52	125.61	119.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2576	0	2548	21	0
2	B	847	0	852	14	0
3	A	12	0	13	6	0
4	A	5	0	0	0	0
5	A	6	0	8	0	0
6	A	238	0	0	1	0
6	B	70	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3754	0	3421	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:ARG:HH12	3:A:400:MES:H82	1.38	0.87
2:B:544:ILE:HD12	2:B:603[A]:LEU:HB3	1.80	0.63
2:B:556:GLU:OE2	2:B:579:ARG:NH1	2.33	0.61
2:B:539:LEU:O	2:B:543:ARG:HG3	2.02	0.59
1:A:210:ARG:NH1	3:A:400:MES:H71	2.20	0.56
2:B:564:HIS:N	2:B:568:GLU:OE1	2.39	0.56
2:B:574:LEU:HB3	2:B:635[B]:MET:HE1	1.90	0.54
1:A:210:ARG:HH12	3:A:400:MES:C8	2.14	0.53
1:A:264:GLN:NE2	6:A:2219:HOH:O	2.35	0.52
1:A:1:MET:HG3	1:A:279:CYS:HB3	1.92	0.52
1:A:261:GLN:HB3	1:A:286:VAL:HG13	1.91	0.51
1:A:320:THR:HG22	1:A:321:GLU:H	1.74	0.51
1:A:294:ARG:HB2	1:A:320:THR:HG23	1.93	0.51
2:B:556:GLU:CD	2:B:579:ARG:HH12	2.17	0.51
1:A:285:GLY:HA3	1:A:305:LYS:HG2	1.95	0.49
2:B:588:ALA:O	2:B:589:ASP:HB2	2.13	0.48
1:A:191:PHE:CD2	3:A:400:MES:H32	2.50	0.47
2:B:559:ASP:HA	2:B:572:ARG:HH12	1.80	0.47
1:A:1:MET:HE3	1:A:6:TRP:HB2	1.97	0.46
1:A:193:ASN:OD1	3:A:400:MES:H61	2.16	0.46
1:A:129:TRP:HA	1:A:152:GLN:HB3	1.98	0.45
2:B:540:GLU:O	2:B:544:ILE:HG12	2.17	0.45
1:A:50:SER:HG	1:A:60:THR:H	1.66	0.43
1:A:228:LEU:HD23	1:A:228:LEU:HA	1.89	0.43
2:B:548:MET:HE1	2:B:583:PHE:CZ	2.53	0.43
1:A:-2:PRO:O	1:A:-1:HIS:ND1	2.51	0.43
1:A:191:PHE:CG	3:A:400:MES:H32	2.53	0.43
1:A:259:VAL:HG22	1:A:265:ILE:HG12	2.01	0.42
2:B:539:LEU:HD23	2:B:539:LEU:HA	1.92	0.42
2:B:544:ILE:O	2:B:548:MET:HG2	2.20	0.42
1:A:102:SER:HB3	2:B:565:ASN:HD22	1.85	0.42
2:B:596:THR:HA	2:B:599:TYR:CD2	2.54	0.41
1:A:17:PRO:HA	1:A:22:TYR:CG	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:THR:HG23	1:A:320:THR:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/324 (102%)	320 (97%)	9 (3%)	1 (0%)	36	25
2	B	102/147 (69%)	97 (95%)	5 (5%)	0	100	100
All	All	432/471 (92%)	417 (96%)	14 (3%)	1 (0%)	43	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/279 (102%)	280 (99%)	4 (1%)	59	49
2	B	93/126 (74%)	92 (99%)	1 (1%)	65	57
All	All	377/405 (93%)	372 (99%)	5 (1%)	59	51

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ILE
1	A	47	PHE
1	A	132	LYS
1	A	320	THR
2	B	556	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	ASN
1	A	264	GLN
1	A	272	ASN
2	B	565	ASN
2	B	598	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	402	-	5,5,5	0.35	0	5,5,5	0.52	0
4	PO4	A	401	-	4,4,4	0.93	0	6,6,6	0.53	0
3	MES	A	400	-	12,12,12	1.65	2 (16%)	15,16,16	1.66	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	402	-	-	2/4/4/4	-
3	MES	A	400	-	-	5/6/14/14	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	400	MES	O2S-S	3.84	1.56	1.45
3	A	400	MES	O1S-S	2.52	1.52	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	400	MES	O1-C6-C5	-3.40	104.45	111.77
3	A	400	MES	O3S-S-C8	3.16	112.19	106.00
3	A	400	MES	O2S-S-C8	2.50	110.51	106.73
3	A	400	MES	C7-N4-C5	-2.36	104.94	111.24

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	400	MES	C7-C8-S-O2S
5	A	402	GOL	O1-C1-C2-C3
5	A	402	GOL	O1-C1-C2-O2
3	A	400	MES	N4-C7-C8-S
3	A	400	MES	C8-C7-N4-C3
3	A	400	MES	C8-C7-N4-C5
3	A	400	MES	C7-C8-S-O3S

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	400	MES	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/324 (100%)	-0.08	9 (2%) 55 58	14, 34, 64, 102	8 (2%)
2	B	103/147 (70%)	-0.00	3 (2%) 53 57	15, 34, 72, 105	3 (2%)
All	All	427/471 (90%)	-0.06	12 (2%) 55 58	14, 34, 67, 105	11 (2%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	320	THR	3.9
2	B	539	LEU	3.8
1	A	321	GLU	3.3
1	A	-1	HIS	2.9
2	B	564	HIS	2.7
1	A	319	PHE	2.7
1	A	315	ASN	2.5
1	A	272	ASN	2.3
1	A	317	ILE	2.3
1	A	229	LYS	2.2
1	A	318	GLN	2.0
2	B	558	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PO4	A	401	5/5	0.80	0.13	48,48,51,53	5
5	GOL	A	402	6/6	0.88	0.11	49,53,58,64	0
3	MES	A	400	12/12	0.97	0.08	15,27,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.