



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 10:50 AM UTC

PDB ID : 3D0A / pdb_00003d0a
Title : Human p53 core domain with hot spot mutation R249S and second site suppressor mutation H168R in sequence-specific complex with DNA
Authors : Suad, O.; Rozenberg, H.; Shakked, Z.
Deposited on : 2008-05-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	12	8% 83% 8% 8%
1	F	12	8% 75% 17% 8%
1	G	12	8% 58% 42%
1	H	12	8% 75% 17% 8%
2	A	200	% 87% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	200	6% 90% 8%
2	C	200	4% 88% 8%
2	D	200	2% 90% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*DCP*DGP*DGP*DGP*DCP*DAP*DTP*DGP*DCP*DCP*DCP*DG)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	12	Total	C	N	O	P	0	0	0
			213	98	39	65	11			
1	F	11	Total	C	N	O	P	0	0	0
			216	102	44	59	11			
1	G	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
1	H	12	Total	C	N	O	P	0	0	0
			227	107	47	63	10			

- Molecule 2 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	196	Total	C	N	O	S	0	11	0
			1577	974	288	295	20			
2	B	197	Total	C	N	O	S	0	11	0
			1550	957	282	292	19			
2	C	193	Total	C	N	O	S	0	12	0
			1554	963	277	294	20			
2	D	194	Total	C	N	O	S	0	7	0
			1522	946	272	284	20			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	ARG	HIS	engineered mutation	UNP P04637
A	249	SER	ARG	engineered mutation	UNP P04637
B	168	ARG	HIS	engineered mutation	UNP P04637
B	249	SER	ARG	engineered mutation	UNP P04637
C	168	ARG	HIS	engineered mutation	UNP P04637
C	249	SER	ARG	engineered mutation	UNP P04637
D	168	ARG	HIS	engineered mutation	UNP P04637
D	249	SER	ARG	engineered mutation	UNP P04637

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0
3	B	1	Total 1	Zn 1	0	0
3	C	1	Total 1	Zn 1	0	0
3	D	1	Total 1	Zn 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	24	Total 24	O 24	0	0
4	F	14	Total 14	O 14	0	0
4	G	12	Total 12	O 12	0	0
4	H	9	Total 9	O 9	0	0
4	A	213	Total 216	O 216	0	3
4	B	113	Total 113	O 113	0	0
4	C	219	Total 220	O 220	0	1
4	D	143	Total 143	O 143	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*DCP*DGP*DGP*DGP*DCP*DAP*DTP*DGP*DCP*DCP*DCP*DG)-3')




- Molecule 1: DNA (5'-D(*DCP*DGP*DGP*DGP*DCP*DAP*DTP*DGP*DCP*DCP*DCP*DG)-3')




- Molecule 1: DNA (5'-D(*DCP*DGP*DGP*DGP*DCP*DAP*DTP*DGP*DCP*DCP*DCP*DG)-3')



- Molecule 1: DNA (5'-D(*DCP*DGP*DGP*DGP*DCP*DAP*DTP*DGP*DCP*DCP*DCP*DG)-3')





- Molecule 2: Cellular tumor antigen p53








- Molecule 2: Cellular tumor antigen p53



- Molecule 2: Cellular tumor antigen p53



- Molecule 2: Cellular tumor antigen p53



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.30Å 57.99Å 78.93Å 83.05° 88.11° 74.08°	Depositor
Resolution (Å)	17.76 – 1.80 17.76 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.6 (17.76-1.80) 97.3 (17.76-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.183 , 0.241 0.184 , 0.240	Depositor DCC
R_{free} test set	4181 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7857	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.50	0/236	1.31	1/362 (0.3%)
1	F	0.48	0/238	1.19	0/360
1	G	0.45	0/272	1.17	1/418 (0.2%)
1	H	0.41	0/252	1.27	4/385 (1.0%)
2	A	0.85	0/1630	0.95	0/2210
2	B	0.65	0/1607	0.87	0/2179
2	C	0.93	0/1618	1.02	2/2194 (0.1%)
2	D	0.74	0/1568	0.88	0/2126
All	All	0.76	0/7421	0.98	8/10234 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	6	DA	O4'-C1'-N9	-6.63	98.45	108.40
2	C	118	THR	N-CA-C	6.33	120.29	111.56
1	E	10	DC	P-O3'-C3'	5.99	129.19	120.20
1	H	6	DA	P-O5'-C5'	5.36	128.05	120.00
1	H	6	DA	N9-C1'-C2'	5.29	121.44	113.50
1	H	2	DG	P-O5'-C5'	5.23	127.85	120.00
1	G	6	DA	O4'-C1'-N9	-5.22	100.57	108.40
2	C	183	SER	N-CA-C	5.06	117.25	111.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	213	0	112	1	0
1	F	216	0	113	1	0
1	G	243	0	135	3	0
1	H	227	0	120	1	0
2	A	1577	0	1521	14	0
2	B	1550	0	1491	14	0
2	C	1554	0	1520	12	0
2	D	1522	0	1466	20	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	216	0	0	3	0
4	B	113	0	0	4	0
4	C	220	0	0	3	0
4	D	143	0	0	1	0
4	E	24	0	0	0	0
4	F	14	0	0	0	0
4	G	12	0	0	0	0
4	H	9	0	0	0	0
All	All	7857	0	6478	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:133:MET:SD	2:D:141[A]:CYS:SG	2.38	1.20
2:A:103:TYR:OH	2:A:264:LEU:HB3	1.61	1.00
2:C:115:HIS:HD2	4:C:586:HOH:O	1.61	0.83
2:A:103:TYR:HH	2:A:264:LEU:HB3	1.38	0.82
2:B:268:ASN:HD22	2:B:269[A]:SER:H	1.28	0.80
2:B:268:ASN:HD22	2:B:269[B]:SER:H	1.30	0.77
2:A:170[A]:THR:HG22	4:A:421:HOH:O	1.83	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:133:MET:CE	2:D:141[A]:CYS:SG	2.74	0.76
2:D:209:ARG:NE	2:D:209:ARG:HA	2.01	0.75
2:A:125[B]:THR:O	2:A:133[B]:MET:SD	2.48	0.72
2:D:192[A]:GLN:HE21	2:D:192[A]:GLN:H	1.37	0.71
2:A:166:SER:HA	2:A:169:MET:HE3	1.74	0.70
2:C:124:CYS:SG	2:C:133[A]:MET:SD	2.89	0.69
2:C:120:LYS:HG3	2:C:280:ARG:HB2	1.74	0.68
2:C:166:SER:HA	2:C:169:MET:HE3	1.74	0.67
2:B:121:SER:O	2:B:122:VAL:HG12	1.93	0.67
1:H:1:DC:H2'	1:H:2:DG:C8	2.28	0.67
1:E:10:DC:H2''	1:E:11:DC:O5'	1.96	0.66
2:C:115:HIS:CD2	4:C:586:HOH:O	2.42	0.64
2:B:248[B]:ARG:NH2	4:B:341:HOH:O	2.33	0.61
2:C:228:ASP:HB3	4:C:600:HOH:O	2.00	0.61
2:B:104:GLN:O	4:B:349:HOH:O	2.16	0.60
2:D:252:LEU:CD2	2:D:271:GLU:HG2	2.34	0.58
2:C:158:ARG:CZ	2:C:217[A]:VAL:HG21	2.34	0.58
2:B:207:ASP:HB2	4:B:405:HOH:O	2.05	0.57
2:D:252:LEU:HD23	2:D:271:GLU:HG2	1.87	0.57
2:B:165[B]:GLN:HB3	2:B:167:GLN:HE21	1.70	0.56
2:D:243[B]:MET:SD	2:D:247:ASN:ND2	2.78	0.56
2:A:175:ARG:HD3	2:A:191:PRO:O	2.06	0.55
1:F:2:DG:H2''	1:F:3:DG:C8	2.43	0.53
2:D:192[A]:GLN:HG3	2:D:214:HIS:CE1	2.44	0.53
2:D:192[A]:GLN:CG	2:D:214:HIS:CE1	2.93	0.52
2:B:248[B]:ARG:HD2	4:B:326:HOH:O	2.08	0.52
2:D:209:ARG:NE	2:D:209:ARG:CA	2.72	0.52
2:D:133:MET:HE1	2:D:141[A]:CYS:SG	2.48	0.51
2:C:120:LYS:HE3	2:C:280:ARG:HD3	1.92	0.50
2:B:165[A]:GLN:HB3	2:B:167:GLN:HE21	1.74	0.50
2:D:175:ARG:HD3	2:D:191:PRO:O	2.12	0.49
2:C:158:ARG:CZ	2:C:217[A]:VAL:CG2	2.91	0.48
2:A:110:ARG:HB2	4:A:384:HOH:O	2.14	0.48
2:D:202:ARG:HD3	4:D:345:HOH:O	2.14	0.47
2:A:116[B]:SER:HB3	4:A:389:HOH:O	2.14	0.47
1:G:11:DC:H2''	1:G:12:DG:C8	2.50	0.47
2:A:103:TYR:OH	2:A:264:LEU:CB	2.49	0.46
2:D:252:LEU:CD2	2:D:271:GLU:CG	2.94	0.46
1:G:3:DG:N7	2:D:120:LYS:NZ	2.62	0.45
2:C:119:ALA:O	2:C:279:GLY:HA3	2.17	0.45
2:D:192[A]:GLN:HG2	2:D:214:HIS:CE1	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:124:CYS:SG	2:A:133[B]:MET:SD	3.16	0.44
2:B:241:SER:HA	2:B:248[B]:ARG:HG3	2.00	0.44
2:D:175:ARG:HG3	2:D:192[B]:GLN:O	2.18	0.43
2:B:259:ASP:OD2	2:B:263:ASN:HB2	2.19	0.43
2:B:168:ARG:HH21	2:B:171:GLU:CD	2.27	0.42
2:A:182[A]:CYS:SG	2:A:184:ASP:OD2	2.78	0.42
2:A:102:THR:HG23	2:A:268[A]:ASN:ND2	2.34	0.42
2:D:103:TYR:CD1	2:D:266:GLY:HA2	2.55	0.42
1:G:4:DG:O6	2:D:120:LYS:NZ	2.48	0.41
2:A:119:ALA:O	2:A:279:GLY:HA3	2.21	0.41
2:B:175:ARG:HD3	2:B:191:PRO:O	2.20	0.41
2:A:256:THR:HG22	2:A:267:ARG:HG3	2.03	0.41
2:C:158:ARG:NH2	2:C:217[A]:VAL:HG21	2.35	0.41
2:C:175:ARG:HD3	2:C:191:PRO:O	2.22	0.40
2:D:243[B]:MET:HE1	2:D:247:ASN:HD21	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	205/200 (102%)	202 (98%)	3 (2%)	0	100	100
2	B	206/200 (103%)	200 (97%)	5 (2%)	1 (0%)	24	14
2	C	203/200 (102%)	198 (98%)	5 (2%)	0	100	100
2	D	199/200 (100%)	195 (98%)	4 (2%)	0	100	100
All	All	813/800 (102%)	795 (98%)	17 (2%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	122	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	180/179 (101%)	176 (98%)	4 (2%)	45	34
2	B	173/179 (97%)	169 (98%)	4 (2%)	44	33
2	C	181/179 (101%)	179 (99%)	2 (1%)	65	60
2	D	170/179 (95%)	168 (99%)	2 (1%)	63	57
All	All	704/716 (98%)	692 (98%)	12 (2%)	59	45

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	183[B]	SER
2	A	183[C]	SER
2	A	238[A]	CYS
2	A	238[B]	CYS
2	B	106	SER
2	B	149	SER
2	B	167	GLN
2	B	268	ASN
2	C	183	SER
2	C	243	MET
2	D	192[A]	GLN
2	D	192[B]	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	104	GLN
2	A	288	ASN
2	B	100	GLN
2	B	104	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	167	GLN
2	B	235	ASN
2	B	239	ASN
2	B	268	ASN
2	C	100	GLN
2	C	115	HIS
2	D	178	HIS
2	D	247	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	12/12 (100%)	0.22	1 (8%) 17 15	26, 35, 85, 86	0
1	F	11/12 (91%)	0.24	1 (9%) 15 13	27, 35, 62, 70	0
1	G	12/12 (100%)	0.28	1 (8%) 17 15	27, 43, 58, 67	0
1	H	12/12 (100%)	0.27	1 (8%) 17 15	29, 41, 65, 72	0
2	A	196/200 (98%)	-0.20	2 (1%) 79 80	7, 23, 40, 52	11 (5%)
2	B	197/200 (98%)	0.38	11 (5%) 30 29	14, 34, 58, 72	11 (5%)
2	C	193/200 (96%)	0.00	9 (4%) 36 35	10, 24, 43, 63	14 (7%)
2	D	194/200 (97%)	0.21	5 (2%) 57 57	12, 32, 51, 67	10 (5%)
All	All	827/848 (97%)	0.11	31 (3%) 45 45	7, 29, 53, 86	46 (5%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	225	VAL	8.5
2	C	226	GLY	3.8
2	C	284	THR	3.6
2	D	95	SER	3.3
2	C	96	SER	3.2
2	B	121	SER	3.1
2	B	122	VAL	3.0
2	B	289	LEU	2.9
2	C	121	SER	2.8
2	B	212	PHE	2.7
2	D	117	GLY	2.7
2	A	103	TYR	2.7
2	B	183	SER	2.6
2	B	290	ARG	2.6
2	B	228	ASP	2.5
2	B	225	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	119	ALA	2.5
1	E	11	DC	2.4
2	C	286	GLU	2.4
2	D	225	VAL	2.4
2	C	283	ARG	2.3
2	B	152	PRO	2.3
2	D	287	GLU	2.2
1	G	1	DC	2.2
2	D	119	ALA	2.2
2	C	287	GLU	2.2
1	H	12	DG	2.1
2	B	109	PHE	2.1
2	C	183	SER	2.1
2	A	288	ASN	2.0
1	F	2	DG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	1	1/1	0.98	0.03	23,23,23,23	0
3	ZN	B	1	1/1	0.99	0.02	26,26,26,26	0
3	ZN	C	1	1/1	0.99	0.02	23,23,23,23	0
3	ZN	D	1	1/1	0.99	0.02	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.