



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:13 PM UTC

PDB ID : 3D0Q / pdb_00003d0q
Title : Crystal structure of calG3 from Micromonospora echinospora determined in space group I222
Authors : Bitto, E.; Singh, S.; Bingman, C.A.; Wesenberg, G.E.; Phillips Jr., G.N.
Deposited on : 2008-05-02
Resolution : 2.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 5% 83% 11% • 5%
1	B	398	 5% 81% 12% • 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein CalG3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	Se	0	0	0
			2858	1809	512	528	1	8			
1	B	377	Total	C	N	O	S	Se	0	0	0
			2858	1809	512	528	1	8			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	GLY	-	expression tag	UNP Q8KND7
A	-20	HIS	-	expression tag	UNP Q8KND7
A	-19	HIS	-	expression tag	UNP Q8KND7
A	-18	HIS	-	expression tag	UNP Q8KND7
A	-17	HIS	-	expression tag	UNP Q8KND7
A	-16	HIS	-	expression tag	UNP Q8KND7
A	-15	HIS	-	expression tag	UNP Q8KND7
A	-14	HIS	-	expression tag	UNP Q8KND7
A	-13	HIS	-	expression tag	UNP Q8KND7
A	-12	HIS	-	expression tag	UNP Q8KND7
A	-11	HIS	-	expression tag	UNP Q8KND7
A	-10	SER	-	expression tag	UNP Q8KND7
A	-9	SER	-	expression tag	UNP Q8KND7
A	-8	GLY	-	expression tag	UNP Q8KND7
A	-7	HIS	-	expression tag	UNP Q8KND7
A	-6	ILE	-	expression tag	UNP Q8KND7
A	-5	GLU	-	expression tag	UNP Q8KND7
A	-4	GLY	-	expression tag	UNP Q8KND7
A	-3	ARG	-	expression tag	UNP Q8KND7
A	-2	HIS	-	expression tag	UNP Q8KND7
A	-1	MSE	-	expression tag	UNP Q8KND7
A	0	ARG	-	expression tag	UNP Q8KND7
A	1	VAL	-	expression tag	UNP Q8KND7
B	-21	GLY	-	expression tag	UNP Q8KND7
B	-20	HIS	-	expression tag	UNP Q8KND7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	HIS	-	expression tag	UNP Q8KND7
B	-18	HIS	-	expression tag	UNP Q8KND7
B	-17	HIS	-	expression tag	UNP Q8KND7
B	-16	HIS	-	expression tag	UNP Q8KND7
B	-15	HIS	-	expression tag	UNP Q8KND7
B	-14	HIS	-	expression tag	UNP Q8KND7
B	-13	HIS	-	expression tag	UNP Q8KND7
B	-12	HIS	-	expression tag	UNP Q8KND7
B	-11	HIS	-	expression tag	UNP Q8KND7
B	-10	SER	-	expression tag	UNP Q8KND7
B	-9	SER	-	expression tag	UNP Q8KND7
B	-8	GLY	-	expression tag	UNP Q8KND7
B	-7	HIS	-	expression tag	UNP Q8KND7
B	-6	ILE	-	expression tag	UNP Q8KND7
B	-5	GLU	-	expression tag	UNP Q8KND7
B	-4	GLY	-	expression tag	UNP Q8KND7
B	-3	ARG	-	expression tag	UNP Q8KND7
B	-2	HIS	-	expression tag	UNP Q8KND7
B	-1	MSE	-	expression tag	UNP Q8KND7
B	0	ARG	-	expression tag	UNP Q8KND7
B	1	VAL	-	expression tag	UNP Q8KND7

- Molecule 2 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (CCD ID: MPO) (formula: $C_7H_{15}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	22	Total	O	0	0
			22	22		
3	B	23	Total	O	0	0
			23	23		

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	106.65Å 119.27Å 155.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.75 – 2.79 26.75 – 2.79	Depositor EDS
% Data completeness (in resolution range)	98.9 (26.75-2.79) 98.9 (26.75-2.79)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.78 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.243 0.193 , 0.236	Depositor DCC
R_{free} test set	1270 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 27.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5787	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2913	0.91	4/3968 (0.1%)
1	B	0.68	1/2913 (0.0%)	0.94	4/3968 (0.1%)
All	All	0.68	1/5826 (0.0%)	0.92	8/7936 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	258	THR	CA-CB	5.87	1.60	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	77	ARG	CA-C-N	-5.68	114.08	119.76
1	B	77	ARG	C-N-CA	-5.68	114.08	119.76
1	B	97	VAL	N-CA-C	5.59	115.73	110.30
1	A	253	ILE	N-CA-CB	-5.34	106.34	112.21
1	A	97	VAL	N-CA-C	5.17	115.31	110.30
1	A	16	ILE	N-CA-C	5.06	115.49	110.23
1	A	312	HIS	N-CA-C	5.03	116.45	110.97
1	B	230	VAL	N-CA-C	5.02	115.79	110.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2858	0	2872	25	0
1	B	2858	0	2872	25	0
2	A	13	0	15	1	0
2	B	13	0	15	1	0
3	A	22	0	0	0	0
3	B	23	0	0	1	0
All	All	5787	0	5774	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285:GLY:HA2	1:A:302:LEU:CD1	2.00	0.92
1:A:285:GLY:HA2	1:A:302:LEU:HD13	1.58	0.84
1:A:31:ILE:HD11	1:A:45:LEU:HD13	1.63	0.81
1:B:225:PHE:HD2	1:B:307:ARG:HH11	1.38	0.71
1:A:67:ASN:OD1	1:A:69:ARG:HG3	1.93	0.69
1:A:285:GLY:CA	1:A:302:LEU:CD1	2.72	0.67
1:A:101:MSE:HE1	1:A:162:VAL:HG11	1.78	0.66
1:B:143:ARG:NH1	1:B:181:GLU:OE2	2.33	0.61
1:B:175:PRO:HD2	1:B:178:LEU:HD12	1.83	0.59
1:A:291:MSE:HE3	1:A:357:MSE:HE2	1.86	0.58
1:A:285:GLY:CA	1:A:302:LEU:HD13	2.33	0.57
1:A:231:GLU:N	1:A:232:PRO:HD2	2.19	0.57
1:A:101:MSE:CE	1:A:162:VAL:HG11	2.35	0.57
1:B:-1:MSE:HE1	1:B:374:ILE:HD11	1.86	0.56
1:B:294:ILE:HD11	1:B:323:ILE:HD12	1.88	0.56
1:B:231:GLU:HB2	1:B:232:PRO:HD3	1.88	0.55
1:A:101:MSE:HE3	1:A:127:ARG:HD3	1.88	0.54
1:A:294:ILE:HD11	1:A:323:ILE:HD12	1.91	0.53
1:A:175:PRO:HD2	1:A:178:LEU:HD12	1.91	0.53
1:B:139:ALA:HB3	1:B:313:THR:HB	1.92	0.52
1:A:244:PHE:HB2	1:A:263:VAL:HG22	1.92	0.51
1:B:-1:MSE:HE1	1:B:374:ILE:CD1	2.41	0.51
1:B:11:HIS:NE2	2:B:401:MPO:O2	2.44	0.50
1:A:67:ASN:ND2	1:A:70:PHE:HB2	2.27	0.50
1:A:138:SER:OG	1:A:139:ALA:N	2.35	0.49
1:B:217:MSE:HE2	1:B:222:LEU:HA	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:ALA:HB3	1:A:313:THR:HB	1.95	0.48
1:B:271:LEU:HD21	1:B:289:THR:HG23	1.95	0.47
1:B:231:GLU:CB	1:B:232:PRO:HD3	2.45	0.47
1:A:30:LEU:HD11	1:A:48:VAL:HG23	1.97	0.47
1:B:241:ASP:HA	3:B:407:HOH:O	2.15	0.46
1:B:141:ARG:NE	1:B:141:ARG:HA	2.31	0.46
1:A:273:THR:OG1	1:B:46:GLU:OE2	2.35	0.45
1:B:227:ILE:H	1:B:227:ILE:HG13	1.59	0.44
1:B:7:PRO:HG2	1:B:54:TYR:CZ	2.53	0.44
1:A:294:ILE:HG22	1:A:357:MSE:HE3	2.00	0.44
1:A:228:GLY:C	1:A:230:VAL:H	2.26	0.43
1:B:171:ILE:HD11	1:B:370:ILE:HD11	2.00	0.42
1:B:305:ASP:CG	1:B:306:PRO:HD2	2.44	0.42
1:B:30:LEU:HD11	1:B:48:VAL:HG23	2.01	0.42
1:B:217:MSE:HE3	1:B:284:HIS:NE2	2.35	0.42
1:B:3:PHE:HB2	1:B:31:ILE:HD13	2.02	0.41
1:A:101:MSE:CE	1:A:127:ARG:HB2	2.51	0.41
1:A:192:VAL:HA	1:A:193:PRO:HD2	1.96	0.41
1:A:11:HIS:NE2	2:A:401:MPO:H71	2.35	0.41
1:B:307:ARG:HE	1:B:307:ARG:HB2	1.55	0.40
1:B:92:VAL:O	1:B:95:PRO:HD2	2.21	0.40
1:A:101:MSE:HE1	1:A:127:ARG:HB2	2.03	0.40
1:B:239:GLU:OE2	1:B:239:GLU:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/398 (94%)	364 (97%)	10 (3%)	1 (0%)	36	66
1	B	375/398 (94%)	364 (97%)	11 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	750/796 (94%)	728 (97%)	21 (3%)	1 (0%)	48 77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/306 (97%)	285 (96%)	12 (4%)	28 63
1	B	297/306 (97%)	279 (94%)	18 (6%)	17 46
All	All	594/612 (97%)	564 (95%)	30 (5%)	21 54

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	VAL
1	A	63	VAL
1	A	69	ARG
1	A	72	GLU
1	A	77	ARG
1	A	143	ARG
1	A	230	VAL
1	A	259	LEU
1	A	284	HIS
1	A	355	GLU
1	A	357	MSE
1	A	373	ARG
1	B	31	ILE
1	B	47	VAL
1	B	137	GLN
1	B	151	SER
1	B	220	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	222	LEU
1	B	227	ILE
1	B	230	VAL
1	B	254	SER
1	B	261	ARG
1	B	266	VAL
1	B	279	THR
1	B	284	HIS
1	B	307	ARG
1	B	319	SER
1	B	329	SER
1	B	341	ILE
1	B	373	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	A	134	GLN
1	A	309	GLN
1	B	134	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	MPO	A	401	-	13,13,13	1.41	1 (7%)	17,17,17	1.61	4 (23%)
2	MPO	B	401	-	13,13,13	1.45	1 (7%)	17,17,17	2.34	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPO	A	401	-	-	6/7/15/15	0/1/1/1
2	MPO	B	401	-	-	4/7/15/15	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	MPO	C1-S1	4.07	1.83	1.77
2	A	401	MPO	C1-S1	3.88	1.83	1.77

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	MPO	O4-C6-C7	-5.93	99.00	111.77
2	B	401	MPO	C7-N1-C4	4.64	118.84	108.84
2	A	401	MPO	C7-N1-C4	3.71	116.82	108.84
2	B	401	MPO	C5-C4-N1	3.09	114.82	110.12
2	B	401	MPO	O2-S1-O1	-2.49	105.74	113.82
2	A	401	MPO	C6-C7-N1	2.40	113.77	110.12
2	B	401	MPO	O1-S1-C1	2.17	110.01	106.73
2	A	401	MPO	O3-S1-C1	2.12	110.16	106.00
2	A	401	MPO	O4-C6-C7	2.03	116.14	111.77

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	MPO	S1-C1-C2-C3
2	B	401	MPO	C1-C2-C3-N1
2	A	401	MPO	C2-C3-N1-C7
2	B	401	MPO	C2-C3-N1-C7
2	A	401	MPO	C2-C3-N1-C4
2	B	401	MPO	C2-C3-N1-C4
2	A	401	MPO	C2-C1-S1-O3
2	A	401	MPO	C2-C1-S1-O2
2	A	401	MPO	S1-C1-C2-C3
2	A	401	MPO	C1-C2-C3-N1

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	MPO	1	0
2	B	401	MPO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/398 (92%)	-0.32	1 (0%) 90 86	2, 9, 23, 34	0
1	B	369/398 (92%)	0.02	18 (4%) 35 27	2, 9, 27, 46	0
All	All	738/796 (92%)	-0.15	19 (2%) 57 47	2, 9, 24, 46	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	211	PRO	5.5
1	B	241	ASP	3.5
1	B	373	ARG	3.4
1	B	275	LEU	3.3
1	B	366	THR	3.3
1	B	272	HIS	2.9
1	B	271	LEU	2.7
1	B	243	ASP	2.7
1	B	213	VAL	2.6
1	B	210	ARG	2.6
1	A	79	ALA	2.4
1	B	227	ILE	2.4
1	B	251	LEU	2.3
1	B	263	VAL	2.3
1	B	220	ILE	2.2
1	B	369	ARG	2.2
1	B	244	PHE	2.1
1	B	258	THR	2.0
1	B	362	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MPO	B	401	13/13	0.84	0.18	23,35,53,56	0
2	MPO	A	401	13/13	0.95	0.17	18,27,41,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.