



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 11:16 PM UTC

PDB ID : 4D00 / pdb_00004d00
Title : Haemagglutinin of H10N8 Influenza Virus Isolated from Humans in Complex with Human Receptor Analogue 6'SLN
Authors : Xiong, X.; Haire, L.F.; Vachieri, S.G.; Collins, P.J.; Walker, P.A.; Martin, S.R.; Zhang, Y.; McCauley, J.W.; Gamblin, S.J.; Skehel, J.J.
Deposited on : 2014-04-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	 2% 91% 6% .
1	C	326	 % 93% 5% .
1	E	326	 % 92% 6% .
2	B	183	 2% 89% 7% . .
2	D	183	 4% 90% 6% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	183	
3	G	3	
3	I	3	
3	K	3	
4	H	2	
4	J	2	
4	L	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	1320	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 12576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HAEMAGGLUTININ HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2444	1511	450	466	17			
1	C	321	Total	C	N	O	S	0	0	0
			2457	1518	452	470	17			
1	E	319	Total	C	N	O	S	0	0	0
			2444	1511	450	466	17			

- Molecule 2 is a protein called HAEMAGGLUTININ HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	175	Total	C	N	O	S	0	0	0
			1409	871	244	286	8			
2	D	176	Total	C	N	O	S	0	0	0
			1416	876	245	287	8			
2	F	173	Total	C	N	O	S	0	0	0
			1394	862	241	283	8			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	3	Total	C	N	O	0	0	0
			46	25	2	19			
3	I	3	Total	C	N	O	0	0	0
			46	25	2	19			
3	K	3	Total	C	N	O	0	0	0
			46	25	2	19			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	J	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	L	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	E	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ni	0	0
			1	1		
6	B	1	Total	Ni	0	0
			1	1		
6	D	1	Total	Ni	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	180	Total	O	0	0
			180	180		
7	A	1	Total	O	0	0
			1	1		
7	B	94	Total	O	0	0
			94	94		
7	C	151	Total	O	0	0
			151	151		
7	C	1	Total	O	0	0
			1	1		
7	D	81	Total	O	0	0
			81	81		
7	E	131	Total	O	0	0
			131	131		
7	E	4	Total	O	0	0
			4	4		
7	F	60	Total	O	0	0
			60	60		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HAEMAGGLUTININ HA1




- Molecule 1: HAEMAGGLUTININ HA1




- Molecule 1: HAEMAGGLUTININ HA1




- Molecule 2: HAEMAGGLUTININ HA1



- Molecule 2: HAEMAGGLUTININ HA1



- Molecule 2: HAEMAGGLUTININ HA1



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.60Å 183.17Å 192.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	132.58 – 2.50 132.58 – 2.50	Depositor EDS
% Data completeness (in resolution range)	92.7 (132.58-2.50) 92.7 (132.58-2.50)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.209 , 0.235 0.211 , 0.237	Depositor DCC
R_{free} test set	5013 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.020 for -h,l,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12576	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, NAG, SIA, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2494	0.64	0/3379
1	C	0.45	0/2507	0.65	0/3398
1	E	0.44	0/2494	0.64	0/3379
2	B	0.48	0/1435	0.72	0/1938
2	D	0.47	0/1442	0.72	1/1948 (0.1%)
2	F	0.48	0/1419	0.71	0/1915
All	All	0.46	0/11791	0.67	1/15957 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	173	ILE	CB-CA-C	-5.30	105.41	112.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2444	0	2395	14	0
1	C	2457	0	2404	12	0
1	E	2444	0	2395	10	0
2	B	1409	0	1308	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1416	0	1317	7	0
2	F	1394	0	1295	5	1
3	G	46	0	40	0	0
3	I	46	0	40	0	0
3	K	46	0	40	1	0
4	H	28	0	25	0	0
4	J	28	0	25	0	0
4	L	28	0	25	0	0
5	A	28	0	26	1	0
5	C	28	0	26	0	0
5	E	28	0	26	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	A	181	0	0	5	0
7	B	94	0	0	0	0
7	C	152	0	0	2	1
7	D	81	0	0	0	0
7	E	135	0	0	2	0
7	F	60	0	0	0	0
All	All	12576	0	11387	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:162:ARG:HD2	1:C:235:ASN:OD1	1.59	1.00
1:C:158:THR:N	7:C:2090:HOH:O	2.00	0.94
2:D:173:ILE:O	2:D:173:ILE:HG22	1.79	0.80
1:C:162:ARG:CD	1:C:235:ASN:OD1	2.34	0.74
1:C:104:GLU:HG3	1:C:256:ARG:HH21	1.60	0.67
1:C:205:ASN:HB3	7:C:2111:HOH:O	1.98	0.62
2:D:169:ASN:OD1	2:D:174:ASN:ND2	2.35	0.59
2:D:173:ILE:O	2:D:173:ILE:CG2	2.50	0.57
1:A:109:ASN:CG	7:A:2088:HOH:O	2.48	0.56
1:E:5:LEU:HD22	2:F:119:TYR:HA	1.88	0.55
1:A:162:ARG:NH1	5:A:1319:NAG:O5	2.40	0.55
1:E:128:ARG:NH1	1:E:136:ASN:OD1	2.40	0.55
1:A:226:HIS:HD2	7:A:2022:HOH:O	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:268:ASN:ND2	7:E:2026:HOH:O	2.42	0.52
1:C:256:ARG:NH2	2:D:64:GLU:OE2	2.43	0.50
1:A:56:PRO:HA	1:A:59:MET:HE2	1.93	0.50
1:A:57:ILE:CD1	1:A:95:VAL:HG23	2.43	0.49
1:E:57:ILE:CD1	1:E:95:VAL:HG23	2.42	0.49
1:C:57:ILE:CD1	1:C:95:VAL:HG23	2.43	0.48
1:A:268:ASN:ND2	7:A:2031:HOH:O	2.42	0.48
1:E:56:PRO:HA	1:E:59:MET:HE2	1.95	0.48
1:A:67:ASP:OD1	1:A:132:ARG:NH1	2.46	0.48
1:A:205:ASN:HB2	7:A:2123:HOH:O	2.14	0.48
1:C:-1:ASP:HB2	1:C:0:PRO:CD	2.44	0.48
2:B:75:HIS:CE1	1:C:101:LYS:HG2	2.49	0.48
1:C:56:PRO:HA	1:C:59:MET:HE2	1.95	0.47
2:D:30:GLN:HE22	2:D:146:ASP:H	1.62	0.47
2:F:30:GLN:HE22	2:F:146:ASP:H	1.63	0.47
2:F:9:PHE:CD1	2:F:10:LEU:HD13	2.51	0.46
2:B:30:GLN:HE22	2:B:146:ASP:H	1.62	0.46
1:A:177:HIS:O	1:A:213:ARG:NH2	2.49	0.46
1:A:5:LEU:HD22	2:B:119:TYR:HA	1.99	0.44
2:B:48:ILE:CD1	2:B:107:THR:HG23	2.48	0.44
2:B:173:ILE:HG12	2:D:164:GLU:HG3	2.00	0.44
2:F:9:PHE:CE1	2:F:10:LEU:HD13	2.52	0.44
1:E:136:ASN:ND2	3:K:1:NAG:H82	2.34	0.43
2:F:48:ILE:CD1	2:F:107:THR:HG23	2.49	0.43
2:D:48:ILE:CD1	2:D:107:THR:HG23	2.48	0.43
1:E:93:VAL:HB	7:E:2055:HOH:O	2.18	0.42
1:A:178:PRO:O	1:A:213:ARG:NH2	2.53	0.42
1:A:198:GLY:HA3	1:E:213:ARG:HG2	2.02	0.41
1:A:109:ASN:CB	7:A:2088:HOH:O	2.69	0.41
1:E:57:ILE:CD1	1:E:95:VAL:CG2	2.98	0.41
1:A:57:ILE:CD1	1:A:95:VAL:CG2	2.99	0.41
1:C:-1:ASP:HB2	1:C:0:PRO:HD3	2.02	0.41
1:C:209:VAL:HG11	1:E:205:ASN:HB2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:32:THR:N	7:C:2090:HOH:O[3_545]	1.94	0.26

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/326 (97%)	312 (98%)	5 (2%)	0	100	100
1	C	319/326 (98%)	314 (98%)	5 (2%)	0	100	100
1	E	317/326 (97%)	312 (98%)	5 (2%)	0	100	100
2	B	173/183 (94%)	169 (98%)	4 (2%)	0	100	100
2	D	174/183 (95%)	170 (98%)	4 (2%)	0	100	100
2	F	171/183 (93%)	167 (98%)	4 (2%)	0	100	100
All	All	1471/1527 (96%)	1444 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/275 (98%)	265 (98%)	5 (2%)	50	76
1	C	271/275 (98%)	268 (99%)	3 (1%)	65	84
1	E	270/275 (98%)	262 (97%)	8 (3%)	36	64
2	B	149/156 (96%)	142 (95%)	7 (5%)	23	47
2	D	150/156 (96%)	147 (98%)	3 (2%)	48	75
2	F	147/156 (94%)	143 (97%)	4 (3%)	39	67
All	All	1257/1293 (97%)	1227 (98%)	30 (2%)	43	70

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	133	ASN
1	A	148	LYS
1	A	184	LYS
1	A	196	SER
2	B	56	VAL
2	B	58	LYS
2	B	60	ASN
2	B	113	SER
2	B	135	LYS
2	B	161	GLN
2	B	173	ILE
1	C	53	ASN
1	C	148	LYS
1	C	196	SER
2	D	39	LYS
2	D	113	SER
2	D	135	LYS
1	E	2	LYS
1	E	8	HIS
1	E	12	ASN
1	E	148	LYS
1	E	152	GLN
1	E	196	SER
1	E	200	SER
1	E	305	ARG
2	F	10	LEU
2	F	60	ASN
2	F	113	SER
2	F	135	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (30) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	226	HIS
1	A	231	GLN
1	A	268	ASN
1	A	297	GLN
1	A	303	ASN
2	B	30	GLN
2	B	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	75	HIS
2	B	174	ASN
1	C	53	ASN
1	C	156	GLN
1	C	205	ASN
1	C	231	GLN
2	D	26	HIS
2	D	30	GLN
2	D	42	GLN
2	D	60	ASN
2	D	95	GLN
1	E	8	HIS
1	E	156	GLN
1	E	182	GLN
1	E	205	ASN
1	E	226	HIS
1	E	231	GLN
1	E	268	ASN
1	E	297	GLN
2	F	12	ASN
2	F	30	GLN
2	F	42	GLN
2	F	95	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	G	1	3	15,15,15	0.48	0	21,21,21	1.48	4 (19%)
3	GAL	G	2	3	11,11,12	0.43	0	15,15,17	0.82	0
3	SIA	G	3	3	20,20,21	0.58	0	21,28,31	1.46	4 (19%)
4	NAG	H	1	4,6,2	14,14,15	0.53	0	17,19,21	1.06	1 (5%)
4	NAG	H	2	4	14,14,15	0.55	0	17,19,21	1.45	3 (17%)
3	NAG	I	1	3	15,15,15	0.46	0	21,21,21	1.29	3 (14%)
3	GAL	I	2	3	11,11,12	0.37	0	15,15,17	0.85	1 (6%)
3	SIA	I	3	3	20,20,21	0.55	0	21,28,31	1.34	5 (23%)
4	NAG	J	1	4,6,2	14,14,15	0.63	0	17,19,21	1.37	1 (5%)
4	NAG	J	2	4	14,14,15	0.51	0	17,19,21	1.71	3 (17%)
3	NAG	K	1	3	15,15,15	0.55	0	21,21,21	1.56	3 (14%)
3	GAL	K	2	3	11,11,12	0.36	0	15,15,17	1.57	3 (20%)
3	SIA	K	3	3	20,20,21	0.53	0	21,28,31	1.31	5 (23%)
4	NAG	L	1	4,6,2	14,14,15	0.69	0	17,19,21	1.05	1 (5%)
4	NAG	L	2	4	14,14,15	0.51	0	17,19,21	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3	-	1/6/26/26	0/1/1/1
3	GAL	G	2	3	-	0/2/19/22	0/1/1/1
3	SIA	G	3	3	-	0/18/34/38	0/1/1/1
4	NAG	H	1	4,6,2	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1
3	NAG	I	1	3	-	4/6/26/26	0/1/1/1
3	GAL	I	2	3	-	0/2/19/22	0/1/1/1
3	SIA	I	3	3	-	0/18/34/38	0/1/1/1
4	NAG	J	1	4,6,2	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	2/6/23/26	0/1/1/1
3	NAG	K	1	3	-	2/6/26/26	0/1/1/1
3	GAL	K	2	3	-	0/2/19/22	0/1/1/1
3	SIA	K	3	3	-	1/18/34/38	0/1/1/1
4	NAG	L	1	4,6,2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	L	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	J	2	NAG	C1-O5-C5	5.21	119.16	112.19
4	J	1	NAG	C1-O5-C5	4.21	117.83	112.19
3	K	2	GAL	C1-C2-C3	4.09	115.59	109.64
4	H	2	NAG	C4-C3-C2	3.74	116.49	111.02
3	K	1	NAG	C3-C2-N2	-3.58	104.03	110.62
3	G	1	NAG	C1-C2-N2	-3.44	106.75	110.73
3	G	3	SIA	O6-C2-C1	3.41	114.15	107.72
3	G	1	NAG	C4-C3-C2	3.30	115.21	110.40
3	I	3	SIA	C4-C5-N5	-3.20	104.14	110.44
3	I	1	NAG	C1-C2-N2	-3.06	107.19	110.73
3	I	1	NAG	C8-C7-N2	2.96	121.03	116.12
4	L	1	NAG	C1-O5-C5	2.96	116.15	112.19
3	K	1	NAG	C8-C7-N2	2.95	121.01	116.12
4	J	2	NAG	C1-C2-N2	2.89	114.98	110.43
3	K	3	SIA	C4-C5-N5	-2.85	104.83	110.44
3	G	3	SIA	O1B-C1-C2	2.83	120.08	112.71
3	G	1	NAG	C3-C4-C5	2.72	115.16	110.23
3	K	1	NAG	C1-C2-N2	-2.69	107.61	110.73
3	G	1	NAG	C3-C2-N2	-2.66	105.72	110.62
3	K	3	SIA	O1B-C1-C2	2.64	119.58	112.71
3	G	3	SIA	C4-C5-N5	-2.64	105.23	110.44
3	K	3	SIA	O1A-C1-C2	-2.52	117.40	122.85
3	I	3	SIA	O1B-C1-C2	2.52	119.26	112.71
3	K	3	SIA	O6-C2-C1	2.47	112.38	107.72
4	J	2	NAG	C4-C3-C2	-2.38	107.52	111.02
3	I	3	SIA	O6-C2-C1	2.37	112.19	107.72
4	H	2	NAG	O5-C1-C2	-2.25	107.81	111.29
4	H	2	NAG	C3-C4-C5	2.24	114.30	110.23
4	H	1	NAG	O5-C1-C2	-2.15	107.96	111.29
3	I	3	SIA	O1A-C1-C2	-2.11	118.28	122.85
3	G	3	SIA	C6-C5-N5	-2.10	107.56	110.91
3	K	3	SIA	C6-C5-N5	-2.09	107.56	110.91
3	I	1	NAG	C3-C2-N2	-2.08	106.79	110.62
3	K	2	GAL	C1-O5-C5	2.07	114.96	112.19
3	I	3	SIA	C6-C5-N5	-2.04	107.65	110.91
3	K	2	GAL	C2-C3-C4	2.03	114.43	110.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	2	GAL	O5-C1-C2	-2.00	106.02	110.79

There are no chirality outliers.

All (14) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	J	2	NAG	C4-C5-C6-O6
4	H	2	NAG	C4-C5-C6-O6
3	I	1	NAG	O5-C5-C6-O6
4	H	2	NAG	O5-C5-C6-O6
4	J	2	NAG	O5-C5-C6-O6
3	I	1	NAG	C4-C5-C6-O6
3	I	1	NAG	C8-C7-N2-C2
3	I	1	NAG	O7-C7-N2-C2
3	K	1	NAG	C8-C7-N2-C2
3	K	1	NAG	O7-C7-N2-C2
4	L	2	NAG	C4-C5-C6-O6
4	L	2	NAG	O5-C5-C6-O6
3	G	1	NAG	C1-C2-N2-C7
3	K	3	SIA	O1A-C1-C2-C3


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	K	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	C	1320	1	14,14,15	0.45	0	17,19,21	1.33	1 (5%)
5	NAG	A	1319	1	14,14,15	0.46	0	17,19,21	1.43	3 (17%)
5	NAG	E	1320	1	14,14,15	0.36	0	17,19,21	2.45	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	1320	1	14,14,15	0.79	0	17,19,21	1.53	3 (17%)
5	NAG	C	1319	1	14,14,15	0.43	0	17,19,21	1.49	4 (23%)
5	NAG	E	1319	1	14,14,15	0.29	0	17,19,21	1.16	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1320	1	-	0/6/23/26	0/1/1/1
5	NAG	A	1319	1	-	2/6/23/26	0/1/1/1
5	NAG	E	1320	1	-	2/6/23/26	0/1/1/1
5	NAG	A	1320	1	1/1/5/7	3/6/23/26	0/1/1/1
5	NAG	C	1319	1	-	3/6/23/26	0/1/1/1
5	NAG	E	1319	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	1320	NAG	C1-O5-C5	7.56	122.32	112.19
5	C	1320	NAG	C1-O5-C5	4.77	118.58	112.19
5	E	1320	NAG	C8-C7-N2	3.95	122.67	116.12
5	C	1319	NAG	C1-O5-C5	3.55	116.94	112.19
5	A	1320	NAG	O5-C5-C4	-3.17	103.11	110.83
5	A	1319	NAG	C8-C7-N2	2.95	121.02	116.12
5	C	1319	NAG	C8-C7-N2	2.83	120.82	116.12
5	E	1319	NAG	C1-O5-C5	2.80	115.94	112.19
5	A	1320	NAG	C1-C2-N2	-2.37	106.70	110.43
5	E	1320	NAG	C2-N2-C7	2.36	126.06	122.90
5	E	1320	NAG	C1-C2-N2	-2.32	106.78	110.43
5	C	1319	NAG	C1-C2-N2	2.31	114.07	110.43
5	A	1319	NAG	C1-O5-C5	2.28	115.25	112.19
5	E	1320	NAG	O7-C7-C8	-2.28	118.00	122.05
5	E	1320	NAG	C4-C3-C2	-2.22	107.77	111.02
5	A	1319	NAG	O5-C1-C2	-2.18	107.92	111.29
5	A	1320	NAG	C4-C3-C2	2.16	114.19	111.02
5	C	1319	NAG	O7-C7-C8	-2.06	118.39	122.05

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	1320	NAG	C1

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1320	NAG	C3-C2-N2-C7
5	A	1319	NAG	C8-C7-N2-C2
5	A	1319	NAG	O7-C7-N2-C2
5	C	1319	NAG	C8-C7-N2-C2
5	C	1319	NAG	O7-C7-N2-C2
5	E	1320	NAG	C8-C7-N2-C2
5	E	1320	NAG	O7-C7-N2-C2
5	A	1320	NAG	O5-C5-C6-O6
5	A	1320	NAG	C4-C5-C6-O6
5	C	1319	NAG	O5-C5-C6-O6
5	E	1319	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1319	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/326 (97%)	-0.12	6 (1%) 66 62	21, 29, 44, 56	0
1	C	321/326 (98%)	-0.04	4 (1%) 76 73	22, 32, 44, 56	0
1	E	319/326 (97%)	-0.14	2 (0%) 85 83	22, 30, 43, 54	0
2	B	175/183 (95%)	-0.07	4 (2%) 61 57	20, 28, 44, 71	0
2	D	176/183 (96%)	-0.09	8 (4%) 38 33	20, 27, 45, 79	0
2	F	173/183 (94%)	-0.08	7 (4%) 42 38	20, 27, 44, 82	0
All	All	1483/1527 (97%)	-0.09	31 (2%) 63 59	20, 29, 44, 82	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	173	ILE	8.6
1	C	-2	ALA	5.5
2	D	176	VAL	4.9
1	C	270	CYS	4.3
2	F	59	THR	3.6
2	B	60	ASN	3.4
1	C	130	CYS	3.3
2	D	58	LYS	3.3
1	E	270	CYS	3.2
2	B	175	PRO	3.0
2	B	59	THR	2.9
2	D	59	THR	2.9
1	A	0	PRO	2.9
2	F	60	ASN	2.8
2	F	172	ASN	2.8
2	F	58	LYS	2.8
1	C	205	ASN	2.7
2	D	60	ASN	2.7
1	A	12	ASN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	109	ASN	2.5
1	A	270	CYS	2.4
1	E	148	LYS	2.4
1	A	305	ARG	2.4
2	D	167	LEU	2.4
2	D	55	LEU	2.2
2	D	56	VAL	2.2
2	F	56	VAL	2.1
2	B	58	LYS	2.1
2	F	11	GLU	2.0
2	D	175	PRO	2.0
1	A	53	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	J	2	14/15	0.59	0.25	63,71,77,81	0
3	NAG	I	1	15/15	0.61	0.25	87,99,101,102	0
3	NAG	G	1	15/15	0.65	0.27	93,105,107,109	0
4	NAG	L	2	14/15	0.71	0.21	62,70,74,74	0
3	GAL	I	2	11/12	0.80	0.15	54,80,90,91	0
3	NAG	K	1	15/15	0.81	0.20	65,74,76,78	0
4	NAG	H	2	14/15	0.83	0.17	61,69,74,75	0
3	GAL	K	2	11/12	0.84	0.13	46,56,60,62	0
3	GAL	G	2	11/12	0.85	0.15	50,72,81,84	0
4	NAG	J	1	14/15	0.91	0.10	32,36,45,52	0
4	NAG	L	1	14/15	0.93	0.09	30,36,42,51	0
4	NAG	H	1	14/15	0.93	0.10	32,37,44,51	0
3	SIA	G	3	20/21	0.95	0.08	34,36,40,41	0
3	SIA	I	3	20/21	0.96	0.06	35,39,45,46	0
3	SIA	K	3	20/21	0.96	0.07	35,38,40,41	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	1320	14/15	0.68	0.19	49,58,59,60	0
5	NAG	A	1320	14/15	0.74	0.21	58,67,69,71	0
5	NAG	C	1319	14/15	0.78	0.18	52,56,65,70	0
5	NAG	E	1320	14/15	0.84	0.14	44,51,54,54	0
5	NAG	A	1319	14/15	0.85	0.12	39,43,48,51	0
5	NAG	E	1319	14/15	0.88	0.11	41,44,48,49	0
6	NI	D	1179	1/1	0.96	0.04	39,39,39,39	0
6	NI	B	1178	1/1	0.98	0.05	38,38,38,38	0
6	NI	A	1321	1/1	0.98	0.03	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.