



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:16 AM UTC

PDB ID : 4D08 / pdb_00004d08
Title : PDE2a catalytic domain in complex with a brain penetrant inhibitor
Authors : Buijnsters, P.; Andres, J.I.; DeAngelis, M.; Langlois, X.; Rombouts, F.; Sanderson, W.; Tresadern, G.; Trabanco, A.; VanHoof, G.; VanRoosbroeck, Y.
Deposited on : 2014-04-24
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	5% 94% • •
1	B	353	9% 93% • 5%
1	C	353	14% 91% • 7%
1	D	353	5% 94% • 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11498 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CGMP-DEPENDENT 3', 5'-CYCLIC PHOSPHODIESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	3	0
			2736	1743	463	505	25			
1	B	337	Total	C	N	O	S	0	0	0
			2699	1723	458	493	25			
1	C	327	Total	C	N	O	S	0	0	0
			2606	1665	440	476	25			
1	D	335	Total	C	N	O	S	0	0	0
			2691	1718	457	491	25			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	576	MET	-	expression tag	UNP O00408
A	577	GLY	-	expression tag	UNP O00408
A	922	ARG	-	expression tag	UNP O00408
A	923	HIS	-	expression tag	UNP O00408
A	924	HIS	-	expression tag	UNP O00408
A	925	HIS	-	expression tag	UNP O00408
A	926	HIS	-	expression tag	UNP O00408
A	927	HIS	-	expression tag	UNP O00408
A	928	HIS	-	expression tag	UNP O00408
B	576	MET	-	expression tag	UNP O00408
B	577	GLY	-	expression tag	UNP O00408
B	922	ARG	-	expression tag	UNP O00408
B	923	HIS	-	expression tag	UNP O00408
B	924	HIS	-	expression tag	UNP O00408
B	925	HIS	-	expression tag	UNP O00408
B	926	HIS	-	expression tag	UNP O00408
B	927	HIS	-	expression tag	UNP O00408
B	928	HIS	-	expression tag	UNP O00408
C	576	MET	-	expression tag	UNP O00408
C	577	GLY	-	expression tag	UNP O00408
C	922	ARG	-	expression tag	UNP O00408

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	923	HIS	-	expression tag	UNP O00408
C	924	HIS	-	expression tag	UNP O00408
C	925	HIS	-	expression tag	UNP O00408
C	926	HIS	-	expression tag	UNP O00408
C	927	HIS	-	expression tag	UNP O00408
C	928	HIS	-	expression tag	UNP O00408
D	576	MET	-	expression tag	UNP O00408
D	577	GLY	-	expression tag	UNP O00408
D	922	ARG	-	expression tag	UNP O00408
D	923	HIS	-	expression tag	UNP O00408
D	924	HIS	-	expression tag	UNP O00408
D	925	HIS	-	expression tag	UNP O00408
D	926	HIS	-	expression tag	UNP O00408
D	927	HIS	-	expression tag	UNP O00408
D	928	HIS	-	expression tag	UNP O00408

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is 1-(5-butoxypyridin-3-yl)-4-methyl-8-(morpholin-4-ylmethyl)[1,2,4]triazolo[4,3-a]quinoxaline (CCD ID: Q2T) (formula: C₂₄H₂₈N₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			32	24	6	2		
3	B	1	Total	C	N	O	0	0
			32	24	6	2		
3	C	1	Total	C	N	O	0	0
			32	24	6	2		
3	D	1	Total	C	N	O	0	0
			32	24	6	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	173	Total	O	0	0
			173	173		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	168	Total 168	O 168	0	0
5	C	130	Total 130	O 130	0	0
5	D	159	Total 159	O 159	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CGMP-DEPENDENT 3', 5'-CYCLIC PHOSPHODIESTERASE




- Molecule 1: CGMP-DEPENDENT 3', 5'-CYCLIC PHOSPHODIESTERASE



- Molecule 1: CGMP-DEPENDENT 3', 5'-CYCLIC PHOSPHODIESTERASE



- Molecule 1: CGMP-DEPENDENT 3', 5'-CYCLIC PHOSPHODIESTERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	56.04Å 74.15Å 92.77Å 109.04° 91.73° 91.34°	Depositor
Resolution (Å)	46.34 – 1.90 46.34 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.0 (46.34-1.90) 92.1 (46.34-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.213 , 0.247 (Not available) , 0.263	Depositor DCC
R_{free} test set	5177 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11498	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, Q2T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2806	0.68	0/3800
1	B	0.44	0/2766	0.69	0/3746
1	C	0.43	0/2671	0.68	0/3619
1	D	0.44	0/2756	0.68	0/3728
All	All	0.44	0/10999	0.68	0/14893

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2736	0	2610	3	0
1	B	2699	0	2581	3	0
1	C	2606	0	2483	2	0
1	D	2691	0	2582	1	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	32	0	28	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	32	0	28	0	0
3	C	32	0	28	0	0
3	D	32	0	28	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	173	0	0	0	0
5	B	168	0	0	0	0
5	C	130	0	0	0	0
5	D	159	0	0	0	0
All	All	11498	0	10368	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:645:MET:HE1	1:A:740:THR:HG21	1.85	0.58
1:A:626:MET:HE1	1:A:690:PHE:CD1	2.50	0.47
1:B:864:GLU:OE2	1:B:892:ARG:NH2	2.48	0.46
1:A:596:ILE:HG21	1:A:600:PHE:CE1	2.54	0.42
1:C:660:HIS:O	1:C:664:VAL:HG23	2.20	0.41
1:D:834:GLY:HA3	1:D:848:MET:O	2.20	0.41
1:B:855:ILE:N	1:B:856:PRO:HD2	2.35	0.41
1:C:856:PRO:HG3	1:C:902:PHE:CE1	2.57	0.40
1:B:778:LYS:N	1:B:778:LYS:HD3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/353 (96%)	338 (99%)	2 (1%)	0	100	100
1	B	335/353 (95%)	333 (99%)	2 (1%)	0	100	100
1	C	325/353 (92%)	320 (98%)	5 (2%)	0	100	100
1	D	331/353 (94%)	330 (100%)	1 (0%)	0	100	100
All	All	1331/1412 (94%)	1321 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/319 (89%)	283 (100%)	0	100	100
1	B	289/319 (91%)	286 (99%)	3 (1%)	68	70
1	C	278/319 (87%)	276 (99%)	2 (1%)	76	78
1	D	289/319 (91%)	288 (100%)	1 (0%)	86	88
All	All	1139/1276 (89%)	1133 (100%)	6 (0%)	81	84

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	585	LEU
1	B	683	ASP
1	B	916	LEU
1	C	722	GLU
1	C	786	VAL
1	D	633	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	733	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	739	ASN
1	A	794	GLN
1	A	859	GLN
1	B	591	GLN
1	B	679	ASN
1	B	842	ASN
1	C	624	GLN
1	C	708	GLN
1	D	591	GLN
1	D	627	ASN
1	D	630	ASN
1	D	733	GLN
1	D	842	ASN
1	D	891	ASN
1	D	911	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	Q2T	A	1918	-	35,36,36	0.22	0	47,50,50	0.63	1 (2%)
3	Q2T	D	1922	-	35,36,36	0.18	0	47,50,50	0.54	0
3	Q2T	B	1917	-	35,36,36	0.22	0	47,50,50	0.56	0
3	Q2T	C	1917	-	35,36,36	0.19	0	47,50,50	0.61	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	Q2T	A	1918	-	-	3/13/21/21	0/5/5/5
3	Q2T	D	1922	-	-	2/13/21/21	0/5/5/5
3	Q2T	B	1917	-	-	2/13/21/21	0/5/5/5
3	Q2T	C	1917	-	-	4/13/21/21	0/5/5/5

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1918	Q2T	C3-N2-C9	2.14	136.52	133.30
3	C	1917	Q2T	C3-N2-C9	2.01	136.33	133.30

There are no chirality outliers.


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1917	Q2T	C12-C15-N16-C17
3	C	1917	Q2T	C12-C15-N16-C21
3	A	1918	Q2T	C12-C15-N16-C21
3	A	1918	Q2T	C12-C15-N16-C17
3	B	1917	Q2T	C23-C22-C9-N8
3	D	1922	Q2T	C23-C22-C9-N8
3	C	1917	Q2T	C23-C22-C9-N8
3	A	1918	Q2T	C23-C22-C9-N8
3	C	1917	Q2T	C27-C22-C9-N8
3	B	1917	Q2T	C27-C22-C9-N8
3	D	1922	Q2T	C27-C22-C9-N8


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	339/353 (96%)	0.57	16 (4%)	36	39	8, 21, 33, 46	3 (0%)
1	B	337/353 (95%)	0.67	32 (9%)	14	14	12, 21, 37, 47	0
1	C	327/353 (92%)	1.04	51 (15%)	5	5	12, 27, 45, 61	0
1	D	335/353 (94%)	0.62	17 (5%)	33	36	12, 21, 32, 39	0
All	All	1338/1412 (94%)	0.72	116 (8%)	16	17	8, 22, 37, 61	3 (0%)

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	791	ASN	5.1
1	C	784	ALA	4.6
1	C	789	ASP	4.5
1	C	590	ILE	4.4
1	C	786	VAL	4.3
1	D	584	LYS	4.3
1	A	747	ASP	4.2
1	C	681	LEU	4.1
1	B	629	ILE	4.0
1	B	588	ASP	3.8
1	C	595	ALA	3.8
1	B	630	ASN	3.7
1	C	785	GLU	3.7
1	B	710	ALA	3.6
1	B	748	HIS	3.6
1	A	636	CYS	3.5
1	C	790	ARG	3.5
1	C	593	VAL	3.4
1	B	580	ASP	3.4
1	B	589	GLY	3.4
1	D	595	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	792	ASN	3.4
1	C	896	THR	3.3
1	C	617	MET	3.2
1	B	747	ASP	3.2
1	C	748	HIS	3.2
1	C	788	TYR	3.2
1	D	594	ALA	3.2
1	B	627	ASN	3.2
1	B	634	ILE	3.1
1	C	591	GLN	3.1
1	C	631	ASN	3.1
1	C	793	LYS	3.1
1	B	722	GLU	3.1
1	D	585	LEU	3.0
1	A	785	GLU	3.0
1	C	684	ILE	3.0
1	A	895	TRP	3.0
1	B	914	ASP	3.0
1	C	627	ASN	3.0
1	A	892	ARG	2.9
1	B	631	ASN	2.9
1	B	636	CYS	2.9
1	D	894	HIS	2.9
1	C	747	ASP	2.9
1	B	632	TYR	2.8
1	D	918	GLU	2.8
1	C	795	HIS	2.8
1	D	586	LEU	2.7
1	C	680	TYR	2.7
1	B	723	GLY	2.7
1	D	617	MET	2.7
1	C	613	ASP	2.6
1	D	920	TYR	2.6
1	C	787	GLY	2.6
1	C	678	THR	2.6
1	C	894	HIS	2.6
1	C	839	ALA	2.6
1	B	785	GLU	2.5
1	A	896	THR	2.5
1	C	914	ASP	2.5
1	C	884	LEU	2.5
1	A	748	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	894	HIS	2.5
1	C	890	SER	2.5
1	C	783	MET	2.5
1	B	709	VAL	2.5
1	B	786	VAL	2.5
1	C	632	TYR	2.4
1	B	628	PHE	2.4
1	D	590	ILE	2.4
1	C	683	ASP	2.4
1	B	824	GLU	2.4
1	C	915	PHE	2.4
1	D	917	ASP	2.4
1	B	916	LEU	2.4
1	C	878	PHE	2.4
1	C	722	GLU	2.3
1	C	903	THR	2.3
1	B	590	ILE	2.3
1	B	887	ARG	2.3
1	C	794	GLN	2.3
1	A	595	ALA	2.3
1	C	634	ILE	2.3
1	C	816	TRP	2.3
1	B	721	SER	2.3
1	D	893	GLU	2.3
1	D	921	GLU	2.3
1	C	799	LEU	2.3
1	C	916	LEU	2.3
1	B	892	ARG	2.2
1	B	717	ALA	2.2
1	A	898	VAL	2.2
1	D	636	CYS	2.2
1	B	720	SER	2.2
1	A	915	PHE	2.2
1	B	915	PHE	2.2
1	C	592	PRO	2.2
1	D	892	ARG	2.2
1	A	851	GLU	2.1
1	B	890	SER	2.1
1	C	679	ASN	2.1
1	C	776	ILE	2.1
1	A	917	ASP	2.1
1	A	598	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	909	SER	2.1
1	C	902	PHE	2.1
1	B	613	ASP	2.1
1	D	634	ILE	2.1
1	A	887	ARG	2.1
1	C	611	PRO	2.1
1	C	879	PRO	2.1
1	A	904	ILE	2.0
1	C	840	MET	2.0
1	A	916	LEU	2.0
1	D	593	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	Q2T	B	1917	32/32	0.90	0.09	14,18,23,24	0
3	Q2T	A	1918	32/32	0.91	0.09	17,18,23,24	0
3	Q2T	C	1917	32/32	0.91	0.09	15,17,23,24	0
3	Q2T	D	1922	32/32	0.93	0.07	15,16,24,26	0
4	MG	C	1918	1/1	0.95	0.04	11,11,11,11	0
4	MG	B	1918	1/1	0.96	0.04	12,12,12,12	0
4	MG	D	1923	1/1	0.96	0.05	11,11,11,11	0
4	MG	A	1919	1/1	0.98	0.03	12,12,12,12	0
2	ZN	B	1001	1/1	0.99	0.04	13,13,13,13	0
2	ZN	D	1001	1/1	0.99	0.04	13,13,13,13	0
2	ZN	C	1001	1/1	1.00	0.03	14,14,14,14	0
2	ZN	A	1001	1/1	1.00	0.03	13,13,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Q2T B 1917:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Q2T A 1918:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Q2T C 1917:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.