



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:11 PM UTC

PDB ID : 6D06 / pdb_00006d06
Title : Human ADAR2d E488Y mutant complexed with dsRNA containing an abasic site opposite the edited base
Authors : Matthews, M.M.; Fisher, A.J.; Beal, P.A.
Deposited on : 2018-04-10
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)
RNA backbone	3983	1112 (2.80-2.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	87% 8% 5%
1	D	403	5% 77% 10% 13%
2	B	23	78% 13% 9%
3	C	23	78% 22%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Double-stranded RNA-specific editase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	0	4	0
			3034	1916	549	558	11			
1	D	352	Total	C	N	O	S	0	0	0
			2754	1746	494	503	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	488	TYR	GLU	engineered mutation	UNP P78563
D	488	TYR	GLU	engineered mutation	UNP P78563


- Molecule 2 is a RNA chain called RNA (5'-R(*GP*CP*UP*CP*GP*CP*GP*AP*UP*GP*CP*UP*(8AZ)P*GP*AP*GP*GP*GP*CP*UP*CP*UP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	23	Total	C	N	O	P	0	0	0
			490	218	88	162	22			

- Molecule 3 is a RNA chain called RNA (5'-R(*CP*AP*GP*AP*GP*CP*CP*CP*CP*CP*NP*AP*GP*CP*AP*UP*CP*GP*CP*GP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	23	Total	C	N	O	P	0	0	0
			477	214	87	154	22			

- Molecule 4 is INOSITOL HEXAKISPHOSPHATE (CCD ID: IHP) (formula: C₆H₁₈O₂₄P₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			36	6	24	6		
4	D	1	Total	C	O	P	0	0
			36	6	24	6		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	D	1	Total	Zn	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	29	Total	O	0	0
			29	29		
6	B	12	Total	O	0	0
			12	12		
6	C	4	Total	O	0	0
			4	4		
6	D	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Double-stranded RNA-specific editase 1




- Molecule 1: Double-stranded RNA-specific editase 1




- Molecule 2: RNA (5'-R(*GP*CP*UP*CP*GP*CP*GP*AP*UP*GP*CP*UP*(8AZ)P*GP*AP*GP*GP*GP*CP*UP*CP*UP*G)-3')



- Molecule 3: RNA (5'-R(*CP*AP*GP*AP*GP*CP*CP*CP*CP*NP*AP*GP*CP*AP*UP*CP*GP*CP*GP*AP*GP*C)-3')





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	174.68Å 63.44Å 132.06Å 90.00° 126.64° 90.00°	Depositor
Resolution (Å)	105.96 – 2.55 105.96 – 2.55	Depositor EDS
% Data completeness (in resolution range)	97.7 (105.96-2.55) 97.7 (105.96-2.55)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.190 , 0.255 0.196 , 0.254	Depositor DCC
R_{free} test set	1862 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtriage
Anisotropy	0.410	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6877	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IHP, ZN, 8AZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	0/3100	1.06	3/4191 (0.1%)
1	D	0.83	0/2809	1.00	1/3794 (0.0%)
2	B	0.47	0/521	1.05	4/809 (0.5%)
3	C	0.55	1/518 (0.2%)	0.97	3/803 (0.4%)
All	All	0.84	1/6948 (0.0%)	1.03	11/9597 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	13	G	O3'-P	-5.27	1.53	1.61

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	481	ARG	NE-CZ-NH2	-7.82	112.16	119.20
2	B	18	G	C3'-C2'-O2'	7.42	121.83	110.70
2	B	10	G	C3'-C2'-O2'	7.04	121.25	110.70
1	A	456	ILE	N-CA-C	6.24	117.73	110.62
2	B	23	G	C2'-C3'-O3'	-6.21	104.38	113.70
2	B	2	C	C1'-C2'-O2'	-6.02	99.37	108.40
1	D	487	GLY	N-CA-C	5.68	123.16	115.32
3	C	6	C	C3'-C2'-O2'	5.36	118.74	110.70
1	A	481	ARG	NE-CZ-NH1	5.28	126.78	121.50
3	C	13	G	C4'-C3'-O3'	-5.26	105.10	113.00
3	C	20	G	C1'-C2'-O2'	5.15	116.12	108.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3034	0	3053	13	0
1	D	2754	0	2788	13	0
2	B	490	0	239	1	0
3	C	477	0	251	0	0
4	A	36	0	6	1	0
4	D	36	0	6	0	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	A	29	0	0	0	0
6	B	12	0	0	0	0
6	C	4	0	0	0	0
6	D	3	0	0	0	0
All	All	6877	0	6343	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:801:IHP:O26	4:A:801:IHP:O21	2.19	0.60
1:D:377:CYS:SG	1:D:483:LYS:HD3	2.43	0.58
1:D:526:VAL:HB	1:D:668:TYR:CE1	2.39	0.57
1:A:400:ARG:HD3	1:A:523:TRP:CE2	2.41	0.55
1:A:503:ASP:OD2	1:D:490:THR:OG1	2.24	0.54
1:D:541:TYR:HB3	1:D:577:ASN:ND2	2.27	0.50
1:D:532:LEU:HB3	1:D:636:TRP:CD1	2.48	0.48
1:A:488[A]:TYR:CD1	1:A:488[A]:TYR:N	2.83	0.47
1:D:400:ARG:HD3	1:D:523:TRP:CZ2	2.50	0.46
1:D:400:ARG:HD3	1:D:523:TRP:CE2	2.50	0.46
1:A:547:LEU:HD11	1:A:556:LEU:HD22	1.99	0.45
1:A:369:ILE:HG21	1:A:403:LEU:HD12	1.98	0.44
1:A:460:HIS:C	1:A:550:LEU:HD22	2.43	0.44
1:D:644:PRO:HD2	1:D:647:LEU:HD12	2.00	0.43
1:D:323:LEU:O	1:D:327:VAL:HG13	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:463:ILE:HD12	1:A:463:ILE:O	2.19	0.43
1:D:380:GLY:HA2	1:D:383:MET:HE3	2.00	0.43
1:D:514:MET:HG3	1:D:687:TRP:CE2	2.53	0.43
1:D:392:ASP:OD2	1:D:483:LYS:NZ	2.48	0.43
1:A:372:SER:HA	1:A:600:VAL:O	2.20	0.42
1:A:600:VAL:HG13	1:A:609:ILE:HB	2.01	0.42
1:A:455:ARG:HD3	2:B:13:8AZ:OP1	2.20	0.41
1:D:537:VAL:HG23	1:D:538:GLU:O	2.20	0.41
1:A:692:THR:HG22	1:A:696:GLN:HE21	1.84	0.41
1:A:373:THR:O	1:A:599:SER:OG	2.32	0.40
1:A:556:LEU:O	1:A:560:MET:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/403 (96%)	368 (95%)	16 (4%)	2 (0%)	24	34
1	D	346/403 (86%)	319 (92%)	23 (7%)	4 (1%)	10	13
All	All	732/806 (91%)	687 (94%)	39 (5%)	6 (1%)	16	22

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	699	LEU
1	A	465	GLU
1	D	462	PRO
1	D	573	LEU
1	A	561	TYR
1	D	652	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/347 (94%)	318 (97%)	10 (3%)	36	53
1	D	300/347 (86%)	283 (94%)	17 (6%)	18	28
All	All	628/694 (90%)	601 (96%)	27 (4%)	26	39

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	328	SER
1	A	340	ASP
1	A	470	ARG
1	A	479	GLN
1	A	488[A]	TYR
1	A	488[B]	TYR
1	A	490	THR
1	A	528	ILE
1	A	652	ILE
1	A	653	THR
1	D	327	VAL
1	D	384	SER
1	D	399	SER
1	D	419	ASP
1	D	461	GLU
1	D	463	ILE
1	D	484	ILE
1	D	493	VAL
1	D	537	VAL
1	D	562	GLN
1	D	565	SER
1	D	568	GLU
1	D	569	ASP
1	D	580	LEU
1	D	651	LYS
1	D	661	SER
1	D	700	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	416	ASN
1	A	417	ASN
1	A	427	GLN
1	A	479	GLN
1	A	696	GLN
1	D	341	ASN
1	D	421	GLN
1	D	427	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	22/23 (95%)	1 (4%)	0
3	C	20/23 (86%)	2 (10%)	0
All	All	42/46 (91%)	3 (7%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	18	G
3	C	7	C
3	C	14	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	8AZ	B	13	5,2	18,24,25	0.53	0	19,35,38	1.02	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8AZ	B	13	5,2	-	0/7/35/36	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	13	8AZ	C1'-N9-N8	2.58	124.20	119.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	13	8AZ	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	IHP	A	801	-	36,36,36	0.99	1 (2%)	60,60,60	1.28	6 (10%)
4	IHP	D	801	-	36,36,36	0.90	1 (2%)	60,60,60	1.20	7 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IHP	A	801	-	-	0/30/54/54	0/1/1/1
4	IHP	D	801	-	-	2/30/54/54	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	801	IHP	P4-O14	2.40	1.63	1.59
4	A	801	IHP	P1-O11	2.39	1.63	1.59

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	801	IHP	O13-C3-C4	3.65	116.52	108.76
4	D	801	IHP	O46-P6-O16	-3.05	93.97	105.85
4	D	801	IHP	O46-P6-O26	2.72	121.43	110.83
4	A	801	IHP	P5-O15-C5	2.53	130.20	123.43
4	A	801	IHP	O11-C1-C6	2.36	113.79	108.76
4	D	801	IHP	O41-P1-O31	2.35	116.61	107.80
4	D	801	IHP	O42-P2-O12	-2.32	96.81	105.85
4	D	801	IHP	O44-P4-O34	2.25	116.25	107.80
4	A	801	IHP	O16-C6-C5	2.24	113.52	108.76
4	A	801	IHP	O41-P1-O21	2.23	119.51	110.83
4	D	801	IHP	O46-P6-O36	2.15	115.87	107.80
4	A	801	IHP	O11-P1-O21	-2.11	101.80	109.33
4	D	801	IHP	C6-C5-C4	2.11	115.05	110.43

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	801	IHP	C2-O12-P2-O32
4	D	801	IHP	C1-O11-P1-O21


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	801	IHP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand IHP A 801




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	384/403 (95%)	-0.15	5 (1%) 75 75	20, 51, 86, 126	4 (1%)
1	D	352/403 (87%)	0.45	19 (5%) 31 31	50, 76, 111, 130	0
2	B	22/23 (95%)	-0.83	0 100 100	45, 60, 81, 90	0
3	C	22/23 (95%)	-0.56	0 100 100	44, 69, 83, 120	0
All	All	780/852 (91%)	0.09	24 (3%) 51 52	20, 63, 103, 130	4 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	363	VAL	5.4
1	D	383	MET	3.3
1	D	700	THR	3.3
1	A	464	LEU	3.1
1	D	512	LEU	3.0
1	D	389	ALA	2.9
1	D	699	LEU	2.8
1	A	463	ILE	2.7
1	D	494	ARG	2.7
1	D	319	LEU	2.7
1	D	488	TYR	2.7
1	A	700	THR	2.6
1	D	463	ILE	2.6
1	D	647	LEU	2.6
1	A	497	ALA	2.3
1	D	653	THR	2.3
1	D	642	LYS	2.3
1	D	654	LYS	2.2
1	A	467	PRO	2.2
1	D	384	SER	2.2
1	D	361	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	418	LYS	2.1
1	D	485	GLU	2.1
1	D	658	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8AZ	B	13	22/23	0.97	0.06	37,41,50,54	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	IHP	D	801	36/36	0.95	0.07	52,73,87,89	0
4	IHP	A	801	36/36	0.98	0.04	34,41,47,48	0
5	ZN	A	802	1/1	1.00	0.02	42,42,42,42	0
5	ZN	D	802	1/1	1.00	0.01	58,58,58,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IHP D 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IHP A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.