



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:52 PM UTC

PDB ID : 1D1I / pdb_00001d1i
Title : MUTATED SHIGA-LIKE TOXIN B SUBUNIT (W34A) COMPLEXED
WITH RECEPTOR GB3 ANALOGUE
Authors : Ling, H.; Boodhoo, A.; Brunton, J.L.; Read, R.J.
Deposited on : 1999-09-17
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	69	87% 12% .
1	B	69	% 91% 9%
1	C	69	% 88% 12%
1	D	69	90% 10%
1	E	69	% 91% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 50%50%
3	G	3	 100%
3	H	3	 100%
3	I	3	 100%
3	J	3	 100%
3	K	3	 100%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SHIGA TOXIN B-CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	69	Total	C	N	O	S	0	0	0
			531	331	89	108	3			
1	B	69	Total	C	N	O	S	0	0	0
			531	331	89	108	3			
1	C	69	Total	C	N	O	S	0	0	0
			531	331	89	108	3			
1	D	69	Total	C	N	O	S	0	0	0
			531	331	89	108	3			
1	E	69	Total	C	N	O	S	0	0	0
			531	331	89	108	3			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	ALA	TRP	engineered mutation	UNP P08027
B	234	ALA	TRP	engineered mutation	UNP P08027
C	334	ALA	TRP	engineered mutation	UNP P08027
D	434	ALA	TRP	engineered mutation	UNP P08027
E	534	ALA	TRP	engineered mutation	UNP P08027


- Molecule 2 is an oligosaccharide called alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	F	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is an oligosaccharide called alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose.

se-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	G	3	Total	C	O	0	0	0
			34	18	16			
3	H	3	Total	C	O	0	0	0
			34	18	16			
3	I	3	Total	C	O	0	0	0
			34	18	16			
3	J	3	Total	C	O	0	0	0
			34	18	16			
3	K	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	35	Total	O	0	0
			35	35		
4	B	40	Total	O	0	0
			40	40		
4	C	41	Total	O	0	0
			41	41		
4	D	41	Total	O	0	0
			41	41		
4	E	39	Total	O	0	0
			39	39		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SHIGA TOXIN B-CHAIN

Chain A:  87% 12%



- Molecule 1: SHIGA TOXIN B-CHAIN

Chain B:  91% 9%



- Molecule 1: SHIGA TOXIN B-CHAIN

Chain C:  88% 12%




- Molecule 1: SHIGA TOXIN B-CHAIN

Chain D:  90% 10%



- Molecule 1: SHIGA TOXIN B-CHAIN

Chain E:  91% 9%



- Molecule 2: alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose

Chain F:  50% 50%



- Molecule 3: alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranos e

Chain G:  100%



- Molecule 3: alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranos e

Chain H:  100%



- Molecule 3: alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranos e

Chain I:  100%



- Molecule 3: alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranos e

Chain J:  100%



- Molecule 3: alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranos e

Chain K:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	44.23Å 44.14Å 53.88Å 106.04° 106.37° 99.22°	Depositor
Resolution (Å)	19.70 – 1.70 19.70 – 1.70	Depositor EDS
% Data completeness (in resolution range)	91.1 (19.70-1.70) 91.0 (19.70-1.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 1.70Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.191 , 0.211 0.170 , 0.192	Depositor DCC
R_{free} test set	1101 reflections (3.03%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.147	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3044	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLA, BGC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	1/538 (0.2%)	1.82	8/726 (1.1%)
1	B	1.07	1/538 (0.2%)	1.73	8/726 (1.1%)
1	C	1.13	1/538 (0.2%)	1.71	8/726 (1.1%)
1	D	1.09	1/538 (0.2%)	1.67	7/726 (1.0%)
1	E	1.15	1/538 (0.2%)	1.72	8/726 (1.1%)
All	All	1.12	5/2690 (0.2%)	1.73	39/3630 (1.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	358	HIS	CD2-NE2	-8.04	1.29	1.37
1	A	158	HIS	CD2-NE2	-7.86	1.29	1.37
1	B	258	HIS	CD2-NE2	-6.83	1.30	1.37
1	E	558	HIS	CD2-NE2	-6.38	1.30	1.37
1	D	458	HIS	CD2-NE2	-6.15	1.31	1.37

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	ARG	CD-NE-CZ	21.86	155.01	124.40
1	B	269	ARG	CD-NE-CZ	21.68	154.75	124.40
1	D	469	ARG	CD-NE-CZ	16.28	147.20	124.40
1	E	569	ARG	CD-NE-CZ	16.20	147.09	124.40
1	C	369	ARG	CD-NE-CZ	16.03	146.84	124.40
1	B	269	ARG	NE-CZ-NH2	-13.38	107.16	119.20
1	A	169	ARG	NE-CZ-NH2	-12.26	108.17	119.20
1	D	469	ARG	NE-CZ-NH2	10.52	128.66	119.20
1	D	469	ARG	NE-CZ-NH1	-10.51	110.99	121.50
1	E	569	ARG	NE-CZ-NH1	-10.42	111.08	121.50
1	C	369	ARG	NE-CZ-NH1	-10.11	111.39	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	569	ARG	NE-CZ-NH2	9.81	128.03	119.20
1	B	269	ARG	NE-CZ-NH1	9.78	131.28	121.50
1	A	169	ARG	NE-CZ-NH1	9.25	130.75	121.50
1	C	369	ARG	NE-CZ-NH2	9.25	127.53	119.20
1	A	158	HIS	CB-CG-CD2	-7.23	121.80	131.20
1	C	358	HIS	CB-CG-CD2	-7.14	121.92	131.20
1	C	344	GLN	OE1-CD-NE2	-7.10	115.50	122.60
1	B	258	HIS	CB-CG-CD2	-7.02	122.07	131.20
1	D	458	HIS	CB-CG-CD2	-6.75	122.43	131.20
1	E	558	HIS	CB-CG-CD2	-6.59	122.63	131.20
1	C	355	ASN	OD1-CG-ND2	-6.38	116.22	122.60
1	C	358	HIS	CB-CG-ND1	6.30	132.16	122.70
1	E	532	ASN	OD1-CG-ND2	-6.28	116.32	122.60
1	E	555	ASN	OD1-CG-ND2	-6.26	116.33	122.60
1	A	158	HIS	CB-CG-ND1	6.10	131.85	122.70
1	E	508	LYS	CA-CB-CG	-6.07	101.96	114.10
1	B	258	HIS	CB-CG-ND1	6.04	131.76	122.70
1	E	558	HIS	CB-CG-ND1	5.98	131.66	122.70
1	D	408	LYS	CA-CB-CG	-5.89	102.33	114.10
1	A	108	LYS	CA-CB-CG	-5.81	102.48	114.10
1	B	208	LYS	CA-CB-CG	-5.73	102.64	114.10
1	A	115	ASN	OD1-CG-ND2	-5.66	116.94	122.60
1	B	244	GLN	OE1-CD-NE2	-5.66	116.94	122.60
1	D	458	HIS	CB-CG-ND1	5.63	131.15	122.70
1	B	259	ASN	CA-CB-CG	-5.43	107.17	112.60
1	C	308	LYS	CA-CB-CG	-5.42	103.25	114.10
1	A	159	ASN	CA-CB-CG	-5.15	107.45	112.60
1	D	432	ASN	OD1-CG-ND2	-5.02	117.58	122.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	531	0	519	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	531	0	519	0	0
1	C	531	0	519	1	0
1	D	531	0	519	1	0
1	E	531	0	519	0	0
2	F	23	0	21	1	0
3	G	34	0	30	0	0
3	H	34	0	30	0	0
3	I	34	0	30	0	0
3	J	34	0	30	0	0
3	K	34	0	30	0	0
4	A	35	0	0	0	0
4	B	40	0	0	0	0
4	C	41	0	0	0	0
4	D	41	0	0	0	0
4	E	39	0	0	0	0
All	All	3044	0	2766	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:VAL:HG21	1:A:144:GLN:HB2	1.90	0.53
1:C:309:VAL:HG13	1:C:322:VAL:HG23	1.98	0.46
1:D:409:VAL:HG13	1:D:422:VAL:HG23	1.99	0.45
1:A:160:GLY:HA2	2:F:2:GLA:O6	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/69 (97%)	66 (98%)	1 (2%)	0	100	100
1	B	67/69 (97%)	66 (98%)	1 (2%)	0	100	100
1	C	67/69 (97%)	66 (98%)	1 (2%)	0	100	100
1	D	67/69 (97%)	65 (97%)	2 (3%)	0	100	100
1	E	67/69 (97%)	66 (98%)	1 (2%)	0	100	100
All	All	335/345 (97%)	329 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/60 (100%)	58 (97%)	2 (3%)	33	17
1	B	60/60 (100%)	59 (98%)	1 (2%)	53	38
1	C	60/60 (100%)	59 (98%)	1 (2%)	53	38
1	D	60/60 (100%)	59 (98%)	1 (2%)	53	38
1	E	60/60 (100%)	59 (98%)	1 (2%)	53	38
All	All	300/300 (100%)	294 (98%)	6 (2%)	48	32

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	VAL
1	A	129	LEU
1	B	229	LEU
1	C	329	LEU
1	D	429	LEU
1	E	529	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN
1	A	144	GLN
1	A	158	HIS
1	B	237	GLN
1	B	244	GLN
1	B	255	ASN
1	C	337	GLN
1	C	344	GLN
1	C	358	HIS
1	D	437	GLN
1	D	444	GLN
1	E	544	GLN
1	E	558	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GAL	F	1	2	12,12,12	0.44	0	17,17,17	0.41	0
2	GLA	F	2	2	11,11,12	0.54	0	15,15,17	0.65	0
3	BGC	G	1	3	12,12,12	0.38	0	17,17,17	0.58	0
3	GAL	G	2	3	11,11,12	0.49	0	15,15,17	0.37	0
3	GLA	G	3	3	11,11,12	0.42	0	15,15,17	0.60	0
3	BGC	H	1	3	12,12,12	0.35	0	17,17,17	0.43	0
3	GAL	H	2	3	11,11,12	0.45	0	15,15,17	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLA	H	3	3	11,11,12	0.45	0	15,15,17	0.57	0
3	BGC	I	1	3	12,12,12	0.49	0	17,17,17	0.55	0
3	GAL	I	2	3	11,11,12	0.56	0	15,15,17	0.36	0
3	GLA	I	3	3	11,11,12	0.34	0	15,15,17	0.60	0
3	BGC	J	1	3	12,12,12	0.29	0	17,17,17	0.42	0
3	GAL	J	2	3	11,11,12	0.54	0	15,15,17	0.41	0
3	GLA	J	3	3	11,11,12	0.46	0	15,15,17	0.66	0
3	BGC	K	1	3	12,12,12	0.48	0	17,17,17	0.59	0
3	GAL	K	2	3	11,11,12	0.58	0	15,15,17	0.53	0
3	GLA	K	3	3	11,11,12	0.43	0	15,15,17	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	F	1	2	-	0/2/22/22	0/1/1/1
2	GLA	F	2	2	-	1/2/19/22	0/1/1/1
3	BGC	G	1	3	-	0/2/22/22	0/1/1/1
3	GAL	G	2	3	-	0/2/19/22	0/1/1/1
3	GLA	G	3	3	-	0/2/19/22	0/1/1/1
3	BGC	H	1	3	-	0/2/22/22	0/1/1/1
3	GAL	H	2	3	-	0/2/19/22	0/1/1/1
3	GLA	H	3	3	-	0/2/19/22	0/1/1/1
3	BGC	I	1	3	-	0/2/22/22	0/1/1/1
3	GAL	I	2	3	-	0/2/19/22	0/1/1/1
3	GLA	I	3	3	-	0/2/19/22	0/1/1/1
3	BGC	J	1	3	-	0/2/22/22	0/1/1/1
3	GAL	J	2	3	-	0/2/19/22	0/1/1/1
3	GLA	J	3	3	-	0/2/19/22	0/1/1/1
3	BGC	K	1	3	-	0/2/22/22	0/1/1/1
3	GAL	K	2	3	-	0/2/19/22	0/1/1/1
3	GLA	K	3	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	2	GLA	O5-C5-C6-O6



There are no ring outliers.



1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	2	GLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	69/69 (100%)	-0.34	0 100 100	10, 16, 25, 36	0
1	B	69/69 (100%)	-0.30	1 (1%) 73 77	10, 17, 26, 37	0
1	C	69/69 (100%)	-0.43	1 (1%) 73 77	11, 15, 24, 38	0
1	D	69/69 (100%)	-0.41	0 100 100	10, 15, 24, 36	0
1	E	69/69 (100%)	-0.33	1 (1%) 73 77	10, 16, 26, 36	0
All	All	345/345 (100%)	-0.36	3 (0%) 81 83	10, 16, 25, 38	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	369	ARG	3.0
1	B	269	ARG	2.5
1	E	569	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLA	F	2	11/12	0.26	0.28	73,75,76,77	0
2	GAL	F	1	12/12	0.48	0.21	77,78,79,79	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BGC	K	1	12/12	0.80	0.12	28,34,37,38	0
3	BGC	I	1	12/12	0.87	0.10	24,31,33,35	0
3	BGC	G	1	12/12	0.89	0.09	26,35,38,39	0
3	BGC	J	1	12/12	0.91	0.08	12,20,23,23	0
3	BGC	H	1	12/12	0.92	0.08	16,21,25,28	0
3	GAL	K	2	11/12	0.92	0.09	19,23,25,26	0
3	GAL	G	2	11/12	0.96	0.06	18,21,23,25	0
3	GLA	G	3	11/12	0.96	0.06	16,17,18,21	0
3	GAL	I	2	11/12	0.96	0.05	17,19,21,22	0
3	GLA	K	3	11/12	0.96	0.05	14,17,19,19	0
3	GLA	J	3	11/12	0.97	0.05	14,16,17,17	0
3	GLA	I	3	11/12	0.97	0.05	13,15,16,17	0
3	GLA	H	3	11/12	0.98	0.04	13,16,18,19	0
3	GAL	J	2	11/12	0.98	0.04	13,14,15,16	0
3	GAL	H	2	11/12	0.98	0.04	13,15,16,17	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.