



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 05:33 AM UTC

PDB ID : 1D1U / pdb_00001d1u
Title : USE OF AN N-TERMINAL FRAGMENT FROM MOLONEY MURINE LEUKEMIA VIRUS REVERSE TRANSCRIPTASE TO FACILITATE CRYSTALLIZATION AND ANALYSIS OF A PSEUDO-16-MER DNA MOLECULE CONTAINING G-A MISPAIRS
Authors : Cote, M.L.; Yohannan, S.; Georgiadis, M.M.
Deposited on : 1999-09-21
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	6	17% 33% 50% 17%
2	C	10	30% 30% 70%
3	A	255	9% 65% 30% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*TP*CP*GP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	6	Total	C	N	O	P	0	0	0
			119	58	20	36	5			

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*CP*GP*GP*CP*AP*CP*GP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			205	97	44	55	9			

- Molecule 3 is a protein called PROTEIN (REVERSE TRANSCRIPTASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	255	Total	C	N	O	S	0	1	0
			2053	1320	357	369	7			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	7	Total	O	0	0
			7	7		
4	C	6	Total	O	0	0
			6	6		
4	A	162	Total	O	0	0
			162	162		

- Molecule 1: DNA (5'-D(*CP*TP*CP*GP*TP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	54.74Å 145.49Å 46.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.00-2.30) 99.0 (50.00-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.89 (at 2.31Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.230 , 0.286 0.229 , 0.286	Depositor DCC
R_{free} test set	861 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2552	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.27	0/132	0.84	0/202
2	C	1.04	1/231 (0.4%)	0.46	0/355
3	A	0.40	0/2110	0.93	7/2877 (0.2%)
All	All	0.49	1/2473 (0.0%)	0.89	7/3434 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	7	DA	O3'-P	15.39	1.84	1.61

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	236	CYS	N-CA-C	-6.23	104.74	112.90
3	A	139	LEU	CA-C-N	6.20	124.12	119.66
3	A	139	LEU	C-N-CA	6.20	124.12	119.66
3	A	43	VAL	N-CA-C	5.70	118.17	111.05
3	A	264	LYS	N-CA-C	-5.49	106.25	113.17
3	A	91	GLN	N-CA-C	-5.30	99.65	108.34
3	A	102	LYS	N-CA-C	5.11	114.98	108.24

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	4	DG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	119	0	70	10	0
2	C	205	0	112	6	0
3	A	2053	0	2065	73	0
4	A	162	0	0	2	0
4	B	7	0	0	0	0
4	C	6	0	0	1	1
All	All	2552	0	2247	83	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:179:ILE:HD13	3:A:183:LEU:HD21	1.29	1.10
2:C:12:DA:H2''	2:C:13:DC:H5'	1.32	1.08
1:B:3:DC:H1'	3:A:116:ARG:HH12	1.40	0.87
3:A:61:ILE:HD11	3:A:117:GLU:HG3	1.63	0.81
3:A:206:ASP:HB3	3:A:250:LEU:HD13	1.63	0.80
3:A:74:ILE:HG23	3:A:111:PRO:HG3	1.62	0.79
3:A:102:LYS:N	3:A:102:LYS:HD3	2.01	0.76
1:B:3:DC:H1'	3:A:116:ARG:NH1	2.00	0.76
1:B:3:DC:H2'	1:B:4:DG:C8	2.21	0.76
3:A:218:ILE:HB	3:A:229:ALA:HB3	1.67	0.75
3:A:161:HIS:HD2	3:A:163:THR:H	1.36	0.73
1:B:1:DC:O5'	3:A:64[B]:TYR:CZ	2.41	0.71
3:A:59:VAL:HG11	3:A:121:ARG:NH2	2.05	0.71
3:A:237:GLN:O	3:A:241:ARG:HG3	1.93	0.67
3:A:59:VAL:HG11	3:A:121:ARG:CZ	2.24	0.67
2:C:12:DA:C2'	2:C:13:DC:H5'	2.20	0.62
3:A:75:LYS:HB3	3:A:76:PRO:HD3	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:31:GLN:NE2	3:A:249:ASN:HD22	1.98	0.62
3:A:161:HIS:CD2	3:A:163:THR:H	2.17	0.61
1:B:2:DT:H5'	3:A:64[A]:TYR:CE1	2.38	0.59
3:A:68:GLN:HG2	3:A:72:LEU:HD22	1.85	0.58
3:A:122:VAL:HG22	3:A:193:LYS:HE2	1.86	0.57
3:A:277:GLN:HA	3:A:277:GLN:HE21	1.72	0.55
3:A:216:ASP:OD2	3:A:216:ASP:N	2.40	0.55
2:C:8:DC:H2'	2:C:9:DG:C8	2.42	0.54
3:A:31:GLN:HE22	3:A:249:ASN:HD22	1.54	0.54
1:B:2:DT:H5'	3:A:64[A]:TYR:HE1	1.73	0.54
3:A:220:LEU:HD22	3:A:227:LEU:HD23	1.88	0.54
3:A:230:ALA:HB3	3:A:236:CYS:HB2	1.90	0.54
3:A:233:GLU:HG2	3:A:237:GLN:HE21	1.73	0.53
3:A:277:GLN:HA	3:A:277:GLN:NE2	2.23	0.53
3:A:31:GLN:HE22	3:A:249:ASN:ND2	2.06	0.53
3:A:77:HIS:O	3:A:81:LEU:HD22	2.08	0.53
3:A:161:HIS:CD2	3:A:163:THR:HG22	2.44	0.52
1:B:1:DC:H2''	1:B:2:DT:H6	1.74	0.52
3:A:100:PRO:HG2	4:A:383:HOH:O	2.10	0.52
3:A:31:GLN:NE2	3:A:249:ASN:ND2	2.57	0.51
3:A:213:GLN:NE2	3:A:213:GLN:HA	2.26	0.51
3:A:61:ILE:HD11	3:A:117:GLU:CG	2.38	0.49
3:A:62:LYS:HA	3:A:62:LYS:HE2	1.93	0.49
3:A:158:LEU:HD11	3:A:199:PHE:HA	1.95	0.48
3:A:41:LEU:C	3:A:41:LEU:HD23	2.39	0.48
3:A:213:GLN:HA	3:A:213:GLN:HE21	1.78	0.48
3:A:131:ASN:HD21	3:A:134:ASN:ND2	2.10	0.48
2:C:9:DG:H1'	2:C:10:DG:C8	2.50	0.47
2:C:13:DC:H4'	4:C:21:HOH:O	2.13	0.47
3:A:41:LEU:O	3:A:251:GLY:HA3	2.15	0.46
3:A:77:HIS:CD2	3:A:80:ARG:NH2	2.83	0.46
3:A:70:ALA:HB1	3:A:100:PRO:CB	2.45	0.46
2:C:13:DC:H2''	2:C:14:DG:OP2	2.16	0.45
3:A:145:TRP:CH2	3:A:233:GLU:HB2	2.52	0.45
3:A:24:THR:HB	3:A:27:SER:OG	2.17	0.45
3:A:141:PRO:HD2	4:A:421:HOH:O	2.15	0.44
3:A:66:MET:HE2	3:A:100:PRO:HG3	1.99	0.44
3:A:102:LYS:N	3:A:102:LYS:CD	2.74	0.44
3:A:26:LEU:C	3:A:26:LEU:HD23	2.43	0.44
3:A:26:LEU:HD23	3:A:26:LEU:O	2.17	0.44
3:A:26:LEU:HA	3:A:33:TRP:NE1	2.33	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:DC:H2''	1:B:2:DT:C6	2.52	0.43
1:B:2:DT:C2'	1:B:3:DC:O5'	2.66	0.43
3:A:214:HIS:N	3:A:215:PRO:HD3	2.33	0.43
3:A:70:ALA:HB1	3:A:100:PRO:HB3	2.00	0.42
3:A:29:PHE:O	3:A:33:TRP:CD1	2.72	0.42
3:A:179:ILE:HD13	3:A:183:LEU:CD2	2.22	0.42
1:B:1:DC:O5'	3:A:64[B]:TYR:CE2	2.65	0.42
3:A:68:GLN:CG	3:A:72:LEU:HD22	2.48	0.42
3:A:158:LEU:HD12	3:A:188:LEU:HD22	2.02	0.41
3:A:39:MET:HE3	3:A:86:ILE:HD11	2.02	0.41
3:A:264:LYS:N	3:A:264:LYS:HD3	2.35	0.41
3:A:98:LEU:HD12	3:A:98:LEU:HA	1.89	0.41
3:A:99:LEU:HD12	3:A:99:LEU:N	2.35	0.41
3:A:49:ILE:O	3:A:51:PRO:HD3	2.20	0.41
3:A:98:LEU:C	3:A:99:LEU:HD12	2.45	0.41
3:A:99:LEU:HA	3:A:100:PRO:HD3	1.80	0.41
3:A:77:HIS:HA	3:A:80:ARG:NH2	2.35	0.41
3:A:197:THR:O	3:A:201:GLU:HG3	2.21	0.41
3:A:41:LEU:HD23	3:A:42:ALA:N	2.36	0.41
3:A:39:MET:HE3	3:A:86:ILE:CD1	2.51	0.40
3:A:103:LYS:HE2	3:A:108:ASP:OD1	2.22	0.40
3:A:101:VAL:C	3:A:102:LYS:HD3	2.47	0.40
3:A:274:LYS:NZ	3:A:275:GLU:OE1	2.53	0.40
3:A:213:GLN:C	3:A:215:PRO:HD3	2.47	0.40
3:A:221:GLN:HG2	3:A:226:LEU:HD22	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:20:HOH:O	4:C:20:HOH:O[2_455]	1.99	0.21

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	254/255 (100%)	238 (94%)	7 (3%)	9 (4%)	3 1

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	108	ASP
3	A	176	GLU
3	A	181	GLY
3	A	223	VAL
3	A	262	CYS
3	A	104	PRO
3	A	175	PRO
3	A	100	PRO
3	A	178	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	225/224 (100%)	210 (93%)	15 (7%)	15 21

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	57	THR
3	A	62	LYS
3	A	64[A]	TYR
3	A	64[B]	TYR
3	A	72	LEU
3	A	81	LEU
3	A	102	LYS
3	A	122	VAL
3	A	144	GLN
3	A	147	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	175	PRO
3	A	176	GLU
3	A	220	LEU
3	A	244	LEU
3	A	264	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	31	GLN
3	A	77	HIS
3	A	79	GLN
3	A	84	GLN
3	A	134	ASN
3	A	144	GLN
3	A	161	HIS
3	A	213	GLN
3	A	214	HIS
3	A	237	GLN
3	A	245	GLN
3	A	249	ASN
3	A	277	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	7:DA	O3'	8:DC	P	1.84

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	6/6 (100%)	1.26	1 (16%) 4 5	46, 51, 65, 72	0
2	C	10/10 (100%)	1.53	3 (30%) 1 1	35, 73, 85, 87	0
3	A	255/255 (100%)	0.37	23 (9%) 15 16	15, 32, 75, 100	1 (0%)
All	All	271/271 (100%)	0.43	27 (9%) 12 14	15, 32, 79, 100	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	106	THR	6.3
3	A	104	PRO	5.5
3	A	107	ASN	5.3
3	A	99	LEU	4.9
3	A	179	ILE	4.6
3	A	177	MET	4.2
3	A	175	PRO	4.2
3	A	64[A]	TYR	4.1
3	A	108	ASP	3.8
3	A	278	ARG	3.5
3	A	100	PRO	3.4
3	A	101	VAL	3.4
3	A	110	ARG	3.3
3	A	102	LYS	3.2
3	A	105	GLY	3.0
3	A	103	LYS	2.9
3	A	176	GLU	2.7
2	C	8	DC	2.7
3	A	216	ASP	2.5
3	A	234	LEU	2.5
2	C	7	DA	2.4
2	C	12	DA	2.3
3	A	26	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	25	TRP	2.2
1	B	6	DG	2.1
3	A	109	TYR	2.0
3	A	180	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.