



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:32 AM UTC

PDB ID : 2D1Z / pdb_00002d1z
Title : Crystal structure of catalytic-site mutant xylanase from *Streptomyces olivaceoviridis* E-86
Authors : Suzuki, R.; Kuno, A.; Fujimoto, Z.; Ito, S.; Kawahara, S.I.; Kaneko, S.; Hasegawa, T.; Taira, K.
Deposited on : 2005-09-02
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	9% 84% 13% .
1	B	436	18% 82% 15% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDO-1,4-BETA-D-XYLANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	0	0
			3233	1988	588	641	16			
1	B	427	Total	C	N	O	S	0	0	0
			3233	1988	588	641	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	SER	ASN	engineered mutation	UNP Q7SI98
A	128	HIS	GLU	engineered mutation	UNP Q7SI98
B	627	SER	ASN	engineered mutation	UNP Q7SI98
B	628	HIS	GLU	engineered mutation	UNP Q7SI98

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	427	Total 427	O 427	0	0
4	B	474	Total 474	O 474	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ENDO-1,4-BETA-D-XYLANASE



• Molecule 1: ENDO-1,4-BETA-D-XYLANASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.51Å 94.06Å 139.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.03 – 1.60 47.03 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.03-1.60) 99.4 (47.03-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.01 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.191 , 0.207 0.191 , 0.207	Depositor DCC
R_{free} test set	6904 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7438	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/3298	0.91	13/4473 (0.3%)
1	B	0.32	0/3298	0.90	11/4473 (0.2%)
All	All	0.32	0/6596	0.90	24/8946 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	THR	N-CA-C	8.72	124.32	112.90
1	B	735	THR	N-CA-C	8.69	124.00	113.41
1	B	537	GLU	N-CA-C	6.81	121.67	113.23
1	B	546	GLU	N-CA-C	6.68	120.53	112.38
1	A	366	LEU	N-CA-C	-6.62	100.24	110.10
1	B	549	ILE	N-CA-C	6.57	116.73	110.42
1	B	866	LEU	N-CA-C	-6.54	100.36	110.10
1	B	583	LEU	N-CA-C	6.44	119.62	111.69
1	A	37	GLU	N-CA-C	6.27	121.01	113.23
1	A	83	LEU	N-CA-C	6.09	119.19	111.69
1	A	46	GLU	N-CA-C	6.04	120.12	112.87
1	A	49	ILE	N-CA-C	5.96	116.14	110.42
1	B	624	ASP	N-CA-C	-5.74	98.36	108.23
1	B	780	PRO	N-CA-C	5.71	122.91	114.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	LEU	N-CA-C	5.71	118.61	110.10
1	A	169	TYR	N-CA-C	-5.71	99.94	109.24
1	B	737	LEU	N-CA-C	5.70	118.60	110.10
1	A	407	LEU	N-CA-C	-5.68	101.44	109.96
1	B	669	TYR	N-CA-C	-5.62	100.03	109.07
1	A	280	PRO	N-CA-C	5.40	122.47	114.80
1	A	234	ILE	N-CA-C	-5.29	99.48	107.73
1	B	734	ILE	N-CA-C	-5.21	99.59	107.73
1	A	124	ASP	N-CA-C	-5.18	99.31	108.23
1	A	86	HIS	N-CA-C	5.10	117.73	111.82

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	669	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3233	0	3043	32	0
1	B	3233	0	3040	56	0
2	A	5	0	0	0	0
3	A	30	0	40	5	0
3	B	36	0	48	3	0
4	A	427	0	0	5	0
4	B	474	0	0	6	0
All	All	7438	0	6171	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:540:MET:HE3	1:B:541:VAL:N	1.90	0.87
1:B:540:MET:CE	1:B:579:ARG:HG3	2.07	0.84
1:B:911:GLY:H	3:B:1946:GOL:H31	1.41	0.83
1:A:336:GLN:HB2	1:A:376:LYS:HE3	1.63	0.80
1:B:540:MET:HE2	1:B:579:ARG:HG3	1.67	0.76
1:A:139:ARG:HB2	3:A:1442:GOL:H11	1.66	0.76
1:B:816:LYS:HE2	1:B:936:THR:HG21	1.68	0.76
1:A:421:GLN:HE22	3:A:1445:GOL:H32	1.53	0.73
1:B:816:LYS:HG2	1:B:823:CYS:SG	2.36	0.65
1:B:910:VAL:HA	1:B:921:GLN:HE21	1.63	0.63
1:B:935:ARG:HA	1:B:936:THR:HB	1.79	0.63
1:A:238:ASP:HB2	1:A:280:PRO:HB2	1.81	0.63
1:B:549:ILE:HD12	4:B:2168:HOH:O	2.00	0.62
1:A:421:GLN:NE2	3:A:1445:GOL:H32	2.18	0.59
1:A:97:SER:O	1:A:100:THR:HG22	2.03	0.58
1:B:789:LYS:HD2	1:B:891:ARG:HH12	1.68	0.58
1:B:837:VAL:HG23	1:B:920:ILE:HB	1.85	0.57
1:A:127:SER:HA	1:A:170:ASN:O	2.05	0.57
1:A:1:ALA:HB1	4:A:1545:HOH:O	2.06	0.56
1:A:389:LYS:HE3	4:A:1585:HOH:O	2.05	0.56
1:B:778:ASP:O	1:B:779:THR:C	2.48	0.56
1:B:935:ARG:HA	1:B:936:THR:C	2.31	0.56
1:B:936:THR:HG22	1:B:936:THR:OXT	2.05	0.56
1:B:738:ASP:HB2	1:B:780:PRO:HB2	1.87	0.56
1:B:540:MET:HE3	1:B:541:VAL:H	1.67	0.55
1:A:411:GLY:H	3:A:1445:GOL:H31	1.71	0.54
1:A:278:ASP:O	1:A:279:THR:C	2.50	0.54
1:B:754:CYS:SG	1:B:763:ILE:HD11	2.48	0.54
1:B:540:MET:HE1	1:B:579:ARG:HG3	1.87	0.54
1:B:899:VAL:HG22	1:B:906:CYS:SG	2.47	0.54
1:B:676:ASN:HB3	1:B:679:TRP:CD2	2.43	0.54
1:B:720:THR:HG23	4:B:2298:HOH:O	2.09	0.53
1:B:789:LYS:HD2	1:B:891:ARG:NH1	2.24	0.52
1:A:234:ILE:HD12	1:A:263:ILE:HG12	1.92	0.50
1:B:779:THR:N	1:B:780:PRO:HD3	2.27	0.50
1:A:55:GLN:HB2	1:A:58:GLN:HB2	1.92	0.50
1:A:337:VAL:HG23	1:A:420:ILE:HB	1.94	0.50
1:B:925:CYS:HA	1:B:931:GLN:OE1	2.11	0.50
1:A:15:TYR:CG	1:A:40:MET:HG3	2.47	0.50
1:A:142:ASN:O	1:A:146:THR:HG23	2.11	0.50
1:A:279:THR:N	1:A:280:PRO:HD3	2.27	0.50
1:B:540:MET:HE3	1:B:540:MET:C	2.34	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:627:SER:HA	1:B:670:ASN:O	2.12	0.49
1:B:820:SER:HA	1:B:914:THR:HB	1.93	0.49
1:B:822:ARG:HG2	1:B:822:ARG:HH11	1.77	0.49
1:B:874:GLY:HA2	1:B:921:GLN:OE1	2.13	0.49
1:A:85:TRP:CZ3	1:A:87:SER:HB3	2.48	0.48
1:B:935:ARG:CA	1:B:936:THR:HB	2.43	0.48
1:B:910:VAL:HG22	1:B:921:GLN:NE2	2.28	0.48
1:A:190:ARG:O	1:A:194:GLN:HG3	2.14	0.48
1:A:80:GLY:HA3	1:A:123:TRP:CE3	2.48	0.48
1:B:909:ALA:HB3	1:B:930:ASN:HB2	1.96	0.48
1:B:911:GLY:H	3:B:1946:GOL:C3	2.17	0.48
1:B:694:GLN:HG3	4:B:2173:HOH:O	2.15	0.47
1:A:1:ALA:HB3	1:A:7:ALA:HB1	1.96	0.47
1:A:179:TRP:HE3	4:A:1682:HOH:O	1.97	0.47
1:B:501:ALA:HB3	4:B:2119:HOH:O	2.13	0.47
1:B:754:CYS:CB	1:B:763:ILE:HD11	2.45	0.47
1:A:436:THR:HG22	1:A:436:THR:OXT	2.15	0.46
1:B:827:PRO:HG3	1:B:838:GLN:HG2	1.97	0.46
1:B:580:GLY:HA3	1:B:623:TRP:CE3	2.51	0.46
1:B:816:LYS:HE2	1:B:936:THR:CG2	2.43	0.46
1:B:719:ARG:O	1:B:723:GLN:HG3	2.16	0.46
1:A:426:SER:O	1:A:427:ASN:HB2	2.15	0.45
1:B:921:GLN:NE2	3:B:1946:GOL:H32	2.31	0.45
3:A:1446:GOL:H32	4:A:1660:HOH:O	2.16	0.45
1:A:141:SER:O	1:A:145:ARG:HG3	2.16	0.45
1:B:502:GLU:HB2	1:B:801:ASN:OD1	2.17	0.45
1:B:613:MET:HE2	1:B:662:ASP:HB3	1.98	0.45
1:A:144:GLN:HE22	1:A:148:ASN:HA	1.82	0.45
1:B:831:THR:HG22	4:B:2306:HOH:O	2.16	0.44
1:A:113:MET:HE2	1:A:162:ASP:HB3	1.99	0.44
1:B:743:SER:OG	1:B:854:ASP:OD2	2.35	0.43
1:B:845:ALA:O	1:B:849:GLN:HG2	2.18	0.43
1:A:432:ARG:HD3	4:A:1552:HOH:O	2.19	0.43
1:B:519:ALA:HB2	1:B:766:TRP:CE3	2.54	0.42
1:B:910:VAL:HA	1:B:921:GLN:NE2	2.32	0.42
1:B:672:TYR:HB3	1:B:705:GLN:NE2	2.34	0.42
1:B:540:MET:HE1	1:B:579:ARG:N	2.35	0.42
1:B:842:CYS:HA	1:B:848:GLN:OE1	2.19	0.41
1:B:763:ILE:HD12	1:B:763:ILE:N	2.35	0.41
1:A:52:THR:O	1:A:59:PHE:HA	2.20	0.41
1:A:293:TYR:CD1	1:A:293:TYR:C	2.99	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:789:LYS:CD	1:B:891:ARG:NH1	2.84	0.41
1:A:168:CYS:HA	1:A:201:CYS:O	2.20	0.41
1:A:172:TYR:HB3	1:A:205:GLN:NE2	2.35	0.41
1:B:822:ARG:HG2	1:B:822:ARG:NH1	2.36	0.41
1:B:822:ARG:HE	1:B:915:ALA:HA	1.84	0.41
1:B:694:GLN:HG2	4:B:2253:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/436 (97%)	411 (97%)	12 (3%)	0	100	100
1	B	423/436 (97%)	411 (97%)	12 (3%)	0	100	100
All	All	846/872 (97%)	822 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/341 (98%)	332 (99%)	2 (1%)	78	66
1	B	334/341 (98%)	331 (99%)	3 (1%)	70	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	668/682 (98%)	663 (99%)	5 (1%)	76	63

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	MET
1	A	280	PRO
1	B	540	MET
1	B	601	LEU
1	B	780	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	194	GLN
1	A	223	GLN
1	A	284	ASN
1	B	511	GLN
1	B	828	ASN
1	B	893	ASN
1	B	921	GLN
1	B	927	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	1443	-	5,5,5	0.14	0	5,5,5	0.33	0
3	GOL	A	1446	-	5,5,5	0.19	0	5,5,5	0.40	0
3	GOL	B	1944	-	5,5,5	0.14	0	5,5,5	0.34	0
3	GOL	B	1941	-	5,5,5	0.14	0	5,5,5	0.32	0
2	SO4	A	1441	-	4,4,4	0.36	0	6,6,6	0.08	0
3	GOL	B	1942	-	5,5,5	0.13	0	5,5,5	0.32	0
3	GOL	B	1945	-	5,5,5	0.15	0	5,5,5	0.35	0
3	GOL	A	1444	-	5,5,5	0.15	0	5,5,5	0.34	0
3	GOL	A	1442	-	5,5,5	0.14	0	5,5,5	0.34	0
3	GOL	A	1445	-	5,5,5	0.18	0	5,5,5	0.35	0
3	GOL	B	1943	-	5,5,5	0.14	0	5,5,5	0.34	0
3	GOL	B	1946	-	5,5,5	0.14	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1443	-	-	0/4/4/4	-
3	GOL	A	1446	-	-	0/4/4/4	-
3	GOL	B	1944	-	-	0/4/4/4	-
3	GOL	B	1941	-	-	0/4/4/4	-
3	GOL	B	1942	-	-	0/4/4/4	-
3	GOL	B	1945	-	-	0/4/4/4	-
3	GOL	A	1444	-	-	0/4/4/4	-
3	GOL	A	1442	-	-	0/4/4/4	-
3	GOL	A	1445	-	-	0/4/4/4	-
3	GOL	B	1943	-	-	0/4/4/4	-
3	GOL	B	1946	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1446	GOL	1	0
3	A	1442	GOL	1	0
3	A	1445	GOL	3	0
3	B	1946	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/436 (97%)	0.39	40 (9%) 14 14	9, 15, 33, 44	0
1	B	427/436 (97%)	0.55	78 (18%) 3 3	8, 13, 46, 58	0
All	All	854/872 (97%)	0.47	118 (13%) 6 6	8, 14, 39, 58	0

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	819	GLY	9.4
1	A	303	GLY	9.2
1	B	501	ALA	8.5
1	B	818	VAL	7.7
1	B	813	GLY	7.5
1	B	936	THR	6.5
1	A	1	ALA	6.3
1	A	436	THR	6.2
1	B	928	GLY	6.2
1	A	92	TRP	6.0
1	B	842	CYS	5.9
1	B	910	VAL	5.2
1	B	845	ALA	5.2
1	B	815	ILE	5.0
1	B	839	LEU	4.9
1	B	822	ARG	4.9
1	B	934	THR	4.8
1	B	814	GLN	4.8
1	A	100	THR	4.8
1	B	933	TRP	4.6
1	B	918	THR	4.5
1	B	820	SER	4.5
1	B	915	ALA	4.5
1	B	914	THR	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	840	TYR	4.4
1	B	817	GLY	4.3
1	B	821	GLY	4.2
1	A	435	ARG	4.2
1	B	919	LEU	4.1
1	A	56	ARG	3.9
1	B	843	HIS	3.9
1	B	823	CYS	3.9
1	B	831	THR	3.9
1	B	932	ARG	3.9
1	B	935	ARG	3.9
1	A	179	TRP	3.9
1	B	916	ASN	3.8
1	B	816	LYS	3.7
1	B	841	ASP	3.7
1	B	927	ASN	3.7
1	B	824	LEU	3.7
1	B	909	ALA	3.7
1	B	913	GLY	3.7
1	B	848	GLN	3.6
1	B	899	VAL	3.6
1	B	929	SER	3.6
1	B	925	CYS	3.5
1	B	844	SER	3.5
1	A	96	LEU	3.4
1	B	896	GLY	3.4
1	B	849	GLN	3.4
1	A	91	GLY	3.4
1	A	146	THR	3.4
1	B	911	GLY	3.4
1	B	917	GLY	3.4
1	B	923	TYR	3.3
1	B	931	GLN	3.3
1	A	98	GLY	3.3
1	B	828	ASN	3.3
1	A	344	SER	3.3
1	B	912	GLY	3.2
1	A	89	GLN	3.2
1	A	145	ARG	3.2
1	B	846	THR	3.2
1	A	97	SER	3.2
1	B	893	ASN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	926	SER	3.1
1	A	95	SER	3.1
1	B	897	SER	3.1
1	A	140	ASP	3.1
1	B	876	LYS	3.0
1	B	826	VAL	3.0
1	B	825	ASP	3.0
1	B	906	CYS	3.0
1	B	908	ASP	3.0
1	A	88	GLN	2.9
1	B	851	THR	2.8
1	A	58	GLN	2.8
1	A	90	PRO	2.8
1	B	850	TRP	2.8
1	B	895	ASP	2.8
1	A	141	SER	2.8
1	A	94	GLN	2.8
1	A	101	LEU	2.8
1	B	898	ILE	2.7
1	B	847	ASN	2.7
1	A	134	GLY	2.7
1	B	907	LEU	2.7
1	B	922	LEU	2.7
1	B	930	ASN	2.6
1	A	143	LEU	2.6
1	B	920	ILE	2.6
1	B	784	ASN	2.6
1	B	892	LEU	2.6
1	B	829	ALA	2.6
1	B	894	SER	2.5
1	A	147	GLY	2.5
1	B	874	GLY	2.5
1	A	50	ASP	2.4
1	A	87	SER	2.4
1	A	99	SER	2.4
1	A	84	ALA	2.4
1	A	93	MET	2.3
1	B	837	VAL	2.3
1	B	838	GLN	2.3
1	A	104	ALA	2.3
1	A	135	SER	2.2
1	A	54	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	891	ARG	2.2
1	A	55	GLN	2.2
1	A	59	PHE	2.2
1	A	139	ARG	2.1
1	B	869	ALA	2.1
1	A	142	ASN	2.1
1	A	57	GLY	2.0
1	B	877	VAL	2.0
1	B	645	ARG	2.0
1	B	924	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	1442	6/6	0.58	0.25	50,52,52,53	0
3	GOL	B	1942	6/6	0.65	0.22	41,43,43,43	0
3	GOL	B	1941	6/6	0.68	0.25	35,42,44,45	0
3	GOL	A	1445	6/6	0.68	0.31	28,33,34,40	0
3	GOL	A	1446	6/6	0.69	0.23	21,27,32,33	0
3	GOL	B	1945	6/6	0.74	0.17	45,45,46,46	0
3	GOL	A	1444	6/6	0.78	0.18	36,37,37,38	0
3	GOL	B	1946	6/6	0.80	0.19	42,43,43,44	0
3	GOL	B	1943	6/6	0.83	0.17	44,45,45,46	0
3	GOL	A	1443	6/6	0.92	0.11	21,23,24,25	0
2	SO4	A	1441	5/5	0.93	0.14	24,26,31,31	0
3	GOL	B	1944	6/6	0.94	0.08	19,21,21,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.