



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:38 PM UTC

PDB ID : 6D1V / pdb_00006d1v
Title : Crystal structure of E. coli RppH-DapF complex, monomer bound to RNA
Authors : Gao, A.; Serganov, A.
Deposited on : 2018-04-12
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)
RNA backbone	3983	1006 (2.22-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	22% 92% 8%
2	B	160	19% 89% 9% ..
3	C	3	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	205	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3839 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diaminopimelate epimerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2113	1327	382	389	15			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P0A6K1

- Molecule 2 is a protein called RNA pyrophosphohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	159	Total	C	N	O	S	0	1	0
			1331	850	242	231	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	SER	-	expression tag	UNP P0A776
B	159	ALA	GLN	conflict	UNP P0A776

- Molecule 3 is a RNA chain called RNA (5'-D*(APC))-R(P*GP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O	P	0	0	0
			58	21	10	22	5			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 5 3 2	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Cl 3 3	0	0
5	B	2	Total Cl 2 2	0	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	3	Total 3	O 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	141	Total 141	O 141	0	0
7	B	138	Total 138	O 138	0	0
7	C	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Diaminopimelate epimerase



- Molecule 2: RNA pyrophosphohydrolase



- Molecule 3: RNA (5'-D*(APC))-R(P*GP*U)-3')



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	160.64Å 191.99Å 51.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.60 – 1.81 61.60 – 1.81	Depositor EDS
% Data completeness (in resolution range)	96.8 (61.60-1.81) 96.7 (61.60-1.81)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 1.81Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.199 , 0.220 0.205 , 0.223	Depositor DCC
R_{free} test set	2000 reflections (2.75%)	wwPDB-VP
Wilson B-factor (Å ²)	38.1	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3839	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, GOL, APC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2156	0.60	2/2921 (0.1%)
2	B	0.44	0/1367	0.67	2/1847 (0.1%)
3	C	0.75	0/29	0.45	0/44
All	All	0.42	0/3552	0.63	4/4812 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	96	LYS	N-CA-C	-6.08	99.32	109.24
1	A	162	ILE	N-CA-CB	-5.54	103.76	111.41
2	B	156	MET	CA-C-O	5.32	126.41	120.82
1	A	144	ALA	CB-CA-C	-5.06	110.72	116.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2113	0	2079	14	0
2	B	1331	0	1315	16	0
3	C	58	0	24	0	0
4	A	23	0	29	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	18	0	24	9	0
5	A	3	0	0	0	0
5	B	2	0	0	0	0
6	B	3	0	0	0	0
7	A	141	0	0	0	0
7	B	138	0	0	1	0
7	C	9	0	0	0	0
All	All	3839	0	3471	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:ASN:HD22	4:A:802:GOL:H2	1.58	0.67
2:B:113:MET:HG3	4:B:206:GOL:H2	1.78	0.66
2:B:47:GLU:HB2	4:B:204:GOL:H11	1.81	0.63
2:B:142:ASP:CA	4:B:205:GOL:H32	2.30	0.62
2:B:142:ASP:N	4:B:205:GOL:H32	2.16	0.61
1:A:90:THR:HA	2:B:156:MET:SD	2.41	0.61
2:B:131:TYR:CZ	2:B:135:GLN:HG3	2.41	0.55
1:A:162:ILE:O	1:A:162:ILE:HG22	2.10	0.52
2:B:142:ASP:OD1	4:B:205:GOL:H12	2.11	0.51
1:A:111:ASP:OD1	1:A:111:ASP:N	2.41	0.50
1:A:91:ASN:ND2	2:B:156:MET:O	2.44	0.50
1:A:85:ARG:HD3	1:A:108:VAL:HG21	1.95	0.48
1:A:22:ASN:CG	4:B:205:GOL:H2	2.39	0.48
1:A:204:LEU:HD23	1:A:242:LEU:HD21	1.96	0.47
1:A:164:VAL:HG11	1:A:170:ALA:HB2	1.97	0.46
1:A:167:VAL:HG13	1:A:195:GLN:HB2	1.97	0.46
2:B:82:ARG:HD2	7:B:374:HOH:O	2.17	0.45
2:B:89:LYS:HE3	2:B:90:PRO:HA	1.98	0.45
1:A:161:VAL:HG21	1:A:224:ALA:HA	1.99	0.45
1:A:137:LYS:CD	1:A:231:GLN:HA	2.47	0.44
2:B:142:ASP:HA	4:B:205:GOL:H32	1.98	0.44
1:A:0:SER:HB3	1:A:273:HIS:NE2	2.34	0.42
2:B:100:PHE:HE2	4:B:204:GOL:H12	1.84	0.42
2:B:13:ILE:HG22	2:B:15:ILE:HG13	2.02	0.41
1:A:182:HIS:CE1	1:A:184:ARG:HG3	2.56	0.41
2:B:142:ASP:HB2	4:B:205:GOL:H31	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:129:TYR:CZ	2:B:148:MET:HE3	2.57	0.40
2:B:93:ILE:H	2:B:93:ILE:HG13	1.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	267 (98%)	6 (2%)	0	100	100
2	B	158/160 (99%)	156 (99%)	2 (1%)	0	100	100
All	All	431/435 (99%)	423 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/225 (99%)	221 (99%)	2 (1%)	70	62
2	B	143/142 (101%)	142 (99%)	1 (1%)	76	70
All	All	366/367 (100%)	363 (99%)	3 (1%)	73	67

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	ASP
1	A	162	ILE
2	B	113	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	44	GLN
1	A	72	GLN
1	A	75	ASN
1	A	256	HIS
2	B	37	GLN
2	B	73	ASN
2	B	97	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	0/3	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 8 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	204	-	5,5,5	0.26	0	5,5,5	0.32	0
4	GOL	A	802	-	5,5,5	0.36	0	5,5,5	0.44	0
4	GOL	A	804	-	5,5,5	0.27	0	5,5,5	0.32	0
4	GOL	A	801	-	5,5,5	0.33	0	5,5,5	0.35	0
4	GOL	B	206	-	5,5,5	0.34	0	5,5,5	0.66	0
4	GOL	B	205	-	5,5,5	0.77	0	5,5,5	0.99	0
4	GOL	A	803	-	4,4,5	0.67	0	4,4,5	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	204	-	-	2/4/4/4	-
4	GOL	A	802	-	-	3/4/4/4	-
4	GOL	A	804	-	-	4/4/4/4	-
4	GOL	A	801	-	-	4/4/4/4	-
4	GOL	B	206	-	-	2/4/4/4	-
4	GOL	B	205	-	-	3/4/4/4	-
4	GOL	A	803	-	-	2/2/2/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	801	GOL	C1-C2-C3-O3
4	A	802	GOL	C1-C2-C3-O3
4	A	804	GOL	C1-C2-C3-O3
4	B	204	GOL	O1-C1-C2-C3
4	B	205	GOL	O1-C1-C2-C3
4	B	205	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	804	GOL	O1-C1-C2-C3
4	B	206	GOL	O1-C1-C2-C3
4	A	801	GOL	O2-C2-C3-O3
4	A	804	GOL	O2-C2-C3-O3
4	B	204	GOL	O1-C1-C2-O2
4	B	205	GOL	O1-C1-C2-O2
4	A	802	GOL	O2-C2-C3-O3
4	B	206	GOL	O1-C1-C2-O2
4	A	803	GOL	O1-C1-C2-O2
4	A	803	GOL	O1-C1-C2-C3
4	A	801	GOL	O1-C1-C2-O2
4	A	804	GOL	O1-C1-C2-O2
4	A	801	GOL	O1-C1-C2-C3
4	A	802	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	204	GOL	2	0
4	A	802	GOL	1	0
4	B	206	GOL	1	0
4	B	205	GOL	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	1.17	60 (21%) 2 2	26, 50, 81, 107	0
2	B	159/160 (99%)	0.81	31 (19%) 3 3	20, 38, 71, 82	1 (0%)
3	C	2/3 (66%)	0.91	0 100 100	45, 45, 45, 62	0
All	All	436/438 (99%)	1.04	91 (20%) 2 2	20, 46, 75, 107	1 (0%)

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	158	LEU	5.8
1	A	147	THR	4.9
2	B	113	MET	4.8
1	A	196	VAL	4.7
1	A	271	PHE	4.7
2	B	0	SER	4.6
1	A	72	GLN	4.5
1	A	164	VAL	4.3
2	B	156	MET	4.2
1	A	171	ALA	4.1
1	A	193	PHE	4.0
2	B	74	TRP	4.0
1	A	169	THR	3.9
1	A	140	ILE	3.9
1	A	149	LEU	3.8
1	A	170	ALA	3.5
1	A	198	LYS	3.5
1	A	71	ALA	3.5
2	B	43	GLY	3.5
1	A	163	GLN	3.4
1	A	162	ILE	3.3
1	A	172	VAL	3.3
1	A	148	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	73	ASN	3.2
1	A	10	GLY	3.1
2	B	42	PRO	3.0
1	A	0	SER	2.9
2	B	1	MET	2.9
1	A	141	MET	2.8
2	B	79	LEU	2.8
1	A	136	GLU	2.8
1	A	211	ALA	2.8
1	A	144	ALA	2.7
2	B	52	ARG	2.7
2	B	89	LYS	2.7
2	B	157	SER	2.7
1	A	214	THR	2.7
2	B	96	LYS	2.7
1	A	160	CYS	2.6
1	A	161	VAL	2.6
1	A	178	VAL	2.6
1	A	9	LEU	2.6
1	A	179	LEU	2.6
2	B	93	ILE	2.6
1	A	135	ALA	2.6
1	A	227	VAL	2.6
1	A	142	ARG	2.6
2	B	114	GLN	2.6
1	A	143	ALA	2.5
1	A	86	LEU	2.5
2	B	88	THR	2.5
1	A	175	LEU	2.5
1	A	165	ASP	2.5
2	B	72	ARG	2.4
2	B	112	ASN	2.4
2	B	2	ILE	2.4
2	B	4	ASP	2.4
1	A	174	THR	2.4
1	A	150	CYS	2.4
2	B	75	LEU	2.4
2	B	90	PRO	2.4
2	B	31	HIS	2.4
2	B	86	TRP	2.3
1	A	251	TRP	2.3
1	A	189	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	212	GLY	2.3
1	A	91	ASN	2.2
2	B	85	ARG	2.2
1	A	183	GLU	2.2
1	A	274	LEU	2.2
2	B	46	ALA	2.2
1	A	137	LYS	2.2
1	A	250	ALA	2.2
1	A	138	THR	2.2
1	A	168	ASP	2.2
1	A	181	SER	2.2
1	A	123	PHE	2.1
1	A	133	ASN	2.1
2	B	41	ASN	2.1
1	A	167	VAL	2.1
1	A	273	HIS	2.1
1	A	231	GLN	2.1
2	B	7	TYR	2.1
2	B	40	ILE	2.1
1	A	197	VAL	2.1
1	A	38	LEU	2.1
1	A	182	HIS	2.0
1	A	146	GLN	2.0
1	A	265	VAL	2.0
1	A	31	ARG	2.0
2	B	66	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	802	6/6	0.67	0.28	62,64,67,68	0
4	GOL	B	204	6/6	0.69	0.21	47,59,61,64	0
4	GOL	B	206	6/6	0.76	0.22	52,58,64,65	0
4	GOL	B	205	6/6	0.77	0.20	45,56,56,57	0
4	GOL	A	804	6/6	0.77	0.24	58,60,67,69	0
4	GOL	A	801	6/6	0.81	0.19	59,61,63,66	0
4	GOL	A	803	5/6	0.83	0.22	50,52,60,61	0
5	CL	A	807	1/1	0.85	0.28	30,30,30,30	0
5	CL	B	208	1/1	0.89	0.21	87,87,87,87	0
5	CL	A	806	1/1	0.91	0.15	76,76,76,76	0
5	CL	A	805	1/1	0.97	0.09	39,39,39,39	0
6	MG	B	201	1/1	0.98	0.05	41,41,41,41	0
5	CL	B	207	1/1	0.99	0.22	60,60,60,60	1
6	MG	B	202	1/1	0.99	0.03	32,32,32,32	0
6	MG	B	203	1/1	0.99	0.06	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.