



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:46 AM UTC

PDB ID : 6D2I / pdb_00006d2i
Title : JAK2 Pseudokinase V617F in complex with AT9283
Authors : Li, Q.; Li, K.; Eck, M.J.
Deposited on : 2018-04-13
Resolution : 3.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	26% 88% 12%
1	B	273	27% 85% 14%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	0	0
			2198	1404	380	403	11			
1	B	272	Total	C	N	O	S	0	0	0
			2191	1399	379	402	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	659	ALA	TRP	engineered mutation	UNP Q506Q0
A	777	ALA	TRP	engineered mutation	UNP Q506Q0
A	794	HIS	PHE	engineered mutation	UNP Q506Q0
B	659	ALA	TRP	engineered mutation	UNP Q506Q0
B	777	ALA	TRP	engineered mutation	UNP Q506Q0
B	794	HIS	PHE	engineered mutation	UNP Q506Q0

- Molecule 2 is 1-cyclopropyl-3-{3-[5-(morpholin-4-ylmethyl)-1H-benzimidazol-2-yl]-1H-pyrazol-4-yl}urea (CCD ID: 35R) (formula: C₁₉H₂₃N₇O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	19	7	2		
2	B	1	Total	C	N	O	0	0
			28	19	7	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase



- Molecule 1: Tyrosine-protein kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.91Å 91.13Å 113.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.26 – 3.19 42.26 – 3.19	Depositor EDS
% Data completeness (in resolution range)	97.9 (42.26-3.19) 98.0 (42.26-3.19)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 3.19Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.271 , 0.290 0.275 , 0.292	Depositor DCC
R_{free} test set	574 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	79.5	Xtriage
Anisotropy	0.446	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	4445	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 35R

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/2247	0.27	0/3037
1	B	0.09	0/2240	0.28	0/3027
All	All	0.09	0/4487	0.28	0/6064

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2198	0	2199	20	0
1	B	2191	0	2190	24	0
2	A	28	0	23	2	0
2	B	28	0	23	3	0
All	All	4445	0	4435	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:629:VAL:HG21	1:A:680:LEU:HB3	1.68	0.74
1:B:677:LYS:HE2	2:B:901:35R:H4	1.74	0.69
1:B:629:VAL:HG21	1:B:680:LEU:HB3	1.78	0.64
1:A:743:LEU:HD12	1:A:746:ILE:HD11	1.80	0.63
1:A:717:PRO:HG3	1:A:754:LEU:HD13	1.83	0.61
1:A:559:ILE:HG13	2:A:901:35R:H6	1.83	0.59
1:B:782:ASN:OD1	1:B:783:LEU:N	2.35	0.59
1:B:726:ASN:HB3	1:B:729:ASN:HB2	1.86	0.57
1:B:559:ILE:HG13	2:B:901:35R:H6	1.86	0.57
1:B:610:VAL:HG21	1:B:680:LEU:HD12	1.87	0.56
1:A:673:ASN:OD1	1:A:715:ARG:NH2	2.37	0.56
1:B:673:ASN:OD1	1:B:715:ARG:NH2	2.40	0.55
1:B:638:LEU:HD11	1:B:746:ILE:HG22	1.89	0.54
1:A:638:LEU:HD11	1:A:746:ILE:HG22	1.88	0.54
1:B:641:ASN:O	1:B:645:ILE:HG12	2.10	0.51
1:B:604:LEU:HD22	1:B:663:PHE:HE2	1.76	0.51
1:B:600:MET:HE3	1:B:669:LEU:HB3	1.93	0.50
1:A:673:ASN:ND2	1:A:700:PRO:O	2.33	0.50
1:B:743:LEU:HD23	1:B:746:ILE:HD11	1.94	0.50
1:A:610:VAL:HG21	1:A:680:LEU:HD12	1.95	0.49
1:A:641:ASN:O	1:A:645:ILE:HG12	2.13	0.48
1:A:649:TRP:CD1	1:A:685:GLU:HB2	2.51	0.45
1:B:579:LEU:HD13	2:B:901:35R:C13	2.45	0.45
1:A:559:ILE:HG23	1:A:579:LEU:HD11	1.99	0.44
1:A:629:VAL:HG22	2:A:901:35R:N12	2.34	0.43
1:B:691:ASN:HA	1:B:692:PRO:HD3	1.88	0.43
1:B:779:GLU:OE2	1:B:782:ASN:ND2	2.52	0.43
1:B:733:ALA:HA	1:B:736:LYS:HD2	2.01	0.43
1:B:542:ASN:HB2	1:B:618:CYS:HA	2.00	0.42
1:B:730:LEU:HD12	1:B:730:LEU:HA	1.93	0.42
1:A:799:ARG:NH1	1:B:627:GLU:OE2	2.53	0.41
1:B:607:LYS:O	1:B:696:LYS:HE2	2.21	0.41
1:B:601:MET:HE3	1:B:699:ASP:HA	2.00	0.41
1:A:772:LEU:HD23	1:A:773:PRO:HD2	2.03	0.41
1:A:595:PHE:CZ	1:A:622:ASN:HB3	2.55	0.41
1:B:559:ILE:HG23	1:B:579:LEU:HD21	2.02	0.41
1:A:683:ARG:NH1	1:A:692:PRO:O	2.51	0.41
1:A:546:ILE:HD11	1:A:565:ARG:HD2	2.02	0.41
1:A:607:LYS:O	1:A:696:LYS:HE2	2.21	0.41
1:B:556:PHE:HB2	1:B:702:ILE:HD12	2.04	0.40
1:A:629:VAL:HA	1:A:682:ILE:HD13	2.03	0.40
1:A:537:PHE:CZ	1:A:615:VAL:HG21	2.57	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:759:SER:HA	1:B:762:LYS:HD2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/273 (99%)	256 (94%)	14 (5%)	1 (0%)	30	62
1	B	270/273 (99%)	258 (96%)	11 (4%)	1 (0%)	30	62
All	All	541/546 (99%)	514 (95%)	25 (5%)	2 (0%)	30	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	618	CYS
1	B	618	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/244 (100%)	243 (100%)	1 (0%)	84	86
1	B	243/244 (100%)	242 (100%)	1 (0%)	84	86
All	All	487/488 (100%)	485 (100%)	2 (0%)	84	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	604	LEU
1	B	572	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	641	ASN
1	A	786	ASN
1	B	548	ASN
1	B	622	ASN
1	B	626	GLN
1	B	641	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	35R	B	901	-	31,32,32	3.07	16 (51%)	41,45,45	3.36	13 (31%)
2	35R	A	901	-	31,32,32	3.08	16 (51%)	41,45,45	3.35	12 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	35R	B	901	-	-	3/16/26/26	0/5/5/5
2	35R	A	901	-	-	3/16/26/26	0/5/5/5

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	35R	C16-N15	8.91	1.54	1.38
2	B	901	35R	C16-N15	8.89	1.54	1.38
2	A	901	35R	C13-C14	6.86	1.55	1.46
2	B	901	35R	C13-C14	6.73	1.54	1.46
2	A	901	35R	C29-N30	-5.82	1.28	1.39
2	B	901	35R	C29-N30	-5.80	1.28	1.39
2	B	901	35R	C9-N10	4.97	1.41	1.34
2	A	901	35R	C9-N10	4.96	1.41	1.34
2	B	901	35R	C2-N3	4.17	1.47	1.35
2	A	901	35R	C2-N3	4.15	1.46	1.35
2	A	901	35R	C2-N7	3.37	1.47	1.39
2	B	901	35R	C14-N30	-3.34	1.28	1.32
2	A	901	35R	C14-N30	-3.32	1.28	1.32
2	B	901	35R	C2-N7	3.32	1.47	1.39
2	B	901	35R	C16-C29	2.82	1.44	1.40
2	A	901	35R	C20-C19	2.82	1.56	1.51
2	B	901	35R	C20-C19	2.81	1.56	1.51
2	A	901	35R	C16-C29	2.81	1.44	1.40
2	A	901	35R	N10-N12	2.78	1.41	1.36
2	B	901	35R	N10-N12	2.73	1.41	1.36
2	A	901	35R	C13-N12	2.51	1.37	1.34
2	B	901	35R	C13-N12	2.48	1.37	1.34
2	A	901	35R	C6-C4	2.47	1.54	1.48
2	B	901	35R	C6-C4	2.45	1.54	1.48
2	A	901	35R	C8-N7	2.43	1.47	1.41
2	B	901	35R	C5-C4	2.35	1.54	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	35R	C8-N7	2.35	1.47	1.41
2	B	901	35R	C18-C17	2.33	1.42	1.38
2	A	901	35R	C5-C4	2.32	1.54	1.48
2	A	901	35R	C18-C17	2.31	1.42	1.38
2	B	901	35R	C27-C26	2.06	1.58	1.50
2	A	901	35R	C27-C26	2.05	1.58	1.50

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	35R	C29-N30-C14	17.14	122.07	104.68
2	A	901	35R	C29-N30-C14	17.12	122.05	104.68
2	B	901	35R	C16-C29-N30	-5.65	101.68	109.42
2	A	901	35R	C16-C29-N30	-5.59	101.76	109.42
2	A	901	35R	N15-C14-N30	-5.00	107.87	113.22
2	B	901	35R	N15-C14-N30	-4.97	107.90	113.22
2	B	901	35R	C16-N15-C14	-4.10	102.04	106.85
2	A	901	35R	C16-N15-C14	-4.04	102.11	106.85
2	A	901	35R	C28-C29-N30	3.46	136.77	129.31
2	B	901	35R	C28-C29-N30	3.45	136.75	129.31
2	A	901	35R	C8-C9-N10	3.27	107.83	104.76
2	B	901	35R	C8-C9-N10	3.22	107.78	104.76
2	A	901	35R	C14-C13-N12	3.16	127.17	119.86
2	B	901	35R	C14-C13-N12	3.12	127.07	119.86
2	B	901	35R	C17-C16-C29	-3.06	118.75	122.10
2	A	901	35R	C13-N12-N10	3.06	107.92	104.73
2	B	901	35R	C13-N12-N10	3.04	107.89	104.73
2	B	901	35R	C9-N10-N12	-2.98	108.03	112.59
2	A	901	35R	C9-N10-N12	-2.97	108.04	112.59
2	A	901	35R	C17-C16-C29	-2.96	118.85	122.10
2	B	901	35R	C5-C4-N3	-2.30	115.50	118.52
2	A	901	35R	C6-C4-N3	-2.25	115.57	118.52
2	B	901	35R	C6-C4-N3	-2.14	115.71	118.52
2	B	901	35R	C17-C16-N15	2.04	134.96	130.83
2	A	901	35R	C17-C16-N15	2.03	134.94	130.83

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	901	35R	O1-C2-N3-C4
2	B	901	35R	O1-C2-N3-C4

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	A	901	35R	N7-C2-N3-C4
2	B	901	35R	N7-C2-N3-C4
2	B	901	35R	C19-C20-N21-C27
2	A	901	35R	C19-C20-N21-C27

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	901	35R	3	0
2	A	901	35R	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/273 (100%)	1.54	70 (25%)  	43, 73, 113, 126	0
1	B	272/273 (99%)	1.57	73 (26%)  	45, 71, 100, 116	0
All	All	545/546 (99%)	1.56	143 (26%)  	43, 72, 108, 126	0

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	787	CYS	8.0
1	A	742	THR	5.9
1	A	733	ALA	5.5
1	B	787	CYS	5.3
1	A	807	SER	4.8
1	B	627	GLU	4.2
1	B	682	ILE	4.2
1	A	802	ILE	4.1
1	B	742	THR	4.0
1	B	633	SER	4.0
1	A	678	ASN	3.9
1	A	746	ILE	3.9
1	B	708	PRO	3.8
1	B	802	ILE	3.8
1	A	795	ARG	3.8
1	B	666	GLU	3.8
1	A	729	ASN	3.7
1	B	701	GLY	3.7
1	A	704	ILE	3.7
1	A	599	SER	3.5
1	B	624	LEU	3.5
1	B	711	ILE	3.5
1	B	678	ASN	3.4
1	B	687	ARG	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	729	ASN	3.4
1	A	732	LEU	3.4
1	A	601	MET	3.4
1	A	633	SER	3.3
1	A	665	GLU	3.3
1	B	612	ASN	3.3
1	A	563	VAL	3.3
1	A	725	GLU	3.3
1	A	673	ASN	3.3
1	B	679	ILE	3.3
1	B	634	LEU	3.3
1	A	615	VAL	3.2
1	B	685	GLU	3.2
1	B	651	LEU	3.2
1	B	704	ILE	3.1
1	B	716	ILE	3.1
1	A	538	HIS	3.1
1	B	599	SER	3.1
1	A	617	PHE	3.1
1	B	631	PHE	3.1
1	B	778	ALA	3.1
1	A	623	ILE	3.1
1	A	712	LEU	3.0
1	B	730	LEU	3.0
1	B	674	VAL	3.0
1	A	793	ASP	3.0
1	B	743	LEU	2.9
1	A	616	CYS	2.9
1	B	780	LEU	2.9
1	A	734	THR	2.9
1	A	728	LYS	2.9
1	A	647	ILE	2.9
1	A	544	ASP	2.8
1	A	693	PRO	2.8
1	B	591	SER	2.7
1	B	601	MET	2.7
1	A	683	ARG	2.7
1	B	672	GLY	2.7
1	B	669	LEU	2.7
1	A	766	TYR	2.6
1	B	712	LEU	2.6
1	B	603	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	743	LEU	2.5
1	B	579	LEU	2.5
1	B	763	LEU	2.5
1	B	710	ASP	2.5
1	A	631	PHE	2.5
1	A	778	ALA	2.5
1	B	577	GLU	2.5
1	A	669	LEU	2.5
1	B	738	SER	2.5
1	B	610	VAL	2.5
1	A	596	GLU	2.5
1	A	785	ASN	2.5
1	B	620	ASP	2.5
1	B	734	THR	2.5
1	A	545	LEU	2.5
1	A	685	GLU	2.4
1	B	635	ASP	2.4
1	B	728	LYS	2.4
1	B	629	VAL	2.4
1	A	660	ALA	2.4
1	A	798	PHE	2.4
1	A	536	VAL	2.4
1	A	627	GLU	2.4
1	A	780	LEU	2.4
1	B	735	ASP	2.4
1	A	539	LYS	2.4
1	B	665	GLU	2.4
1	B	593	SER	2.4
1	A	597	ALA	2.4
1	A	772	LEU	2.4
1	A	773	PRO	2.4
1	B	736	LYS	2.3
1	A	799	ARG	2.3
1	B	605	SER	2.3
1	B	647	ILE	2.3
1	B	570	TYR	2.3
1	B	615	VAL	2.3
1	B	598	ALA	2.3
1	B	617	PHE	2.3
1	A	621	GLU	2.3
1	B	684	GLU	2.3
1	A	792	PRO	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	594	PHE	2.3
1	A	634	LEU	2.2
1	B	592	GLU	2.2
1	A	711	ILE	2.2
1	B	587	HIS	2.2
1	B	767	GLU	2.2
1	B	638	LEU	2.2
1	B	537	PHE	2.2
1	A	794	HIS	2.2
1	B	692	PRO	2.2
1	B	625	VAL	2.1
1	B	739	PHE	2.1
1	A	639	LYS	2.1
1	B	733	ALA	2.1
1	A	577	GLU	2.1
1	A	569	ASP	2.1
1	A	537	PHE	2.1
1	B	548	ASN	2.1
1	B	655	LYS	2.1
1	A	609	LEU	2.1
1	A	804	ASP	2.1
1	B	789	ASP	2.1
1	A	559	ILE	2.0
1	A	774	ALA	2.0
1	A	701	GLY	2.0
1	A	708	PRO	2.0
1	A	610	VAL	2.0
1	B	541	ARG	2.0
1	B	732	LEU	2.0
1	A	542	ASN	2.0
1	A	770	HIS	2.0
1	A	614	GLY	2.0
1	B	762	LYS	2.0
1	A	684	GLU	2.0
1	B	754	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	35R	B	901	28/28	0.76	0.22	51,62,99,112	0
2	35R	A	901	28/28	0.82	0.21	44,55,92,105	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 35R B 901:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 35R A 901:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.