



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 11:11 PM UTC

PDB ID : 7D2Z / pdb_00007d2z
Title : Structure of sybody SR31 in complex with the SARS-CoV-2 S Receptor Binding domain (RBD)
Authors : Li, T.; Cai, H.; Yao, H.; Qin, W.; Li, D.
Deposited on : 2020-09-17
Resolution : 1.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	 6% 69% 12% 19%
2	B	213	 4% 92% 8%
3	C	7	 29% 71%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	B	706	-	-	X	-
5	GOL	B	712	-	-	X	-
6	ACT	B	701	-	-	X	-

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 2913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SR31 against SARS-CoV-2 RBD, non neutralizing.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	117	915	580	155	173	7	0	4	0


- Molecule 2 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	212	1674	1072	278	316	8	0	1	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	327	ALA	-	expression tag	UNP P0DTC2
B	328	GLY	-	expression tag	UNP P0DTC2
B	329	SER	-	expression tag	UNP P0DTC2
B	532	GLY	-	expression tag	UNP P0DTC2
B	533	THR	-	expression tag	UNP P0DTC2
B	534	LEU	-	expression tag	UNP P0DTC2
B	535	GLU	-	expression tag	UNP P0DTC2
B	536	VAL	-	expression tag	UNP P0DTC2
B	537	LEU	-	expression tag	UNP P0DTC2
B	538	PHE	-	expression tag	UNP P0DTC2
B	539	GLN	-	expression tag	UNP P0DTC2

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	7	81	46	2	33	0	0	0

- Molecule 4 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	3	1	2	0	0
4	A	1	3	1	2	0	0
4	A	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0
4	B	1	3	1	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
7	A	35	Total	O	0	0
			35	35		
7	B	131	Total	O	0	0
			131	131		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SR31 against SARS-CoV-2 RBD, non neutralizing



- Molecule 2: Spike protein S1



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.39Å 92.39Å 101.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.62 – 1.97 19.62 – 1.97	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.62-1.97) 99.9 (19.62-1.97)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.97Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.182 , 0.207 0.182 , 0.208	Depositor DCC
R_{free} test set	1738 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	42.3	Xtrriage
Anisotropy	0.279	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2913	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, FMT, GOL, ACT, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/948	0.49	0/1283
2	B	0.32	0/1723	0.51	0/2343
All	All	0.30	0/2671	0.51	0/3626

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	915	0	896	12	0
2	B	1674	0	1598	20	0
3	C	81	0	70	0	0
4	A	9	0	3	0	0
4	B	30	0	10	4	0
5	A	6	0	8	1	0
5	B	24	0	32	8	0
6	A	4	0	3	0	0
6	B	4	0	3	2	0
7	A	35	0	0	0	0
7	B	131	0	0	3	0
All	All	2913	0	2623	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 32 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:514:SER:HB2	5:B:712:GOL:H31	1.49	0.92
2:B:455:LEU:HD13	6:B:701:ACT:H2	1.74	0.69
1:A:13:GLN:HG2	1:A:114:SER:HB2	1.76	0.68
2:B:430:THR:O	5:B:712:GOL:H11	2.05	0.56
2:B:408:ARG:HG3	7:B:826:HOH:O	2.04	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/144 (83%)	117 (98%)	2 (2%)	0	100	100
2	B	211/213 (99%)	202 (96%)	9 (4%)	0	100	100
All	All	330/357 (92%)	319 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/113 (85%)	96 (100%)	0	100	100
2	B	184/183 (100%)	184 (100%)	0	100	100
All	All	280/296 (95%)	280 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	110	GLN
2	B	334	ASN
2	B	519	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	2,3	14,14,15	0.59	0	17,19,21	0.51	0
3	NAG	C	2	3	14,14,15	0.26	0	17,19,21	0.38	0
3	BMA	C	3	3	11,11,12	0.78	0	15,15,17	0.99	1 (6%)
3	MAN	C	4	3	11,11,12	0.70	0	15,15,17	0.89	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	C	5	3	11,11,12	1.08	1 (9%)	15,15,17	1.20	1 (6%)
3	FUC	C	6	3	10,10,11	1.18	0	14,14,16	0.89	1 (7%)
3	FUC	C	7	3	10,10,11	1.30	2 (20%)	14,14,16	0.96	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	1/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	FUC	C	6	3	-	-	0/1/1/1
3	FUC	C	7	3	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	7	FUC	C1-C2	2.69	1.58	1.52
3	C	5	MAN	C1-C2	2.45	1.58	1.52
3	C	7	FUC	C2-C3	2.40	1.56	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	MAN	C1-O5-C5	3.55	116.94	112.19
3	C	3	BMA	C1-O5-C5	3.16	116.42	112.19
3	C	7	FUC	C1-C2-C3	2.32	113.02	109.64
3	C	4	MAN	C1-O5-C5	2.30	115.27	112.19
3	C	6	FUC	O5-C5-C4	2.05	113.25	109.55

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	4	MAN	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FMT	B	710	-	2,2,2	0.66	0	1,1,1	0.11	0
4	FMT	B	704	-	2,2,2	0.66	0	1,1,1	0.13	0
4	FMT	B	705	-	2,2,2	0.70	0	1,1,1	0.14	0
4	FMT	B	703	-	2,2,2	0.69	0	1,1,1	0.23	0
4	FMT	B	715	-	2,2,2	0.50	0	1,1,1	0.17	0
6	ACT	A	604	-	3,3,3	1.55	0	3,3,3	1.57	1 (33%)
4	FMT	B	714	-	2,2,2	0.64	0	1,1,1	0.01	0
4	FMT	B	713	-	2,2,2	0.68	0	1,1,1	0.18	0
4	FMT	A	602	-	2,2,2	0.70	0	1,1,1	0.10	0
4	FMT	B	706	-	2,2,2	0.60	0	1,1,1	0.03	0
4	FMT	B	711	-	2,2,2	0.72	0	1,1,1	0.30	0
4	FMT	B	702	-	2,2,2	0.69	0	1,1,1	0.17	0
5	GOL	B	708	-	5,5,5	1.35	1 (20%)	5,5,5	1.05	0
6	ACT	B	701	-	3,3,3	1.34	0	3,3,3	1.59	1 (33%)
5	GOL	B	707	-	5,5,5	0.86	0	5,5,5	1.05	0
5	GOL	B	712	-	5,5,5	1.28	0	5,5,5	0.56	0
5	GOL	B	709	-	5,5,5	0.82	0	5,5,5	1.19	0
5	GOL	A	603	-	5,5,5	1.26	1 (20%)	5,5,5	0.75	0
4	FMT	A	601	-	2,2,2	0.64	0	1,1,1	0.00	0
4	FMT	A	605	-	2,2,2	0.73	0	1,1,1	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	708	-	-	4/4/4/4	-
5	GOL	B	707	-	-	2/4/4/4	-
5	GOL	B	712	-	-	2/4/4/4	-
5	GOL	B	709	-	-	2/4/4/4	-
5	GOL	A	603	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	708	GOL	C1-C2	2.14	1.59	1.51
5	A	603	GOL	C1-C2	2.05	1.59	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	701	ACT	OXT-C-O	2.24	130.35	122.03
6	A	604	ACT	OXT-C-O	2.05	129.65	122.03

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	707	GOL	O1-C1-C2-C3
5	B	708	GOL	C1-C2-C3-O3
5	B	709	GOL	C1-C2-C3-O3
5	B	709	GOL	O2-C2-C3-O3
5	B	707	GOL	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	704	FMT	1	0
4	B	714	FMT	1	0
4	B	706	FMT	2	0
6	B	701	ACT	2	0
5	B	707	GOL	3	0
5	B	712	GOL	5	0
5	A	603	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	117/144 (81%)	0.41	8 (6%) 23 31	23, 49, 82, 95	4 (3%)
2	B	212/213 (99%)	0.02	9 (4%) 40 50	29, 41, 75, 90	1 (0%)
All	All	329/357 (92%)	0.16	17 (5%) 33 42	23, 44, 77, 95	5 (1%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	486	PHE	4.5
2	B	328	GLY	3.6
2	B	330	PRO	3.5
2	B	332	ILE	3.0
1	A	-2	SER	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	C	4	11/12	0.28	0.14	100,112,116,119	0
3	BMA	C	3	11/12	0.67	0.13	90,94,108,109	0
3	FUC	C	6	10/11	0.68	0.15	85,93,101,103	0
3	MAN	C	5	11/12	0.75	0.13	92,96,106,106	0
3	NAG	C	1	14/15	0.90	0.09	51,70,83,84	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	FUC	C	7	10/11	0.90	0.10	46,56,60,60	0
3	NAG	C	2	14/15	0.92	0.09	58,71,77,82	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	FMT	B	711	3/3	0.65	0.18	72,72,73,73	0
5	GOL	B	709	6/6	0.65	0.17	60,66,74,83	0
5	GOL	A	603	6/6	0.69	0.19	43,56,64,64	0
4	FMT	A	605	3/3	0.72	0.15	66,66,67,73	0
6	ACT	A	604	4/4	0.76	0.17	58,63,67,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	FMT	B	715	3/3	0.81	0.27	43,43,46,50	0
5	GOL	B	707	6/6	0.81	0.14	47,62,66,72	0
5	GOL	B	708	6/6	0.82	0.14	51,60,66,68	0
4	FMT	B	703	3/3	0.83	0.13	68,68,68,71	0
4	FMT	B	714	3/3	0.84	0.24	56,56,61,61	0
4	FMT	B	704	3/3	0.87	0.23	54,54,57,62	0
6	ACT	B	701	4/4	0.87	0.22	45,45,61,62	0
4	FMT	B	702	3/3	0.88	0.11	59,59,68,69	0
4	FMT	A	602	3/3	0.88	0.12	59,59,61,64	0
4	FMT	B	713	3/3	0.89	0.14	49,49,60,73	0
4	FMT	B	706	3/3	0.90	0.14	48,48,51,66	0
5	GOL	B	712	6/6	0.90	0.10	37,50,56,63	0
4	FMT	B	710	3/3	0.90	0.22	49,49,59,61	0
4	FMT	A	601	3/3	0.90	0.12	44,44,55,56	0
4	FMT	B	705	3/3	0.93	0.10	50,50,58,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.