



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:18 PM UTC

PDB ID : 3D3K / pdb_00003d3k
Title : Crystal structure of human Edc3p
Authors : Ling, S.H.M.
Deposited on : 2008-05-12
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	6% 82% 8% 10%
1	B	259	6% 79% 10% 10%
1	C	259	4% 78% 11% 10%
1	D	259	5% 76% 12% 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enhancer of mRNA-decapping protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1802	1158	312	321	11			
1	B	233	Total	C	N	O	S	0	0	0
			1802	1158	312	321	11			
1	C	233	Total	C	N	O	S	0	0	0
			1802	1158	312	321	11			
1	D	233	Total	C	N	O	S	0	0	0
			1802	1158	312	321	11			



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	89	Total	O	0	0
			89	89		
2	B	94	Total	O	0	0
			94	94		
2	C	84	Total	O	0	0
			84	84		
2	D	81	Total	O	0	0
			81	81		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Enhancer of mRNA-decapping protein 3




• Molecule 1: Enhancer of mRNA-decapping protein 3




• Molecule 1: Enhancer of mRNA-decapping protein 3



• Molecule 1: Enhancer of mRNA-decapping protein 3





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	94.38Å 169.20Å 163.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 20.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.20) 99.8 (20.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.03 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.228 , 0.256 0.223 , 0.253	Depositor DCC
R_{free} test set	3376 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.000 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7556	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.9357e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1844	0.79	0/2507
1	B	0.54	0/1844	0.80	0/2507
1	C	0.56	0/1844	0.81	0/2507
1	D	0.54	0/1844	0.83	1/2507 (0.0%)
All	All	0.55	0/7376	0.81	1/10028 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	436	ARG	N-CA-C	5.38	122.26	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1802	0	1833	10	0
1	B	1802	0	1833	15	0
1	C	1802	0	1833	20	0
1	D	1802	0	1833	24	0
2	A	89	0	0	1	0
2	B	94	0	0	2	0
2	C	84	0	0	4	0
2	D	81	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7556	0	7332	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:371:PHE:HD2	1:D:371:PHE:H	1.07	1.01
1:C:371:PHE:H	1:C:371:PHE:HD2	1.00	0.98
1:C:434:GLN:HG2	2:C:573:HOH:O	1.67	0.93
1:C:371:PHE:HD2	1:C:371:PHE:N	1.68	0.91
1:D:371:PHE:HD2	1:D:371:PHE:N	1.79	0.80
1:C:371:PHE:N	1:C:371:PHE:CD2	2.42	0.79
1:D:422:GLN:HE21	1:D:424:TRP:HE1	1.33	0.76
1:D:334:ARG:HG2	1:D:361:ASP:O	1.85	0.75
1:B:334:ARG:HG2	1:B:361:ASP:O	1.88	0.73
1:A:334:ARG:HG2	1:A:361:ASP:O	1.88	0.72
1:D:434:GLN:HG3	2:D:571:HOH:O	1.92	0.69
1:D:422:GLN:NE2	1:D:424:TRP:HE1	1.95	0.63
1:C:274:ASP:OD1	1:C:506:HIS:HE1	1.82	0.62
1:B:370:ASN:ND2	2:B:512:HOH:O	2.32	0.61
1:D:274:ASP:OD1	1:D:506:HIS:HE1	1.84	0.60
1:D:371:PHE:N	1:D:371:PHE:CD2	2.52	0.59
1:B:355:ARG:HD3	1:B:384:LEU:HB3	1.85	0.59
1:C:371:PHE:HE1	2:C:547:HOH:O	1.87	0.58
1:C:334:ARG:HG2	1:C:361:ASP:O	2.03	0.58
1:B:459:LEU:HD11	1:B:477:CYS:HB2	1.88	0.55
1:D:422:GLN:HE21	1:D:424:TRP:NE1	2.01	0.55
1:D:459:LEU:HD11	1:D:477:CYS:HB2	1.89	0.55
1:D:413:CYS:HB3	1:D:416:ASN:HD22	1.72	0.54
1:C:274:ASP:OD1	1:C:506:HIS:CE1	2.61	0.53
1:A:355:ARG:HD3	1:A:384:LEU:HB3	1.90	0.53
1:B:370:ASN:OD1	1:D:422:GLN:NE2	2.42	0.52
1:C:413:CYS:HB3	1:C:416:ASN:HD22	1.74	0.52
1:B:307:MET:HE2	1:B:478:ASP:O	2.10	0.51
1:A:370:ASN:ND2	2:A:576:HOH:O	2.43	0.51
1:A:435:ASN:O	1:A:436:ARG:HB2	2.10	0.50
1:C:434:GLN:CG	2:C:573:HOH:O	2.40	0.49
1:D:343:PRO:HD2	1:D:419:LEU:HD21	1.95	0.49
1:B:274:ASP:OD1	1:B:506:HIS:HE1	1.96	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:459:LEU:HD11	1:C:477:CYS:HB2	1.96	0.48
1:D:274:ASP:OD1	1:D:506:HIS:CE1	2.66	0.48
1:A:497:PHE:HB3	1:A:500:LYS:O	2.14	0.48
1:D:416:ASN:O	1:D:419:LEU:HB2	2.13	0.48
1:D:260:ARG:HH21	1:D:494:HIS:CD2	2.32	0.47
1:D:497:PHE:HB3	1:D:500:LYS:O	2.14	0.47
1:B:419:LEU:HD11	1:B:424:TRP:HZ2	1.80	0.47
1:D:434:GLN:CG	2:D:571:HOH:O	2.57	0.47
1:B:370:ASN:HB2	2:B:512:HOH:O	2.14	0.47
1:C:307:MET:HE2	1:C:478:ASP:O	2.15	0.46
1:C:497:PHE:HB3	1:C:500:LYS:O	2.16	0.46
1:C:304:ARG:NH2	2:C:582:HOH:O	2.49	0.45
1:D:304:ARG:NH2	2:D:550:HOH:O	2.50	0.45
1:A:459:LEU:HD11	1:A:477:CYS:HB2	1.99	0.44
1:B:279:VAL:HG11	1:B:505:LEU:HD22	2.00	0.44
1:B:497:PHE:HB3	1:B:500:LYS:O	2.17	0.44
1:A:337:VAL:HG22	1:A:406:LEU:HB3	1.99	0.43
1:A:370:ASN:OD1	1:C:424:TRP:CD1	2.71	0.43
1:D:416:ASN:HB3	1:D:419:LEU:HD13	1.99	0.43
1:B:413:CYS:SG	1:B:414:PRO:HD2	2.57	0.43
1:C:355:ARG:HD3	1:C:384:LEU:HB3	2.00	0.43
1:D:407:VAL:HG23	1:D:437:ALA:HB3	2.02	0.42
1:B:337:VAL:HG22	1:B:406:LEU:HB3	2.02	0.42
1:C:371:PHE:CZ	1:C:374:MET:HG2	2.55	0.42
1:C:407:VAL:HG23	1:C:437:ALA:HB3	2.02	0.42
1:B:393:VAL:HG11	1:B:398:ASP:HB3	2.02	0.42
1:A:413:CYS:SG	1:A:414:PRO:HD2	2.60	0.41
1:C:344:HIS:CG	1:C:413:CYS:HB2	2.55	0.41
1:D:344:HIS:CG	1:D:413:CYS:HB2	2.56	0.40
1:A:360:HIS:HE1	1:B:500:LYS:O	2.04	0.40
1:C:343:PRO:HD2	1:C:419:LEU:HD21	2.03	0.40
1:D:355:ARG:HD3	1:D:384:LEU:HB3	2.04	0.40
1:D:339:LEU:HD12	1:D:341:CYS:SG	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/259 (87%)	220 (98%)	5 (2%)	0	100	100
1	B	225/259 (87%)	222 (99%)	3 (1%)	0	100	100
1	C	225/259 (87%)	220 (98%)	4 (2%)	1 (0%)	30	34
1	D	225/259 (87%)	220 (98%)	4 (2%)	1 (0%)	30	34
All	All	900/1036 (87%)	882 (98%)	16 (2%)	2 (0%)	43	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	322	GLY
1	D	322	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/227 (89%)	197 (97%)	6 (3%)	36	49
1	B	203/227 (89%)	197 (97%)	6 (3%)	36	49
1	C	203/227 (89%)	194 (96%)	9 (4%)	25	34
1	D	203/227 (89%)	192 (95%)	11 (5%)	20	25
All	All	812/908 (89%)	780 (96%)	32 (4%)	28	39

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	TYR
1	A	279	VAL
1	A	296	LYS
1	A	339	LEU
1	A	370	ASN
1	A	373	LYS
1	B	258	TYR
1	B	279	VAL
1	B	296	LYS
1	B	339	LEU
1	B	366	LEU
1	B	373	LYS
1	C	258	TYR
1	C	279	VAL
1	C	296	LYS
1	C	339	LEU
1	C	371	PHE
1	C	373	LYS
1	C	374	MET
1	C	401	THR
1	C	419	LEU
1	D	258	TYR
1	D	279	VAL
1	D	296	LYS
1	D	302	GLU
1	D	339	LEU
1	D	371	PHE
1	D	373	LYS
1	D	374	MET
1	D	375	LEU
1	D	419	LEU
1	D	436	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	422	GLN
1	A	487	GLN
1	A	506	HIS
1	B	333	GLN
1	B	349	GLN
1	B	416	ASN
1	B	506	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	360	HIS
1	C	416	ASN
1	C	422	GLN
1	C	506	HIS
1	D	360	HIS
1	D	416	ASN
1	D	422	GLN
1	D	506	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/259 (89%)	0.33	16 (6%) 23 20	19, 27, 41, 50	0
1	B	233/259 (89%)	0.28	15 (6%) 25 22	17, 27, 43, 53	0
1	C	233/259 (89%)	0.17	10 (4%) 40 36	18, 27, 38, 49	0
1	D	233/259 (89%)	0.24	13 (5%) 30 27	18, 26, 41, 50	0
All	All	932/1036 (89%)	0.26	54 (5%) 29 25	17, 27, 41, 53	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	323	PRO	6.1
1	D	371	PHE	5.7
1	C	322	GLY	5.3
1	B	258	TYR	5.1
1	C	323	PRO	5.1
1	A	322	GLY	4.8
1	B	322	GLY	4.7
1	D	323	PRO	4.5
1	A	258	TYR	4.4
1	B	323	PRO	4.4
1	C	371	PHE	4.3
1	D	418	PHE	4.2
1	B	447	HIS	4.0
1	C	418	PHE	3.9
1	B	418	PHE	3.8
1	B	416	ASN	3.7
1	A	333	GLN	3.5
1	C	258	TYR	3.5
1	B	268	SER	3.3
1	B	333	GLN	3.3
1	C	321	GLY	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	436	ARG	3.2
1	A	447	HIS	3.2
1	A	416	ASN	3.2
1	C	419	LEU	3.1
1	A	417	VAL	3.1
1	A	446	VAL	3.1
1	C	434	GLN	3.0
1	A	436	ARG	3.0
1	B	302	GLU	2.9
1	B	434	GLN	2.9
1	D	401	THR	2.9
1	D	434	GLN	2.8
1	D	258	TYR	2.8
1	A	418	PHE	2.8
1	C	447	HIS	2.7
1	D	322	GLY	2.6
1	A	268	SER	2.6
1	B	415	GLU	2.6
1	A	434	GLN	2.5
1	D	447	HIS	2.4
1	B	446	VAL	2.4
1	A	370	ASN	2.4
1	D	268	SER	2.4
1	A	260	ARG	2.4
1	A	415	GLU	2.4
1	D	419	LEU	2.3
1	A	321	GLY	2.3
1	D	334	ARG	2.3
1	B	420	ARG	2.2
1	D	333	GLN	2.1
1	B	507	SER	2.1
1	C	436	ARG	2.1
1	B	453	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.