



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 10:26 PM EDT

PDB ID : 6D3M / pdb_00006d3m
Title : FT_T dioxygenase with bound quizalofop
Authors : Rydel, T.J.; Halls, C.E.
Deposited on : 2018-04-16
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	
1	B	295	
1	F	295	
1	J	295	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FT_T dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	2	0
			2300	1461	396	432	11			
1	B	285	Total	C	N	O	S	0	3	0
			2295	1460	392	432	11			
1	F	286	Total	C	N	O	S	0	1	0
			2292	1457	394	430	11			
1	J	286	Total	C	N	O	S	0	1	0
			2292	1457	394	430	11			

- Molecule 2 is COBALT (II) ION (CCD ID: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Co	0	0
			1	1		
2	B	1	Total	Co	0	0
			1	1		
2	F	1	Total	Co	0	0
			1	1		
2	J	1	Total	Co	0	0
			1	1		

- Molecule 3 is (2R)-2-{4-[(6-chloroquinoxalin-2-yl)oxy]phenoxy}propanoic acid (CCD ID: FTJ) (formula: C₁₇H₁₃ClN₂O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			24	17	1	2	4		
3	B	1	Total	C	Cl	N	O	0	0
			24	17	1	2	4		
3	F	1	Total	C	Cl	N	O	0	0
			24	17	1	2	4		
3	J	1	Total	C	Cl	N	O	0	0
			24	17	1	2	4		

- Molecule 4 is 2-OXOGLUTARIC ACID (CCD ID: AKG) (formula: $C_5H_6O_5$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 10 5 5	0	0
4	B	1	Total C O 10 5 5	0	0
4	F	1	Total C O 10 5 5	0	0
4	J	1	Total C O 10 5 5	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0
5	F	1	Total Cl 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	206	Total O 206 206	0	0
6	B	177	Total O 177 177	0	0
6	F	186	Total O 186 186	0	0
6	J	190	Total O 190 190	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FT_T dioxygenase

Chain A: 




- Molecule 1: FT_T dioxygenase

Chain B: 



- Molecule 1: FT_T dioxygenase

Chain F: 



- Molecule 1: FT_T dioxygenase

Chain J: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	137.78Å 137.78Å 148.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.10 – 2.03 45.10 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.2 (45.10-2.03) 99.4 (45.10-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.177 , 0.217 0.180 , 0.219	Depositor DCC
R_{free} test set	5165 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.469 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	10080	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0964e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FTJ, AKG, CL, CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.25	4/2363 (0.2%)	1.16	9/3217 (0.3%)
1	B	1.27	3/2364 (0.1%)	1.16	3/3218 (0.1%)
1	F	1.27	2/2355 (0.1%)	1.16	4/3206 (0.1%)
1	J	1.26	3/2355 (0.1%)	1.16	4/3206 (0.1%)
All	All	1.26	12/9437 (0.1%)	1.16	20/12847 (0.2%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	122	ALA	CA-CB	8.44	1.67	1.53
1	F	156	ALA	CA-CB	6.72	1.63	1.53
1	J	125	VAL	CA-CB	6.28	1.61	1.54
1	B	123	ALA	CA-CB	6.15	1.64	1.53
1	B	125	VAL	CA-CB	6.13	1.61	1.53
1	A	86	ILE	CA-CB	-5.77	1.48	1.54
1	J	119	ALA	CA-CB	5.48	1.60	1.53
1	A	122	ALA	CA-CB	5.44	1.62	1.53
1	B	285	ARG	N-CA	-5.37	1.39	1.45
1	A	163	VAL	C-O	5.36	1.29	1.24
1	J	288	VAL	CA-CB	5.25	1.60	1.54
1	A	156	ALA	CA-CB	5.20	1.61	1.53

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	86	ILE	CB-CA-C	-7.16	101.41	111.28
1	F	288	VAL	CB-CA-C	7.14	120.55	110.84
1	A	35	GLU	CA-C-N	-7.07	112.70	119.85
1	A	35	GLU	C-N-CA	-7.07	112.70	119.85
1	A	273	VAL	CA-C-N	-6.21	115.38	121.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	273	VAL	C-N-CA	-6.21	115.38	121.65
1	A	86	ILE	CB-CA-C	-5.87	103.17	111.28
1	F	125	VAL	CB-CA-C	5.79	118.51	110.99
1	J	131	VAL	CA-C-N	5.70	125.65	119.78
1	J	131	VAL	C-N-CA	5.70	125.65	119.78
1	F	179	ASN	N-CA-C	5.63	120.28	112.45
1	A	125	VAL	CG1-CB-CG2	5.54	123.00	110.80
1	A	212	ARG	CB-CG-CD	-5.32	99.07	111.30
1	A	288	VAL	CB-CA-C	5.27	118.68	110.83
1	B	235	SER	N-CA-C	5.16	116.59	111.07
1	J	186	SER	N-CA-C	-5.14	108.03	114.56
1	J	125	VAL	CG1-CB-CG2	5.13	122.09	110.80
1	A	125	VAL	CB-CA-C	5.07	117.96	110.77
1	F	17	VAL	CB-CA-C	5.07	118.38	110.83
1	B	125	VAL	CB-CA-C	5.01	117.50	110.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2300	0	2229	13	0
1	B	2295	0	2228	13	0
1	F	2292	0	2224	23	0
1	J	2292	0	2224	16	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	F	1	0	0	0	0
2	J	1	0	0	0	0
3	A	24	0	0	0	0
3	B	24	0	0	0	0
3	F	24	0	0	0	0
3	J	24	0	0	0	0
4	A	10	0	4	0	0
4	B	10	0	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	10	0	4	0	0
4	J	10	0	4	0	0
5	A	1	0	0	0	0
5	F	1	0	0	0	0
6	A	206	0	0	3	0
6	B	177	0	0	2	0
6	F	186	0	0	5	0
6	J	190	0	0	4	0
All	All	10080	0	8921	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:GLU:HG2	6:B:478:HOH:O	1.36	1.21
1:J:91:GLU:HG2	6:J:498:HOH:O	1.52	1.07
1:B:181:ARG:HG2	1:B:181:ARG:HH11	1.22	1.02
1:F:181:ARG:HG2	1:F:181:ARG:HH11	1.29	0.97
1:F:127:ARG:HD2	6:F:401:HOH:O	1.65	0.96
1:F:86:ILE:HD11	1:F:282:TYR:OH	1.68	0.93
1:F:10:ASN:HD22	1:F:11:LYS:H	1.16	0.93
1:A:181:ARG:HH11	1:A:181:ARG:HG2	1.33	0.92
1:F:86:ILE:HD11	1:F:282:TYR:HH	1.37	0.88
1:J:181:ARG:HG2	1:J:181:ARG:HH11	1.40	0.86
1:F:91:GLU:HG2	6:F:483:HOH:O	1.82	0.79
1:A:16:ASP:HB3	1:A:18:GLN:HE22	1.47	0.78
1:B:181:ARG:HH11	1:B:181:ARG:CG	1.98	0.76
1:F:181:ARG:HH11	1:F:181:ARG:CG	1.99	0.75
1:F:16:ASP:HB3	1:F:18:GLN:HE22	1.51	0.74
1:A:181:ARG:HH11	1:A:181:ARG:CG	2.03	0.72
1:J:181:ARG:HH11	1:J:181:ARG:CG	2.02	0.72
1:B:181:ARG:HG2	1:B:181:ARG:NH1	2.01	0.69
1:F:10:ASN:ND2	1:F:11:LYS:H	1.90	0.69
1:B:16:ASP:HB3	1:B:18:GLN:HE22	1.61	0.65
1:J:181:ARG:HG2	1:J:181:ARG:NH1	2.12	0.65
1:J:16:ASP:HB3	1:J:18:GLN:HE22	1.62	0.64
1:B:96:ARG:HH22	1:F:13:ARG:HH21	1.48	0.61
1:F:181:ARG:HG2	1:F:181:ARG:NH1	2.07	0.60
1:F:86:ILE:CD1	1:F:282:TYR:OH	2.48	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:GLU:HG2	6:A:490:HOH:O	2.06	0.55
1:A:174:LEU:O	1:A:178:THR:HG23	2.06	0.55
1:F:192:VAL:HG23	6:F:524:HOH:O	2.07	0.55
1:A:147:TRP:CZ3	1:A:204:VAL:HG21	2.43	0.54
1:F:234:LYS:O	1:F:238:GLN:HG2	2.08	0.53
1:F:184:ASN:HB2	6:F:522:HOH:O	2.09	0.52
1:A:181:ARG:HG2	1:A:181:ARG:NH1	2.11	0.51
1:F:141:LEU:C	1:F:141:LEU:HD12	2.36	0.51
1:F:181:ARG:CG	1:F:181:ARG:NH1	2.68	0.50
1:A:181:ARG:CG	1:A:181:ARG:NH1	2.71	0.49
1:B:50:HIS:CE1	1:B:209:VAL:HG11	2.47	0.49
1:J:86:ILE:HG13	1:J:89:TYR:HD1	1.77	0.49
1:A:86:ILE:HG22	1:A:87:GLU:N	2.29	0.47
1:J:82:ILE:HG12	1:J:187:VAL:HG13	1.96	0.47
1:J:147:TRP:CZ3	1:J:204:VAL:HG21	2.49	0.47
1:A:141:LEU:C	1:A:141:LEU:HD12	2.40	0.47
1:B:109:ASP:HB2	6:B:408:HOH:O	2.14	0.47
1:A:87:GLU:HG3	6:A:423:HOH:O	2.16	0.46
1:B:181:ARG:CG	1:B:181:ARG:NH1	2.67	0.46
1:J:178:THR:HB	6:J:425:HOH:O	2.16	0.44
1:B:11:LYS:HG3	1:B:12:TYR:N	2.29	0.44
1:J:109:ASP:HB2	6:J:424:HOH:O	2.18	0.43
1:J:178:THR:HG22	1:J:179:ASN:H	1.82	0.43
1:F:79:PRO:HD3	1:F:91:GLU:OE1	2.18	0.43
1:J:115:THR:HA	1:J:120:PRO:HG3	2.01	0.43
1:B:276:TYR:O	1:B:277:ALA:C	2.63	0.42
1:F:125:VAL:HB	1:F:262:VAL:HG22	2.01	0.42
1:J:98:GLU:HB2	1:J:101:GLU:HG3	2.00	0.42
1:F:35:GLU:HB3	1:F:36:PRO:HD2	2.02	0.41
1:J:115:THR:HA	1:J:120:PRO:CG	2.48	0.41
1:A:234:LYS:O	1:A:238:GLN:HG2	2.20	0.41
1:J:104:ARG:HG2	6:J:423:HOH:O	2.20	0.41
1:F:110:TRP:CE2	1:F:271:ARG:HD2	2.55	0.41
1:A:100[A]:ASN:CG	6:A:479:HOH:O	2.63	0.41
1:B:125:VAL:HB	1:B:262:VAL:HG22	2.02	0.41
1:F:121:PRO:HB3	1:F:287:THR:CG2	2.50	0.40
1:F:127:ARG:CD	6:F:401:HOH:O	2.43	0.40
1:J:276:TYR:O	1:J:277:ALA:C	2.63	0.40
1:B:110:TRP:CE2	1:B:271:ARG:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/295 (97%)	278 (97%)	8 (3%)	0	100	100
1	B	286/295 (97%)	280 (98%)	6 (2%)	0	100	100
1	F	285/295 (97%)	280 (98%)	5 (2%)	0	100	100
1	J	285/295 (97%)	278 (98%)	7 (2%)	0	100	100
All	All	1142/1180 (97%)	1116 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/256 (98%)	243 (97%)	8 (3%)	34	30
1	B	251/256 (98%)	242 (96%)	9 (4%)	31	26
1	F	250/256 (98%)	242 (97%)	8 (3%)	34	30
1	J	250/256 (98%)	240 (96%)	10 (4%)	28	22
All	All	1002/1024 (98%)	967 (96%)	35 (4%)	32	27

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	11	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	17	VAL
1	A	104	ARG
1	A	125	VAL
1	A	181	ARG
1	A	248	ASP
1	A	288	VAL
1	B	11	LYS
1	B	17	VAL
1	B	73	ARG
1	B	125	VAL
1	B	179	ASN
1	B	181	ARG
1	B	288	VAL
1	B	294	SER
1	B	295	ARG
1	F	10	ASN
1	F	13	ARG
1	F	17	VAL
1	F	73	ARG
1	F	125	VAL
1	F	179	ASN
1	F	181	ARG
1	F	288	VAL
1	J	11	LYS
1	J	17	VAL
1	J	91	GLU
1	J	125	VAL
1	J	178	THR
1	J	181	ARG
1	J	235	SER
1	J	248	ASP
1	J	288	VAL
1	J	295	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	50	HIS
1	A	176	GLN
1	B	18	GLN
1	B	50	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	64	ASN
1	B	219	GLN
1	B	226	GLN
1	F	10	ASN
1	F	18	GLN
1	F	50	HIS
1	F	64	ASN
1	F	226	GLN
1	J	18	GLN
1	J	50	HIS
1	J	64	ASN
1	J	176	GLN
1	J	219	GLN
1	J	238	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 6 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	AKG	A	303	2	9,9,9	1.95	1 (11%)	11,11,11	1.86	4 (36%)
3	FTJ	F	302	-	26,26,26	1.23	1 (3%)	36,36,36	2.38	9 (25%)
3	FTJ	J	302	-	26,26,26	1.36	3 (11%)	36,36,36	1.90	7 (19%)
4	AKG	F	303	2	9,9,9	2.05	3 (33%)	11,11,11	1.75	3 (27%)
3	FTJ	B	302	-	26,26,26	1.14	1 (3%)	36,36,36	2.00	7 (19%)
4	AKG	B	303	2	9,9,9	1.87	2 (22%)	11,11,11	1.90	4 (36%)
4	AKG	J	303	2	9,9,9	1.64	1 (11%)	11,11,11	2.36	6 (54%)
3	FTJ	A	302	-	26,26,26	1.40	3 (11%)	36,36,36	1.89	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AKG	A	303	2	-	1/9/9/9	-
3	FTJ	F	302	-	-	8/12/12/12	0/3/3/3
3	FTJ	J	302	-	-	8/12/12/12	0/3/3/3
4	AKG	F	303	2	-	2/9/9/9	-
3	FTJ	B	302	-	-	8/12/12/12	0/3/3/3
4	AKG	B	303	2	-	1/9/9/9	-
4	AKG	J	303	2	-	2/9/9/9	-
3	FTJ	A	302	-	-	8/12/12/12	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	303	AKG	C2-C1	-4.70	1.46	1.53
4	B	303	AKG	C2-C1	-4.64	1.46	1.53
4	F	303	AKG	C2-C1	-4.60	1.46	1.53
4	J	303	AKG	C2-C1	-4.05	1.47	1.53
3	A	302	FTJ	O1-C7	-2.87	1.33	1.36
4	F	303	AKG	C3-C2	2.76	1.54	1.51
3	J	302	FTJ	O2-C15	2.60	1.47	1.43
3	J	302	FTJ	O3-C16	2.52	1.29	1.22
3	A	302	FTJ	C3-CL1	2.49	1.80	1.74
3	J	302	FTJ	C3-CL1	2.43	1.80	1.74
3	F	302	FTJ	C3-CL1	2.36	1.79	1.74
3	A	302	FTJ	O3-C16	2.27	1.28	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	303	AKG	O3-C5	2.15	1.29	1.22
4	F	303	AKG	O3-C5	2.15	1.29	1.22
3	B	302	FTJ	O3-C16	2.10	1.28	1.22

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	302	FTJ	C7-C8-N2	-10.11	116.17	121.89
3	A	302	FTJ	C7-C8-N2	-8.48	117.09	121.89
3	B	302	FTJ	C7-C8-N2	-8.24	117.23	121.89
3	J	302	FTJ	C7-C8-N2	-7.42	117.69	121.89
3	B	302	FTJ	C9-O1-C7	4.13	130.10	119.06
3	F	302	FTJ	C9-O1-C7	3.97	129.67	119.06
3	A	302	FTJ	C9-O1-C7	3.89	129.45	119.06
3	F	302	FTJ	C8-N2-C1	3.83	121.40	116.96
4	J	303	AKG	C3-C2-C1	3.83	122.37	115.86
3	J	302	FTJ	C9-O1-C7	3.67	128.88	119.06
4	J	303	AKG	O1-C1-C2	-3.57	117.37	121.81
3	B	302	FTJ	C7-N1-C6	3.47	118.44	115.85
4	B	303	AKG	O5-C2-C1	-3.45	114.74	119.67
4	A	303	AKG	O1-C1-C2	-3.40	117.58	121.81
4	F	303	AKG	O1-C1-C2	-3.35	117.64	121.81
3	J	302	FTJ	C8-N2-C1	3.21	120.67	116.96
4	B	303	AKG	C3-C2-C1	3.18	121.26	115.86
3	F	302	FTJ	C12-O2-C15	-3.11	113.33	118.40
4	J	303	AKG	O2-C1-C2	3.05	122.06	113.59
3	J	302	FTJ	C7-N1-C6	3.03	118.12	115.85
3	B	302	FTJ	O2-C15-C17	2.99	113.20	106.97
3	F	302	FTJ	C14-C9-C10	2.95	124.46	120.16
3	B	302	FTJ	C8-N2-C1	2.94	120.36	116.96
3	J	302	FTJ	O2-C15-C17	2.89	112.98	106.97
3	F	302	FTJ	C7-N1-C6	2.89	118.01	115.85
3	F	302	FTJ	C13-C14-C9	-2.88	116.45	119.73
4	F	303	AKG	O5-C2-C1	-2.82	115.64	119.67
3	B	302	FTJ	C17-C15-C16	2.80	114.68	109.42
3	A	302	FTJ	C8-N2-C1	2.72	120.10	116.96
4	J	303	AKG	O5-C2-C1	-2.67	115.85	119.67
4	F	303	AKG	C3-C2-C1	2.64	120.34	115.86
3	J	302	FTJ	C2-C1-N2	2.61	121.01	118.01
4	B	303	AKG	O1-C1-C2	-2.56	118.63	121.81
3	A	302	FTJ	C14-C9-C10	2.54	123.87	120.16
4	A	303	AKG	C3-C2-C1	2.51	120.12	115.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	302	FTJ	C17-C15-C16	2.50	114.11	109.42
4	B	303	AKG	O2-C1-C2	2.49	120.50	113.59
3	F	302	FTJ	O2-C15-C17	2.45	112.07	106.97
4	J	303	AKG	O3-C5-C4	-2.44	115.35	123.09
4	A	303	AKG	O3-C5-C4	-2.41	115.44	123.09
4	J	303	AKG	O4-C5-C4	2.30	121.27	114.00
3	F	302	FTJ	O4-C16-C15	2.30	122.34	113.96
4	A	303	AKG	O2-C1-C2	2.25	119.83	113.59
3	A	302	FTJ	O2-C15-C17	2.17	111.49	106.97
3	B	302	FTJ	C2-C1-N2	2.02	120.33	118.01

There are no chirality outliers.

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	FTJ	O2-C15-C16-O4
3	B	302	FTJ	O2-C15-C16-O4
3	B	302	FTJ	O2-C15-C16-O3
3	B	302	FTJ	C17-C15-C16-O4
3	B	302	FTJ	C17-C15-C16-O3
3	F	302	FTJ	O2-C15-C16-O4
3	F	302	FTJ	O2-C15-C16-O3
3	F	302	FTJ	C17-C15-C16-O4
3	F	302	FTJ	C17-C15-C16-O3
3	J	302	FTJ	O2-C15-C16-O4
4	A	303	AKG	C1-C2-C3-C4
4	B	303	AKG	C1-C2-C3-C4
4	F	303	AKG	C1-C2-C3-C4
3	A	302	FTJ	C17-C15-C16-O4
3	A	302	FTJ	C17-C15-C16-O3
3	J	302	FTJ	C17-C15-C16-O4
3	J	302	FTJ	C17-C15-C16-O3
3	A	302	FTJ	O2-C15-C16-O3
3	J	302	FTJ	O2-C15-C16-O3
3	A	302	FTJ	C16-C15-O2-C12
3	F	302	FTJ	C16-C15-O2-C12
3	J	302	FTJ	C16-C15-O2-C12
3	A	302	FTJ	C17-C15-O2-C12
3	B	302	FTJ	C17-C15-O2-C12
3	F	302	FTJ	C17-C15-O2-C12
3	J	302	FTJ	C17-C15-O2-C12
3	J	302	FTJ	C11-C12-O2-C15

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	302	FTJ	C11-C12-O2-C15
3	B	302	FTJ	C16-C15-O2-C12
4	J	303	AKG	C1-C2-C3-C4
3	J	302	FTJ	C13-C12-O2-C15
3	A	302	FTJ	C11-C12-O2-C15
3	B	302	FTJ	C13-C12-O2-C15
3	A	302	FTJ	C13-C12-O2-C15
3	F	302	FTJ	C11-C12-O2-C15
3	F	302	FTJ	C13-C12-O2-C15
4	J	303	AKG	C3-C4-C5-O4
4	F	303	AKG	O5-C2-C3-C4


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand FTJ J 302



Ligand FTJ B 302





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/295 (96%)	-1.26	0 100 100	20, 35, 49, 77	2 (0%)
1	B	285/295 (96%)	-1.24	0 100 100	20, 35, 51, 78	3 (1%)
1	F	286/295 (96%)	-1.24	0 100 100	27, 35, 49, 77	1 (0%)
1	J	286/295 (96%)	-1.25	0 100 100	28, 35, 51, 78	1 (0%)
All	All	1143/1180 (96%)	-1.25	0 100 100	20, 35, 51, 78	7 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FTJ	A	302	24/24	0.98	0.05	38,51,72,87	0
3	FTJ	B	302	24/24	0.98	0.06	39,58,84,93	0
3	FTJ	F	302	24/24	0.98	0.05	37,48,69,84	0
3	FTJ	J	302	24/24	0.98	0.05	42,58,79,90	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	AKG	B	303	10/10	0.99	0.03	33,34,38,39	0
4	AKG	F	303	10/10	0.99	0.02	28,32,34,34	0
4	AKG	J	303	10/10	0.99	0.03	34,37,40,40	0
2	CO	J	301	1/1	1.00	0.01	32,32,32,32	0
4	AKG	A	303	10/10	1.00	0.02	29,32,35,36	0
2	CO	A	301	1/1	1.00	0.01	30,30,30,30	0
2	CO	B	301	1/1	1.00	0.02	32,32,32,32	0
2	CO	F	301	1/1	1.00	0.01	30,30,30,30	0
5	CL	A	304	1/1	1.00	0.03	35,35,35,35	0
5	CL	F	304	1/1	1.00	0.03	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FTJ A 302:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FTJ B 302:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FTJ F 302:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FTJ J 302:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.