



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:09 AM UTC

PDB ID : 8D32 / pdb_00008d32
Title : Mycobacterium tuberculosis pduO-type ATP:cobalamin adenosyltransferase bound to 5-deoxyadenosylrhodibalamin and PPPi
Authors : Mascarenhas, R.N.; Ruetz, M.; Koutmos, M.; Banerjee, R.
Deposited on : 2022-05-31
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	19% 96% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SF0	A	201	X	-	-	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 3027 atoms, of which 1469 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Corrinoid adenosyltransferase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	186	2783	879	1381	254	266	3	0	3	0

- Molecule 2 is [5-(5,6-dimethyl-1H-benzimidazol-1-yl)-4-hydroxy-2-(hydroxymethyl)oxolan-3-yl 1-{3-[3,14,19-tris(2-amino-2-oxoethyl)-8,13,18-tris(3-amino-3-oxopropyl)-1,4,6,9,9,14,16,19-octamethyl-20,21,22,23-tetraazapentacyclo[15.2.1.1 2,5 .1 7,10 .1 12,15]tricoso-5(23),6,10,12(21),15,17(20)-hexaen-4-yl-kappa 4 N 20 ,N 21 ,N 22 ,N 23]propanamido}propan-2-yl hydrogenato phosphate]rhodium (non-preferred name) (CCD ID: SF0) (formula: C₆₂H₈₉N₁₃O₁₄PRh) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	C	H	N	O	P	Rh		
2	A	1	179	62	88	13	14	1	1	0	0

- Molecule 3 is 5'-DEOXYADENOSINE (CCD ID: 5AD) (formula: C₁₀H₁₃N₅O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			18	10	5	3		

- Molecule 4 is TRIPHOSPHATE (CCD ID: 3PO) (formula: $\text{H}_5\text{O}_{10}\text{P}_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			13	10	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	32	Total	O	0	0
			32	32		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Corrinoid adenosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	62.62Å 62.62Å 267.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.05 – 1.85 38.05 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.05-1.85) 99.6 (38.05-1.85)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.85Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.218 , 0.246 (Not available) , 0.270	Depositor DCC
R_{free} test set	898 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 29.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3027	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5AD, MG, 3PO, SF0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.18	0/1446	0.33	0/1979

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1402	1381	1370	0	0
2	A	91	88	0	0	0
3	A	18	0	11	1	0
4	A	13	0	0	1	0
5	A	2	0	0	0	0
6	A	32	0	0	0	0
All	All	1558	1469	1381	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:202:5AD:H5'2	4:A:203:3PO:O1G	1.79	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/193 (97%)	183 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/152 (97%)	147 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	SF0	A	201	-	92,101,101	0.88	3 (3%)	146,166,166	1.00	8 (5%)
3	5AD	A	202	-	20,20,20	1.53	4 (20%)	28,30,30	1.62	7 (25%)
4	3PO	A	203	5	10,12,12	2.73	1 (10%)	17,20,20	1.96	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF0	A	201	-	1/1/38/38	16/56/223/223	0/3/11/11
3	5AD	A	202	-	-	3/4/20/20	0/3/3/3
4	3PO	A	203	5	-	2/12/12/12	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	203	3PO	PB-O3A	-8.14	1.50	1.59
2	A	201	SF0	RH01-N19	-4.96	2.00	2.04
3	A	202	5AD	O2'-C2'	-3.25	1.34	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	SF0	C90-C91	2.97	1.59	1.51
3	A	202	5AD	O3'-C3'	-2.86	1.35	1.43
2	A	201	SF0	C91-N29	-2.74	1.23	1.32
3	A	202	5AD	C4-N9	2.18	1.42	1.37
3	A	202	5AD	C8-N9	2.11	1.41	1.37

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	SF0	C68-N22-C64	4.33	111.90	108.09
3	A	202	5AD	N3-C2-N1	3.81	134.35	128.58
3	A	202	5AD	C5'-C4'-C3'	-3.69	111.83	115.70
4	A	203	3PO	O2G-PG-O3B	3.53	116.47	104.64
2	A	201	SF0	C52-C50-N18	3.14	106.70	105.30
2	A	201	SF0	C64-C63-C62	2.76	125.47	121.26
3	A	202	5AD	C4-N9-C8	-2.76	102.84	105.74
2	A	201	SF0	C68-C32-N17	2.71	105.12	102.19
3	A	202	5AD	C2-N1-C6	-2.68	114.33	118.73
4	A	203	3PO	O3G-PG-O2G	2.62	117.63	107.80
2	A	201	SF0	C83-C63-C64	-2.47	116.17	120.36
2	A	201	SF0	O08-C30-N18	2.42	112.74	108.09
3	A	202	5AD	C1'-N9-C8	2.34	132.29	127.09
4	A	203	3PO	O3A-PA-O1A	-2.32	98.84	111.04
4	A	203	3PO	O5'-PA-O2A	2.24	116.21	107.80
3	A	202	5AD	N9-C8-N7	2.24	117.11	113.94
4	A	203	3PO	O3B-PG-O1G	-2.19	99.53	111.04
2	A	201	SF0	C75-C48-C46	2.19	113.88	107.11
4	A	203	3PO	O3G-PG-O1G	-2.08	102.74	110.83
3	A	202	5AD	C2-N3-C4	-2.05	106.84	111.83
2	A	201	SF0	C66-C65-C67	2.01	115.88	112.99

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	201	SF0	N20

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	SF0	C87-C88-O04-P02
2	A	201	SF0	C89-C88-O04-P02
2	A	201	SF0	C37-O03-P02-O04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	201	SF0	O08-C41-C44-O09
2	A	201	SF0	C37-C41-C44-O09
2	A	201	SF0	O15-C86-N28-C87
2	A	201	SF0	C77-C78-C79-O13
2	A	201	SF0	N28-C87-C88-C89
2	A	201	SF0	C85-C86-N28-C87
3	A	202	5AD	C2'-C1'-N9-C8
4	A	203	3PO	PA-O3A-PB-O2B
2	A	201	SF0	C80-C81-C82-O14
2	A	201	SF0	C77-C78-C79-N26
2	A	201	SF0	O08-C30-N18-C50
2	A	201	SF0	C80-C81-C82-N27
3	A	202	5AD	C2'-C1'-N9-C4
2	A	201	SF0	C34-C30-N18-C50
3	A	202	5AD	O4'-C1'-N9-C8
2	A	201	SF0	C37-O03-P02-O05
2	A	201	SF0	C37-O03-P02-O06
4	A	203	3PO	PB-O3B-PG-O1G


There are no ring outliers.

2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	5AD	1	0
4	A	203	3PO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand SF0 A 201



Ligand 5AD A 202





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	186/193 (96%)	0.88	36 (19%) 3 2	19, 39, 58, 65	1 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	181	ASP	4.5
1	A	179	ASP	4.4
1	A	182	VAL	3.7
1	A	175	VAL	3.7
1	A	4	HIS	3.5
1	A	180	GLY	3.5
1	A	86	LYS	3.3
1	A	3	VAL	3.2
1	A	110	GLY	3.1
1	A	176	ALA	3.0
1	A	188	GLY	3.0
1	A	109	ALA	3.0
1	A	178	PRO	3.0
1	A	54	HIS	2.9
1	A	123	SER	2.9
1	A	57	THR	2.8
1	A	118	VAL	2.7
1	A	122	GLY	2.6
1	A	112	PRO	2.5
1	A	8	ILE	2.5
1	A	177	ASN	2.5
1	A	64	ARG	2.5
1	A	185	ARG	2.4
1	A	106	ALA	2.3
1	A	113	ALA	2.3
1	A	83	GLU	2.2
1	A	59	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	117	PHE	2.2
1	A	121	GLY	2.2
1	A	58	GLN	2.2
1	A	51	ALA	2.2
1	A	103	TRP	2.1
1	A	111	LEU	2.1
1	A	60	THR	2.0
1	A	84	ASN	2.0
1	A	107	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	A	205	1/1	0.86	0.13	24,24,24,24	0
3	5AD	A	202	18/18	0.91	0.09	21,26,29,29	0
4	3PO	A	203	13/13	0.95	0.09	27,28,35,35	0
2	SF0	A	201	91/91	0.96	0.11	26,38,57,65	0
5	MG	A	204	1/1	0.97	0.09	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 5AD A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3PO A 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SF0 A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.