



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 12:17 PM EDT

PDB ID : 8D3J / pdb_00008d3j
Title : Crystal structure of human Apoptosis-Inducing Factor (AIF) complexed with 6-fluoro-2-methylquinolin-4-amine
Authors : Brosey, C.A.; Tainer, J.A.
Deposited on : 2022-06-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	5% 78% • 20%
1	B	543	6% 78% • 20%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13224 atoms, of which 6404 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Apoptosis-inducing factor 1, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	436	Total	C	H	N	O	S	0	1	0
			6433	2084	3174	569	595	11			
1	B	434	Total	C	H	N	O	S	0	1	0
			6374	2060	3146	569	588	11			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	77	MET	-	initiating methionine	UNP O95831
A	614	LEU	-	expression tag	UNP O95831
A	615	GLU	-	expression tag	UNP O95831
A	616	VAL	-	expression tag	UNP O95831
A	617	LEU	-	expression tag	UNP O95831
A	618	PHE	-	expression tag	UNP O95831
A	619	GLN	-	expression tag	UNP O95831
B	77	MET	-	initiating methionine	UNP O95831
B	614	LEU	-	expression tag	UNP O95831
B	615	GLU	-	expression tag	UNP O95831
B	616	VAL	-	expression tag	UNP O95831
B	617	LEU	-	expression tag	UNP O95831
B	618	PHE	-	expression tag	UNP O95831
B	619	GLN	-	expression tag	UNP O95831

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	10	2	6	2	0	0


- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
3	A	1	83	27	30	9	15	2	0	0
3	B	1	83	27	30	9	15	2	0	0


- Molecule 4 is 6-fluoro-2-methylquinolin-4-amine (CCD ID: QEF) (formula: $C_{10}H_9FN_2$)

(labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	H	N	0
			22	10	1	9	2	
4	B	1	Total	C	F	H	N	0
			22	10	1	9	2	

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O S	0	0
			5	4 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	90	Total	O	0	0
			90	90		
6	B	82	Total	O	0	0
			82	82		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.47Å 115.32Å 123.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.44 – 2.40 84.44 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (84.44-2.40) 99.8 (84.44-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.42 (at 2.40Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.192 , 0.214 0.201 , 0.223	Depositor DCC
R_{free} test set	2637 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	44.4	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13224	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, QEF, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/3325	0.27	0/4517
1	B	0.10	0/3296	0.27	0/4474
All	All	0.10	0/6621	0.27	0/8991

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3259	3174	3172	10	0
1	B	3228	3146	3140	8	0
2	A	4	6	6	0	0
3	A	53	30	31	0	0
3	B	53	30	31	0	0
4	A	13	9	0	0	0
4	B	13	9	0	0	0
5	A	15	0	0	0	0
5	B	10	0	0	0	0
6	A	90	0	0	1	0
6	B	82	0	0	0	0
All	All	6820	6404	6380	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:MET:SD	1:B:250:GLN:NE2	2.62	0.72
1:A:581:ILE:HD11	1:A:610:ILE:HD11	1.91	0.53
1:B:129:PRO:HD3	1:B:214:ALA:HB1	1.93	0.51
1:A:172:ARG:N	1:A:173:PRO:CD	2.76	0.48
1:B:172:ARG:N	1:B:173:PRO:CD	2.77	0.48
1:A:482:PHE:CZ	1:A:494:ALA:HB3	2.50	0.47
1:B:267:LEU:HD12	1:B:398:VAL:HG21	1.96	0.47
1:A:148:ARG:NH1	6:A:802:HOH:O	2.46	0.46
1:A:267:LEU:HD12	1:A:398:VAL:HG21	1.96	0.46
1:A:158:ARG:NH1	1:A:217:LEU:O	2.47	0.45
1:B:267:LEU:HD21	1:B:306:ILE:HG21	2.00	0.44
1:A:368:ILE:N	1:A:368:ILE:HD12	2.33	0.43
1:A:565:ILE:HD12	1:A:565:ILE:N	2.33	0.43
1:A:128:ALA:HB2	1:A:228:LEU:HD22	2.01	0.42
1:B:368:ILE:N	1:B:368:ILE:HD12	2.35	0.42
1:B:565:ILE:N	1:B:565:ILE:HD12	2.35	0.42
1:B:353:MET:HG3	1:B:363:VAL:HG11	2.02	0.41
1:A:267:LEU:HD11	1:A:398:VAL:HG11	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	429/543 (79%)	416 (97%)	13 (3%)	0	100	100
1	B	427/543 (79%)	414 (97%)	13 (3%)	0	100	100
All	All	856/1086 (79%)	830 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/445 (73%)	325 (100%)	0	100	100
1	B	320/445 (72%)	320 (100%)	0	100	100
All	All	645/890 (72%)	645 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	235	GLN
1	A	432	ASN
1	A	598	HIS
1	B	250	GLN
1	B	454	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FAD	A	702	-	58,58,58	1.44	4 (6%)	85,89,89	0.92	3 (3%)
5	SO4	A	704	-	4,4,4	0.23	0	6,6,6	0.09	0
5	SO4	B	703	-	4,4,4	0.24	0	6,6,6	0.08	0
4	QEF	A	703	-	14,14,14	1.37	2 (14%)	19,20,20	1.07	1 (5%)
5	SO4	B	704	-	4,4,4	0.24	0	6,6,6	0.07	0
2	EDO	A	701	-	3,3,3	0.40	0	2,2,2	0.44	0
5	SO4	A	705	-	4,4,4	0.24	0	6,6,6	0.11	0
5	SO4	A	706	-	4,4,4	0.24	0	6,6,6	0.08	0
3	FAD	B	701	-	58,58,58	1.45	4 (6%)	85,89,89	0.91	3 (3%)
4	QEF	B	702	-	14,14,14	1.39	3 (21%)	19,20,20	1.11	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	702	-	-	3/34/50/50	0/6/6/6
4	QEF	A	703	-	-	-	0/2/2/2
2	EDO	A	701	-	-	0/1/1/1	-
3	FAD	B	701	-	-	4/34/50/50	0/6/6/6
4	QEF	B	702	-	-	-	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	701	FAD	C9-C9A	6.24	1.49	1.39
3	A	702	FAD	C9-C9A	6.18	1.49	1.39
3	B	701	FAD	C6-C5X	6.02	1.49	1.40
3	A	702	FAD	C6-C5X	5.98	1.49	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	FAD	O5'-C5'	3.26	1.56	1.44
3	B	701	FAD	O5'-C5'	3.24	1.56	1.44
4	B	702	QEF	C04-N03	-2.67	1.33	1.37
4	A	703	QEF	C04-N03	-2.65	1.33	1.37
4	B	702	QEF	C11-C10	-2.39	1.37	1.43
4	A	703	QEF	C11-C10	-2.33	1.37	1.43
3	A	702	FAD	C1'-C2'	2.10	1.55	1.52
3	B	701	FAD	C1'-C2'	2.08	1.55	1.52
4	B	702	QEF	C09-C10	-2.04	1.38	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	FAD	C4-N3-C2	-2.68	120.89	125.64
3	B	701	FAD	C4-N3-C2	-2.57	121.08	125.64
3	B	701	FAD	C5'-C4'-C3'	-2.39	107.70	112.22
3	B	701	FAD	C4'-C3'-C2'	-2.29	109.76	113.57
3	A	702	FAD	C5'-C4'-C3'	-2.27	107.93	112.22
4	B	702	QEF	C13-C02-N03	-2.17	120.13	122.60
3	A	702	FAD	O3B-C3B-C2B	-2.02	105.36	111.82
4	A	703	QEF	C13-C02-N03	-2.01	120.31	122.60

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	702	FAD	C3B-C4B-C5B-O5B
3	A	702	FAD	O4B-C4B-C5B-O5B
3	B	701	FAD	C3B-C4B-C5B-O5B
3	B	701	FAD	PA-O3P-P-O5'
3	B	701	FAD	O4B-C4B-C5B-O5B
3	A	702	FAD	C2B-C1B-N9A-C8A
3	B	701	FAD	C2B-C1B-N9A-C8A


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	436/543 (80%)	0.29	26 (5%)	27 24	23, 55, 86, 111	1 (0%)
1	B	434/543 (79%)	0.42	33 (7%)	20 16	38, 59, 91, 133	0
All	All	870/1086 (80%)	0.35	59 (6%)	23 20	23, 57, 90, 133	1 (0%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	489	ASP	5.6
1	B	570	ASP	5.5
1	B	509	ALA	5.0
1	A	183	ASP	5.0
1	A	199	LYS	4.9
1	B	599	GLU	4.8
1	A	487	GLY	4.6
1	A	127	LYS	4.5
1	A	561	GLY	4.1
1	B	561	GLY	4.1
1	B	183	ASP	4.0
1	A	184	ASP	4.0
1	A	618	PHE	3.8
1	B	618	PHE	3.7
1	B	486	LEU	3.5
1	B	129	PRO	3.5
1	B	221	GLU	3.4
1	A	619	GLN	3.3
1	A	463[A]	ARG	3.3
1	B	490	VAL	3.3
1	A	560	TYR	3.2
1	B	615	GLU	3.2
1	B	569	ARG	3.2
1	A	509	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	194	LYS	3.1
1	B	199	LYS	3.1
1	B	611	HIS	3.1
1	A	570	ASP	3.1
1	A	599	GLU	3.0
1	B	222	ASN	3.0
1	B	220	ILE	2.8
1	A	273	ALA	2.8
1	B	571	LYS	2.8
1	A	490	VAL	2.7
1	B	195	GLN	2.6
1	A	215	GLN	2.6
1	A	384	LYS	2.5
1	B	488	PRO	2.5
1	B	477	TRP	2.5
1	B	215	GLN	2.5
1	B	596	GLU	2.4
1	A	195	GLN	2.4
1	A	597	GLN	2.4
1	B	497	LEU	2.4
1	B	619	GLN	2.4
1	B	182	SER	2.4
1	B	265	ARG	2.3
1	B	274	GLY	2.3
1	B	600	ASP	2.3
1	B	616	VAL	2.3
1	A	611	HIS	2.2
1	B	573	VAL	2.2
1	A	370	GLN	2.1
1	A	376	SER	2.1
1	A	271	ASP	2.1
1	A	488	PRO	2.1
1	B	272	ARG	2.1
1	B	370	GLN	2.0
1	A	616	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	A	705	5/5	0.70	0.15	68,70,101,101	0
5	SO4	B	704	5/5	0.70	0.12	81,89,94,108	0
5	SO4	A	706	5/5	0.77	0.10	95,96,110,111	0
2	EDO	A	701	4/4	0.94	0.14	42,50,58,66	0
5	SO4	B	703	5/5	0.95	0.15	51,55,72,76	0
4	QEF	A	703	13/13	0.95	0.08	37,46,55,55	0
5	SO4	A	704	5/5	0.96	0.13	51,57,64,65	0
4	QEF	B	702	13/13	0.97	0.07	39,45,54,54	0
3	FAD	B	701	53/53	0.97	0.07	35,45,57,71	0
3	FAD	A	702	53/53	0.97	0.07	31,41,52,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around QEF A 703:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around QEF B 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.