



## wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 05:29 PM UTC

PDB ID : 7D44 / pdb\_00007d44  
EMDB ID : EMD-30569  
Title : eIF2B-eIF2(aP), aP2 complex  
Authors : Kashiwagi, K.; Ito, T.  
Deposited on : 2020-09-22  
Resolution : 4.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	7587 ( 3.50 - 4.50 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	58% (Green), 38% (Yellow), 4% (Grey)
1	B	305	52% (Green), 43% (Yellow), 5% (Grey)
2	C	351	52% (Green), 38% (Yellow), 10% (Grey)
2	D	351	55% (Green), 38% (Yellow), 8% (Grey)

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	452	 51% 25% 24%
3	F	452	 45% 31% 24%
4	G	523	 36% 31% 33%
4	H	523	 39% 28% 33%
5	I	721	 31% 28% 41%
5	J	721	 33% 26% 40%
6	K	315	 52% 44%
6	L	315	 10% 51% 44%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 28391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Translation initiation factor eIF-2B subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	293	Total	C	N	O	S	0	0
			2262	1449	377	424	12		
1	B	291	Total	C	N	O	S	0	0
			2247	1442	373	420	12		

- Molecule 2 is a protein called Translation initiation factor eIF-2B subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	317	Total	C	N	O	S	0	0
			2476	1565	435	461	15		
2	D	324	Total	C	N	O	S	0	0
			2538	1605	445	473	15		

- Molecule 3 is a protein called Translation initiation factor eIF-2B subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	343	Total	C	N	O	S	0	0
			2430	1540	423	452	15		
3	F	343	Total	C	N	O	S	0	0
			2429	1540	423	451	15		

- Molecule 4 is a protein called Translation initiation factor eIF-2B subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	G	353	Total	C	N	O	S	0	0
			2750	1740	490	506	14		
4	H	353	Total	C	N	O	S	0	0
			2750	1740	490	506	14		

- Molecule 5 is a protein called Translation initiation factor eIF-2B subunit epsilon.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	I	427	Total 3358	C 2115	N 591	O 637	S 15	0	0
5	J	429	Total 3373	C 2124	N 593	O 641	S 15	0	0


- Molecule 6 is a protein called Eukaryotic translation initiation factor 2 subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
6	K	176	Total 889	C 529	N 179	O 180	P 1	0	0
6	L	176	Total 889	C 529	N 179	O 180	P 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Translation initiation factor eIF-2B subunit alpha




- Molecule 1: Translation initiation factor eIF-2B subunit alpha




- Molecule 2: Translation initiation factor eIF-2B subunit beta







• Molecule 2: Translation initiation factor eIF-2B subunit beta




• Molecule 3: Translation initiation factor eIF-2B subunit gamma






• Molecule 3: Translation initiation factor eIF-2B subunit gamma



• Molecule 4: Translation initiation factor eIF-2B subunit delta







ARG  
ALA  
GLY  
LEU  
LEU  
LEU  
ASN  
CYS  
SER  
THR  
THR  
GLU  
THR  
ASP  
ASN  
MET  
GLU  
PRO  
PRO  
ILE  
ILE  
LYS  
ILE  
ILE  
ASN  
ARG  
GLN  
LEU  
LEU  
ILE  
ILE  
ALA  
PRO  
PRO  
LEU  
LEU  
ARG  
TYR  
VAL  
GLU  
MET  
MET  
THR  
THR  
THR  
THR  
THR  
THR  
GLY  
GLU  
GLY  
LEU  
LEU  
SER  
SER  
VAL  
VAL  
LEU  
SER  
SER  
GLN  
ALA  
ALA  
MET  
ALA  
ALA  
VAL

MET  
GLU  
PRO  
LYS  
VAL  
VAL  
THR  
THR  
ASP  
THR  
ASP  
ASN  
GLU  
THR  
THR  
GLU  
LEU  
ALA  
ALA  
GLN  
GLN  
MET  
MET  
GLU  
ARG  
ARG  
LEU  
GLU  
ARG  
GLU  
ASN  
ASN  
ALA  
ALA  
THR  
THR  
THR  
THR  
VAL  
VAL  
ASP  
GLY  
LEU  
LEU  
ASP  
ASP  
ASP  
ASP  
ALA  
GLU  
GLU  
GLU  
LEU  
LEU  
SER  
MET  
GLU  
ALA  
LEU  
ASP

● Molecule 6: Eukaryotic translation initiation factor 2 subunit 1



MET  
PRO  
GLY  
LEU  
SER  
G5  
A27  
G30  
L35  
L36  
M40  
E49  
L50  
S51  
R56  
I62  
R63  
P91  
E92  
E93  
A94  
K96  
C97  
E98  
D99  
K103  
S104  
K105  
Y108  
S109  
I110  
L111  
R112  
H113  
V114  
A115  
E116  
V117  
L118  
T121  
K122  
D123  
E124  
Q125  
L126  
E127  
S128

D138  
P144  
G147  
S157  
D158  
P159  
S160  
I161  
E171  
V174  
M178  
I179  
M190  
ARG  
ARG  
LEU  
LEU  
THR  
THR  
PRO  
GLN  
SER  
GLN  
ALA  
VAL  
VAL  
LEU  
VAL  
LYS  
SER  
LYS  
ILE  
ILE  
GLN  
ARG  
I95  
ALA  
ALA  
ASP  
ASP  
ILE  
ILE  
GLU  
GLU  
VAL  
VAL  
ALA  
ALA  
CYS  
TYR  
TYR  
TYR  
GLY  
GLY  
GLY  
GLY  
GLY  
ILE  
ILE  
ASP  
ASP  
PHE  
PHE  
ALA  
ALA  
VAL  
VAL  
ASN  
ASN  
VAL  
VAL  
GLN  
GLN  
MET  
MET  
LEU  
LEU  
ARG  
ARG  
LYS  
LYS  
ALA  
ALA  
VAL  
VAL  
LEU  
LEU  
ASN  
ASN  
CYS  
CYS  
THR  
THR  
THR  
THR  
GLU  
GLU  
ASN

MET  
PRO  
ILE  
ILE  
LYS  
ASN  
ASN  
LEU  
LEU  
ILE  
ILE  
ALA  
ALA  
PRO  
PRO  
ARG  
ARG  
TYR  
VAL  
VAL  
MET  
MET  
THR  
THR  
THR  
THR  
THR  
THR  
LEU  
LEU  
GLU  
GLU  
ARG  
ARG  
THR  
THR  
GLY  
GLY  
LEU  
LEU  
SER  
SER  
GLN  
GLN  
ALA  
ALA  
MET  
MET  
VAL  
VAL  
ILE  
ILE  
GLU  
GLU  
TYR  
TYR  
LYS  
LYS  
GLU  
GLU  
ARG  
ARG  
GLY  
GLY  
VAL  
VAL  
PHE  
PHE  
ALA  
ALA  
ASN  
ASN  
VAL  
VAL  
GLN  
GLN  
MET  
MET  
GLU  
GLU  
PRO  
PRO  
LYS  
LYS  
VAL  
VAL  
VAL  
VAL  
THR  
THR  
ASP  
ASP  
THR  
THR  
ASP

GLU  
THR  
GLU  
LEU  
ALA  
ARG  
GLN  
MET  
GLU  
ARG  
LEU  
GLU  
ARG  
GLU  
ALA  
ASN  
GLU  
VAL  
VAL  
ASP  
ASP  
GLY  
ASP  
ASP  
ASP  
ALA  
ALA  
GLU  
GLU  
ALA  
ALA  
GLU  
ASP

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	80921	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.166	Depositor
Minimum map value	-0.073	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.015	Depositor
Map size ( $\text{\AA}$ )	441.0, 441.0, 441.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.47, 1.47, 1.47	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/2296	0.54	0/3099
1	B	0.27	0/2281	0.50	0/3078
2	C	0.30	0/2522	0.54	2/3411 (0.1%)
2	D	0.32	0/2587	0.58	0/3498
3	E	0.22	0/2464	0.49	0/3346
3	F	0.23	0/2463	0.53	0/3344
4	G	0.30	0/2802	0.54	0/3809
4	H	0.30	0/2802	0.54	0/3809
5	I	0.26	0/3427	0.51	0/4662
5	J	0.29	0/3443	0.55	0/4685
6	K	0.23	0/879	0.68	5/1224 (0.4%)
6	L	0.17	0/879	0.43	0/1224
All	All	0.28	0/28845	0.53	7/39189 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	K	0	1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	K	12	PHE	CA-C-N	8.94	131.02	119.84
6	K	12	PHE	C-N-CA	8.94	131.02	119.84
6	K	14	GLU	N-CA-C	5.79	116.72	108.74
6	K	14	GLU	CA-C-N	5.68	128.24	120.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	K	14	GLU	C-N-CA	5.68	128.24	120.35

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
6	K	14	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2262	0	2333	97	0
1	B	2247	0	2320	108	0
2	C	2476	0	2493	114	0
2	D	2538	0	2538	106	0
3	E	2430	0	2197	93	0
3	F	2429	0	2196	109	0
4	G	2750	0	2811	134	0
4	H	2750	0	2811	129	0
5	I	3358	0	3325	156	0
5	J	3373	0	3336	154	0
6	K	889	0	401	5	0
6	L	889	0	402	11	0
All	All	28391	0	27163	1140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 1140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:L:35:LEU:O	6:L:40:ASN:HA	1.67	0.94
2:C:247:ALA:HB3	2:C:251:THR:HG21	1.52	0.91
1:B:205:GLY:HA3	1:B:276:THR:O	1.73	0.89

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:198:CYS:O	2:C:201:HIS:HB2	1.73	0.89
1:A:201:VAL:HG13	1:A:207:ILE:HD11	1.54	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/305 (95%)	254 (88%)	35 (12%)	0	100	100
1	B	287/305 (94%)	259 (90%)	28 (10%)	0	100	100
2	C	313/351 (89%)	271 (87%)	42 (13%)	0	100	100
2	D	320/351 (91%)	271 (85%)	48 (15%)	1 (0%)	36	70
3	E	333/452 (74%)	260 (78%)	73 (22%)	0	100	100
3	F	333/452 (74%)	262 (79%)	71 (21%)	0	100	100
4	G	351/523 (67%)	299 (85%)	52 (15%)	0	100	100
4	H	351/523 (67%)	305 (87%)	46 (13%)	0	100	100
5	I	425/721 (59%)	345 (81%)	80 (19%)	0	100	100
5	J	427/721 (59%)	344 (81%)	82 (19%)	1 (0%)	43	75
6	K	173/315 (55%)	144 (83%)	29 (17%)	0	100	100
6	L	173/315 (55%)	143 (83%)	30 (17%)	0	100	100
All	All	3775/5334 (71%)	3157 (84%)	616 (16%)	2 (0%)	49	81

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	127	ALA
5	J	249	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/260 (95%)	248 (100%)	0	100	100
1	B	247/260 (95%)	246 (100%)	1 (0%)	84	84
2	C	269/298 (90%)	268 (100%)	1 (0%)	84	84
2	D	276/298 (93%)	275 (100%)	1 (0%)	84	84
3	E	214/398 (54%)	214 (100%)	0	100	100
3	F	214/398 (54%)	213 (100%)	1 (0%)	81	82
4	G	309/444 (70%)	307 (99%)	2 (1%)	78	81
4	H	309/444 (70%)	309 (100%)	0	100	100
5	I	381/626 (61%)	381 (100%)	0	100	100
5	J	383/626 (61%)	381 (100%)	2 (0%)	81	82
6	K	3/279 (1%)	3 (100%)	0	100	100
6	L	3/279 (1%)	3 (100%)	0	100	100
All	All	2856/4610 (62%)	2848 (100%)	8 (0%)	84	85

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	J	400	VAL
5	J	133	ASP
4	G	214	TYR
3	F	161	ASP
4	G	394	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 65 such sidechains are listed below:

Mol	Chain	Res	Type
5	I	416	ASN
5	J	72	ASN
4	G	166	GLN
3	F	283	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
5	J	158	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	SEP	K	51	6	8,9,10	1.61	1 (12%)	7,12,14	1.18	1 (14%)
6	SEP	L	51	6	8,9,10	1.60	1 (12%)	7,12,14	1.25	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SEP	K	51	6	-	2/6/8/10	-
6	SEP	L	51	6	-	4/6/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	K	51	SEP	P-O1P	3.54	1.61	1.50
6	L	51	SEP	P-O1P	3.50	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
6	L	51	SEP	OG-CB-CA	2.64	110.72	108.14
6	K	51	SEP	OG-CB-CA	2.47	110.55	108.14

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	K	51	SEP	N-CA-CB-OG
6	L	51	SEP	N-CA-CB-OG
6	L	51	SEP	CB-OG-P-O1P
6	L	51	SEP	CB-OG-P-O2P
6	L	51	SEP	CB-OG-P-O3P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-30569. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 154



Y Index: 140




Z Index: 157


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views

This section was not generated.


## 6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 410 nm<sup>3</sup>; this corresponds to an approximate mass of 370 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of 0.250 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-30569 and PDB model 7D44. Per-residue inclusion information can be found in section 3 on page 6.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).



























## 9.4 Atom inclusion [i](#)



At the recommended contour level, 96% of all backbone atoms, 94% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9360	 0.4210
A	 0.9580	 0.4330
B	 0.9460	 0.4380
C	 0.9460	 0.4460
D	 0.9570	 0.4470
E	 0.8900	 0.3600
F	 0.9160	 0.3530
G	 0.9490	 0.4450
H	 0.9340	 0.4420
I	 0.9500	 0.4290
J	 0.9540	 0.4400
K	 0.9530	 0.3840
L	 0.7650	 0.3590

