



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 02:20 PM UTC

PDB ID : 3D68 / pdb_00003d68
Title : Crystal structure of a T325I/T329I/H333Y/H335Q mutant of Thrombin-Activatable Fibrinolysis Inhibitor (TAFI-IIYQ)
Authors : Brondijk, T.H.C.; Huizinga, E.G.
Deposited on : 2008-05-19
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	424	 2% 68% 23% •• 5%
1	B	424	 2% 64% 24% 5% • 5%
1	C	424	 2% 64% 23% 6% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	603	X	-	-	-
2	NAG	B	602	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carboxypeptidase B2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3249	2084	552	601	12	0	0	0
1	B	401	3249	2084	552	601	12	0	0	0
1	C	401	3249	2084	552	601	12	0	0	0

There are 87 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	GLY	-	expression tag	UNP Q96IY4
A	-21	SER	-	expression tag	UNP Q96IY4
A	-20	HIS	-	expression tag	UNP Q96IY4
A	-19	HIS	-	expression tag	UNP Q96IY4
A	-18	HIS	-	expression tag	UNP Q96IY4
A	-17	HIS	-	expression tag	UNP Q96IY4
A	-16	HIS	-	expression tag	UNP Q96IY4
A	-15	HIS	-	expression tag	UNP Q96IY4
A	-14	ASP	-	expression tag	UNP Q96IY4
A	-13	TYR	-	expression tag	UNP Q96IY4
A	-12	ASP	-	expression tag	UNP Q96IY4
A	-11	ILE	-	expression tag	UNP Q96IY4
A	-10	PRO	-	expression tag	UNP Q96IY4
A	-9	SER	-	expression tag	UNP Q96IY4
A	-8	SER	-	expression tag	UNP Q96IY4
A	-7	GLU	-	expression tag	UNP Q96IY4
A	-6	ASN	-	expression tag	UNP Q96IY4
A	-5	LEU	-	expression tag	UNP Q96IY4
A	-4	TYR	-	expression tag	UNP Q96IY4
A	-3	PHE	-	expression tag	UNP Q96IY4
A	-2	GLN	-	expression tag	UNP Q96IY4
A	-1	GLY	-	expression tag	UNP Q96IY4
A	0	SER	-	expression tag	UNP Q96IY4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	expression tag	UNP Q96IY4
A	147	THR	ALA	SEE REMARK 999	UNP Q96IY4
A	325	ILE	THR	engineered mutation	UNP Q96IY4
A	329	ILE	THR	engineered mutation	UNP Q96IY4
A	333	TYR	HIS	engineered mutation	UNP Q96IY4
A	335	GLN	HIS	engineered mutation	UNP Q96IY4
B	-22	GLY	-	expression tag	UNP Q96IY4
B	-21	SER	-	expression tag	UNP Q96IY4
B	-20	HIS	-	expression tag	UNP Q96IY4
B	-19	HIS	-	expression tag	UNP Q96IY4
B	-18	HIS	-	expression tag	UNP Q96IY4
B	-17	HIS	-	expression tag	UNP Q96IY4
B	-16	HIS	-	expression tag	UNP Q96IY4
B	-15	HIS	-	expression tag	UNP Q96IY4
B	-14	ASP	-	expression tag	UNP Q96IY4
B	-13	TYR	-	expression tag	UNP Q96IY4
B	-12	ASP	-	expression tag	UNP Q96IY4
B	-11	ILE	-	expression tag	UNP Q96IY4
B	-10	PRO	-	expression tag	UNP Q96IY4
B	-9	SER	-	expression tag	UNP Q96IY4
B	-8	SER	-	expression tag	UNP Q96IY4
B	-7	GLU	-	expression tag	UNP Q96IY4
B	-6	ASN	-	expression tag	UNP Q96IY4
B	-5	LEU	-	expression tag	UNP Q96IY4
B	-4	TYR	-	expression tag	UNP Q96IY4
B	-3	PHE	-	expression tag	UNP Q96IY4
B	-2	GLN	-	expression tag	UNP Q96IY4
B	-1	GLY	-	expression tag	UNP Q96IY4
B	0	SER	-	expression tag	UNP Q96IY4
B	1	ALA	-	expression tag	UNP Q96IY4
B	147	THR	ALA	SEE REMARK 999	UNP Q96IY4
B	325	ILE	THR	engineered mutation	UNP Q96IY4
B	329	ILE	THR	engineered mutation	UNP Q96IY4
B	333	TYR	HIS	engineered mutation	UNP Q96IY4
B	335	GLN	HIS	engineered mutation	UNP Q96IY4
C	-22	GLY	-	expression tag	UNP Q96IY4
C	-21	SER	-	expression tag	UNP Q96IY4
C	-20	HIS	-	expression tag	UNP Q96IY4
C	-19	HIS	-	expression tag	UNP Q96IY4
C	-18	HIS	-	expression tag	UNP Q96IY4
C	-17	HIS	-	expression tag	UNP Q96IY4
C	-16	HIS	-	expression tag	UNP Q96IY4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-15	HIS	-	expression tag	UNP Q96IY4
C	-14	ASP	-	expression tag	UNP Q96IY4
C	-13	TYR	-	expression tag	UNP Q96IY4
C	-12	ASP	-	expression tag	UNP Q96IY4
C	-11	ILE	-	expression tag	UNP Q96IY4
C	-10	PRO	-	expression tag	UNP Q96IY4
C	-9	SER	-	expression tag	UNP Q96IY4
C	-8	SER	-	expression tag	UNP Q96IY4
C	-7	GLU	-	expression tag	UNP Q96IY4
C	-6	ASN	-	expression tag	UNP Q96IY4
C	-5	LEU	-	expression tag	UNP Q96IY4
C	-4	TYR	-	expression tag	UNP Q96IY4
C	-3	PHE	-	expression tag	UNP Q96IY4
C	-2	GLN	-	expression tag	UNP Q96IY4
C	-1	GLY	-	expression tag	UNP Q96IY4
C	0	SER	-	expression tag	UNP Q96IY4
C	1	ALA	-	expression tag	UNP Q96IY4
C	147	THR	ALA	SEE REMARK 999	UNP Q96IY4
C	325	ILE	THR	engineered mutation	UNP Q96IY4
C	329	ILE	THR	engineered mutation	UNP Q96IY4
C	333	TYR	HIS	engineered mutation	UNP Q96IY4
C	335	GLN	HIS	engineered mutation	UNP Q96IY4

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		


- Molecule 4 is ARGinine (CCD ID: ARG) (formula: C₆H₁₅N₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			
4	A	1	Total	12	6	4	2	0	0
4	B	1	Total	12	6	4	2	0	0
4	C	1	Total	12	6	4	2	0	0

- Molecule 1: Carboxypeptidase B2

Chain C: 2%
64% 23% 6% 5%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	159.46Å 159.46Å 139.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.03 – 2.80 49.03 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.03-2.80) 100.0 (49.03-2.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.3.0008	Depositor
R, R_{free}	0.188 , 0.232 0.194 , 0.239	Depositor DCC
R_{free} test set	2582 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	69.9	Xtrriage
Anisotropy	0.317	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 70.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9940	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.05	3/3336 (0.1%)	1.12	10/4532 (0.2%)
1	B	1.10	1/3336 (0.0%)	1.15	20/4532 (0.4%)
1	C	1.08	3/3336 (0.1%)	1.13	14/4532 (0.3%)
All	All	1.08	7/10008 (0.1%)	1.13	44/13596 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
1	C	0	5
All	All	0	11

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	380	LYS	C-O	-6.07	1.19	1.24
1	C	88	THR	N-CA	5.78	1.53	1.46
1	B	88	THR	N-CA	5.76	1.53	1.46
1	A	343	ALA	CA-CB	-5.51	1.47	1.53
1	A	380	LYS	C-O	-5.49	1.19	1.24

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	366	ASP	N-CA-C	-10.04	89.41	110.80
1	C	67	ILE	N-CA-C	7.43	115.27	107.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	297	PRO	CA-C-N	7.39	135.65	121.54
1	B	297	PRO	C-N-CA	7.39	135.65	121.54
1	B	243	CYS	N-CA-C	7.00	121.15	112.47

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	297	PRO	Peptide
1	A	326	SER	Peptide
1	A	365	ARG	Peptide
1	A	37	ALA	Peptide
1	B	297	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3249	0	3167	75	0
1	B	3249	0	3167	96	0
1	C	3249	0	3168	85	0
2	A	56	0	52	4	0
2	B	56	0	52	4	0
2	C	42	0	39	7	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	12	0	12	3	0
4	B	12	0	12	0	0
4	C	12	0	12	0	0
All	All	9940	0	9681	262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:TRP:HE1	1:B:215:MET:HE2	1.12	1.09
1:C:176:ILE:HG21	1:C:190:LEU:HD11	1.35	1.02
1:A:325:ILE:HD12	1:A:388:ALA:HB2	1.36	1.01
1:B:351:ILE:HD12	1:B:356:ILE:HG12	1.43	0.99
1:B:8:ALA:HB2	1:B:72:LEU:HD21	1.46	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/424 (94%)	358 (90%)	31 (8%)	10 (2%)	4	16
1	B	399/424 (94%)	359 (90%)	29 (7%)	11 (3%)	4	14
1	C	399/424 (94%)	366 (92%)	22 (6%)	11 (3%)	4	14
All	All	1197/1272 (94%)	1083 (90%)	82 (7%)	32 (3%)	4	15

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	38	ASP
1	A	42	LYS
1	A	89	VAL
1	A	219	ASN
1	A	291	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	354/375 (94%)	330 (93%)	24 (7%)	14	42
1	B	354/375 (94%)	321 (91%)	33 (9%)	8	27
1	C	354/375 (94%)	317 (90%)	37 (10%)	6	22
All	All	1062/1125 (94%)	968 (91%)	94 (9%)	9	29

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	3	SER
1	C	175	HIS
1	C	25	THR
1	C	100	GLN
1	C	243	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	279	ASN
1	B	400	ASN
1	C	308	HIS
1	B	335	GLN
1	C	5	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 17 ligands modelled in this entry, 3 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	603	1	14,14,15	1.65	2 (14%)	17,19,21	2.49	7 (41%)
2	NAG	A	601	1	14,14,15	1.00	1 (7%)	17,19,21	2.52	3 (17%)
2	NAG	C	604	1	14,14,15	0.73	0	17,19,21	1.97	7 (41%)
2	NAG	B	601	1	14,14,15	0.69	0	17,19,21	1.75	2 (11%)
2	NAG	B	603	1	14,14,15	0.66	0	17,19,21	2.05	2 (11%)
2	NAG	A	602	1	14,14,15	0.66	0	17,19,21	1.28	3 (17%)
2	NAG	C	601	1	14,14,15	0.97	1 (7%)	17,19,21	2.95	3 (17%)
2	NAG	A	604	1	14,14,15	0.81	0	17,19,21	2.32	8 (47%)
2	NAG	B	602	1	14,14,15	0.73	0	17,19,21	2.01	5 (29%)
2	NAG	B	604	1	14,14,15	0.79	0	17,19,21	1.93	6 (35%)
2	NAG	C	602	1	14,14,15	1.56	2 (14%)	17,19,21	2.69	7 (41%)
4	ARG	A	650	-	10,11,11	1.09	1 (10%)	9,13,13	1.77	2 (22%)
4	ARG	B	650	-	10,11,11	0.85	0	9,13,13	1.56	2 (22%)
4	ARG	C	650	-	10,11,11	0.89	0	9,13,13	1.59	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	603	1	1/1/5/7	4/6/23/26	0/1/1/1
2	NAG	A	601	1	-	4/6/23/26	0/1/1/1
2	NAG	C	604	1	-	4/6/23/26	0/1/1/1
2	NAG	B	601	1	-	4/6/23/26	0/1/1/1
2	NAG	B	603	1	-	4/6/23/26	0/1/1/1
2	NAG	A	602	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	601	1	-	4/6/23/26	0/1/1/1
2	NAG	A	604	1	-	4/6/23/26	0/1/1/1
2	NAG	B	602	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	B	604	1	-	2/6/23/26	0/1/1/1
2	NAG	C	602	1	-	4/6/23/26	0/1/1/1
4	ARG	A	650	-	-	2/11/11/11	-
4	ARG	B	650	-	-	2/11/11/11	-
4	ARG	C	650	-	-	3/11/11/11	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	603	NAG	C1-C2	4.91	1.59	1.52
2	C	602	NAG	C1-C2	4.37	1.58	1.52
2	A	601	NAG	C1-C2	3.31	1.56	1.52
2	C	602	NAG	C2-N2	2.66	1.50	1.46
2	C	601	NAG	C1-C2	2.54	1.55	1.52

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	601	NAG	C1-O5-C5	10.76	126.61	112.19
2	A	601	NAG	C1-O5-C5	8.85	124.04	112.19
2	C	602	NAG	C1-O5-C5	7.04	121.62	112.19
2	B	603	NAG	C1-O5-C5	6.72	121.19	112.19
2	A	603	NAG	O5-C1-C2	-5.64	102.56	111.29

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	603	NAG	C1
2	B	602	NAG	C1

5 of 43 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	NAG	C8-C7-N2-C2
2	A	601	NAG	O7-C7-N2-C2
2	A	603	NAG	C8-C7-N2-C2
2	A	603	NAG	O7-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	601	NAG	C8-C7-N2-C2

There are no ring outliers.

8 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	NAG	2	0
2	C	604	NAG	2	0
2	B	603	NAG	2	0
2	C	601	NAG	1	0
2	A	604	NAG	2	0
2	B	604	NAG	2	0
2	C	602	NAG	4	0
4	A	650	ARG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/424 (94%)	0.32	7 (1%) 69 60	50, 67, 77, 95	0
1	B	401/424 (94%)	0.19	8 (1%) 65 56	51, 67, 81, 96	0
1	C	401/424 (94%)	0.25	7 (1%) 69 60	43, 67, 75, 97	0
All	All	1203/1272 (94%)	0.26	22 (1%) 67 58	43, 67, 77, 97	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	ALA	6.0
1	B	1	ALA	4.2
1	A	91	PRO	4.1
1	A	1	ALA	4.0
1	B	148	ALA	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	C	602	14/15	0.47	0.15	79,82,84,84	0
2	NAG	B	602	14/15	0.56	0.15	73,75,78,79	0
2	NAG	A	603	14/15	0.59	0.13	57,67,69,69	0
2	NAG	C	601	14/15	0.64	0.15	75,76,79,80	0
2	NAG	B	603	14/15	0.74	0.12	66,67,70,71	0
2	NAG	A	604	14/15	0.77	0.15	58,62,63,63	0
2	NAG	A	601	14/15	0.79	0.11	75,76,77,78	0
2	NAG	B	604	14/15	0.81	0.11	68,72,74,74	0
2	NAG	A	602	14/15	0.82	0.11	84,86,88,88	0
2	NAG	C	604	14/15	0.84	0.09	57,60,64,66	0
4	ARG	A	650	12/12	0.85	0.19	67,70,74,75	0
4	ARG	C	650	12/12	0.85	0.18	65,69,77,80	0
4	ARG	B	650	12/12	0.86	0.17	72,74,77,78	0
2	NAG	B	601	14/15	0.87	0.11	58,63,65,66	0
3	ZN	B	501	1/1	0.98	0.05	73,73,73,73	0
3	ZN	C	501	1/1	0.99	0.05	69,69,69,69	0
3	ZN	A	501	1/1	0.99	0.05	71,71,71,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.