



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 08:10 PM UTC

PDB ID : 3D6N / pdb_00003d6n
Title : Crystal Structure of Aquifex Dihydroorotase Activated by Aspartate Transcarbamoylase
Authors : Edwards, B.F.P.
Deposited on : 2008-05-20
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	16% 85% 12% .
2	B	291	4% 86% 12% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydroorotase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	422	Total	C	N	O	S	0	0	0
			3256	2071	554	615	16			


- Molecule 2 is a protein called Aspartate carbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	291	Total	C	N	O	S	0	0	0
			2376	1541	395	436	4			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CITRATE ANION (CCD ID: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 13	C 6	O 7	0	0
4	B	1	Total 13	C 6	O 7	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	209	Total 209	O 209	0	0
5	B	213	Total 213	O 213	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dihydroorotase



• Molecule 2: Aspartate carbamoyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	158.44Å 158.44Å 233.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	65.00 – 2.30 65.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (65.00-2.30) 100.0 (65.00-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.80	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.164 , 0.204 0.163 , 0.202	Depositor DCC
R_{free} test set	2544 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6081	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	3/3310 (0.1%)	1.09	5/4472 (0.1%)
2	B	1.23	6/2431 (0.2%)	1.17	8/3274 (0.2%)
All	All	1.14	9/5741 (0.2%)	1.12	13/7746 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	1	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	ARG	CA-C	7.50	1.62	1.52
2	B	45	SER	CA-C	-6.33	1.47	1.53
2	B	143	GLU	CB-CG	6.06	1.70	1.52
1	A	232	HIS	CA-C	-5.89	1.47	1.53
1	A	190	VAL	CA-CB	5.88	1.62	1.54
1	A	97	ASN	C-O	-5.59	1.20	1.25
2	B	38	VAL	CA-CB	5.53	1.60	1.54
2	B	251	GLY	N-CA	5.39	1.52	1.44
2	B	264	VAL	CA-CB	5.18	1.61	1.54

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	231	GLN	CA-C-N	-10.00	107.34	121.33
1	A	231	GLN	C-N-CA	-10.00	107.34	121.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	231	GLN	N-CA-C	8.71	121.93	111.82
2	B	123	THR	N-CA-CB	-7.87	98.54	110.73
2	B	123	THR	OG1-CB-CG2	7.51	124.33	109.30
2	B	249	HIS	CA-C-N	7.03	128.62	119.84
2	B	249	HIS	C-N-CA	7.03	128.62	119.84
1	A	8	ASN	OD1-CG-ND2	6.49	129.09	122.60
2	B	250	PRO	CA-C-N	-6.18	112.17	121.87
2	B	250	PRO	C-N-CA	-6.18	112.17	121.87
1	A	378	PHE	N-CA-C	5.65	118.28	109.52
2	B	68	GLY	N-CA-C	-5.22	108.41	115.36
2	B	73	THR	N-CA-C	5.04	121.53	110.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	123	THR	CB

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	72	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3256	0	3362	36	0
2	B	2376	0	2393	29	0
3	A	1	0	0	0	0
4	A	13	0	5	1	0
4	B	13	0	5	0	0
5	A	209	0	0	7	0
5	B	213	0	0	7	1
All	All	6081	0	5765	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:ARG:HD3	5:A:579:HOH:O	1.36	1.24
2:B:69:SER:HB3	2:B:72:SER:HA	1.20	1.09
2:B:1:MET:HE1	5:B:425:HOH:O	1.60	1.00
2:B:70:GLU:OE1	5:B:491:HOH:O	1.81	0.97
1:A:178:MET:HE3	5:A:468:HOH:O	1.68	0.94
2:B:2:ARG:HG3	2:B:289:LEU:HD22	1.58	0.85
2:B:69:SER:CB	2:B:72:SER:HA	2.05	0.84
2:B:211:TRP:HZ3	2:B:234:PHE:O	1.64	0.79
1:A:232:HIS:NE2	5:A:577:HOH:O	2.15	0.79
1:A:201:GLY:O	2:B:123:THR:CG2	2.33	0.77
2:B:2:ARG:HG3	2:B:289:LEU:CD2	2.17	0.74
1:A:59:ASP:OD2	1:A:304:THR:HG23	1.91	0.71
2:B:69:SER:HB3	2:B:72:SER:CA	2.12	0.70
1:A:339:ARG:NH2	1:A:406:LYS:O	2.26	0.69
2:B:126:HIS:HD2	5:B:427:HOH:O	1.76	0.69
2:B:2:ARG:HH11	2:B:2:ARG:HG2	1.58	0.67
2:B:70:GLU:HA	5:B:494:HOH:O	1.97	0.65
1:A:179:ASP:OD2	1:A:217:ASP:OD2	2.15	0.65
2:B:2:ARG:HH11	2:B:2:ARG:CG	2.12	0.62
1:A:93:MET:HE1	1:A:178:MET:HE1	1.80	0.62
1:A:264:THR:HG21	1:A:284:LYS:HE3	1.83	0.61
2:B:23:LYS:O	2:B:27:GLU:HG3	1.99	0.61
1:A:276:ARG:NH2	5:A:547:HOH:O	2.27	0.58
1:A:70:THR:HA	1:A:73:GLU:O	2.05	0.57
2:B:2:ARG:CG	2:B:289:LEU:CD2	2.83	0.56
1:A:201:GLY:O	2:B:123:THR:HG21	2.07	0.54
4:A:424:FLC:OA1	4:A:424:FLC:CGC	2.56	0.53
2:B:2:ARG:HG2	2:B:2:ARG:NH1	2.22	0.53
1:A:387:ASN:OD1	1:A:389:GLU:HG2	2.09	0.52
2:B:219:GLN:HG3	2:B:226:SER:HA	1.92	0.52
1:A:388:GLU:OE1	1:A:396:ARG:NH2	2.43	0.52
1:A:343:ILE:HB	1:A:347:LYS:HG2	1.91	0.51
2:B:211:TRP:CZ3	2:B:234:PHE:O	2.55	0.50
2:B:43:GLU:HB3	2:B:97:ARG:HG2	1.92	0.50
2:B:122:GLY:O	2:B:159:ARG:HD3	2.11	0.50
1:A:120:ARG:NH2	5:A:562:HOH:O	2.45	0.50
1:A:7:LYS:HA	1:A:23:ASP:OD1	2.13	0.49
1:A:261:LEU:HD13	1:A:337:LEU:HD11	1.95	0.49
1:A:208:GLU:OE2	2:B:186:ARG:NH2	2.43	0.48
1:A:329:THR:HG23	5:A:530:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:ARG:NE	5:A:547:HOH:O	2.36	0.48
1:A:1:MET:SD	1:A:44:GLU:HG3	2.53	0.47
2:B:2:ARG:HG2	2:B:289:LEU:HD23	1.96	0.47
1:A:210:GLU:OE1	1:A:234:SER:OG	2.28	0.47
1:A:59:ASP:OD2	1:A:304:THR:CG2	2.61	0.47
1:A:60:ILE:HA	1:A:91:VAL:HB	1.97	0.47
1:A:95:ASN:HB3	1:A:153:ASP:O	2.15	0.47
1:A:13:ASP:HB2	1:A:367:LEU:HD12	1.96	0.47
1:A:410:ILE:HA	1:A:422:ASP:HB2	1.96	0.46
2:B:2:ARG:CG	2:B:289:LEU:HD23	2.46	0.46
1:A:194:GLY:HA2	2:B:187:ASP:OD2	2.15	0.46
2:B:267:GLU:HB2	5:B:416:HOH:O	2.16	0.45
1:A:93:MET:CE	1:A:178:MET:HE1	2.46	0.45
2:B:126:HIS:CD2	5:B:427:HOH:O	2.61	0.43
1:A:126:THR:HG21	1:A:131:ARG:HG3	1.99	0.43
1:A:208:GLU:H	1:A:208:GLU:HG3	1.32	0.43
2:B:94:VAL:HB	2:B:116:LEU:HD23	1.99	0.43
1:A:126:THR:HA	1:A:151:THR:O	2.20	0.42
1:A:284:LYS:HE2	1:A:287:ARG:HH11	1.84	0.42
2:B:113:ASN:HB2	5:B:482:HOH:O	2.21	0.41
1:A:171:SER:HB3	1:A:226:GLY:HA2	2.04	0.40
1:A:388:GLU:CD	1:A:388:GLU:H	2.29	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:487:HOH:O	5:B:492:HOH:O[17_555]	2.11	0.09

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/422 (100%)	398 (95%)	19 (4%)	3 (1%)	18	23
2	B	289/291 (99%)	276 (96%)	11 (4%)	2 (1%)	18	23
All	All	709/713 (99%)	674 (95%)	30 (4%)	5 (1%)	18	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	72	SER
1	A	37	ASN
1	A	306	HIS
2	B	250	PRO
1	A	326	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/362 (100%)	345 (95%)	17 (5%)	23	35
2	B	262/262 (100%)	252 (96%)	10 (4%)	29	44
All	All	624/624 (100%)	597 (96%)	27 (4%)	26	39

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	5	ILE
1	A	36	LYS
1	A	37	ASN
1	A	40	VAL
1	A	143	LYS
1	A	208	GLU
1	A	232	HIS
1	A	284	LYS
1	A	329	THR
1	A	347	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	368	LYS
1	A	369	LEU
1	A	388	GLU
1	A	389	GLU
1	A	396	ARG
1	A	415	ASP
2	B	2	ARG
2	B	29	LYS
2	B	39	LEU
2	B	73	THR
2	B	106	LYS
2	B	123	THR
2	B	189	GLU
2	B	192	LYS
2	B	193	VAL
2	B	220	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
2	B	126	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FLC	A	424	3	12,12,12	1.94	2 (16%)	17,17,17	1.47	3 (17%)
4	FLC	B	292	-	12,12,12	1.41	2 (16%)	17,17,17	1.25	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FLC	A	424	3	-	10/16/16/16	-
4	FLC	B	292	-	-	7/16/16/16	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	424	FLC	CB-CBC	4.43	1.58	1.53
4	A	424	FLC	CA-CB	3.13	1.57	1.54
4	B	292	FLC	CB-CBC	2.87	1.56	1.53
4	B	292	FLC	OA2-CAC	-2.00	1.24	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	424	FLC	OB2-CBC-CB	3.13	119.14	113.14
4	B	292	FLC	OB2-CBC-CB	2.92	118.75	113.14
4	A	424	FLC	OB1-CBC-CB	-2.82	116.63	122.09
4	B	292	FLC	OB1-CBC-CB	-2.29	117.66	122.09
4	A	424	FLC	OG1-CGC-CG	-2.22	116.68	122.95
4	B	292	FLC	OG1-CGC-CG	-2.08	117.06	122.95

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	424	FLC	CG-CB-CBC-OB1
4	A	424	FLC	CG-CB-CBC-OB2
4	A	424	FLC	OHB-CB-CBC-OB1
4	A	424	FLC	OHB-CB-CBC-OB2
4	A	424	FLC	CA-CB-CG-CGC
4	B	292	FLC	CG-CB-CBC-OB1
4	B	292	FLC	CG-CB-CBC-OB2
4	B	292	FLC	OHB-CB-CBC-OB1
4	B	292	FLC	OHB-CB-CBC-OB2
4	A	424	FLC	OHB-CB-CG-CGC
4	A	424	FLC	CAC-CA-CB-CBC
4	A	424	FLC	CBC-CB-CG-CGC
4	B	292	FLC	CAC-CA-CB-CBC
4	B	292	FLC	CAC-CA-CB-OHB
4	A	424	FLC	CAC-CA-CB-CG
4	B	292	FLC	CAC-CA-CB-CG
4	A	424	FLC	CAC-CA-CB-OHB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	424	FLC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/422 (100%)	1.16	66 (15%) 5 6	24, 38, 49, 68	0
2	B	291/291 (100%)	-0.26	12 (4%) 41 43	18, 26, 44, 78	0
All	All	713/713 (100%)	0.58	78 (10%) 10 12	18, 32, 48, 78	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	8.7
1	A	2	LEU	6.6
2	B	73	THR	5.6
2	B	74	VAL	5.0
1	A	39	LEU	4.5
1	A	26	VAL	4.2
1	A	41	PRO	4.1
1	A	43	ALA	4.0
1	A	6	VAL	3.9
1	A	42	GLU	3.8
1	A	4	LEU	3.8
1	A	376	THR	3.8
1	A	27	GLU	3.7
1	A	40	VAL	3.7
2	B	72	SER	3.5
1	A	384	TRP	3.5
2	B	2	ARG	3.4
1	A	5	ILE	3.3
2	B	75	LYS	3.3
1	A	24	ILE	3.2
2	B	71	SER	3.1
2	B	69	SER	3.0
2	B	68	GLY	2.9
1	A	374	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	8	ASN	2.8
1	A	136	ILE	2.7
2	B	291	THR	2.7
1	A	132	LYS	2.7
1	A	316	LEU	2.7
1	A	337	LEU	2.7
1	A	33	LYS	2.7
1	A	32	LYS	2.6
2	B	76	GLY	2.6
1	A	3	LYS	2.6
1	A	45	ILE	2.6
1	A	133	GLY	2.5
1	A	46	ILE	2.5
1	A	18	LEU	2.5
1	A	25	LEU	2.5
1	A	47	ASP	2.5
1	A	50	GLY	2.5
1	A	36	LYS	2.5
1	A	35	ASP	2.3
1	A	398	THR	2.3
1	A	34	ILE	2.3
1	A	188	TYR	2.3
1	A	130	GLY	2.3
1	A	131	ARG	2.3
1	A	49	LYS	2.3
2	B	211	TRP	2.3
1	A	134	LYS	2.3
1	A	317	VAL	2.3
1	A	261	LEU	2.2
1	A	28	ASN	2.2
1	A	44	GLU	2.2
1	A	413	ILE	2.2
1	A	115	SER	2.2
1	A	128	THR	2.2
1	A	53	VAL	2.2
1	A	10	TYR	2.1
1	A	38	ILE	2.1
1	A	186	LEU	2.1
1	A	212	ILE	2.1
1	A	385	ILE	2.1
1	A	31	ILE	2.1
1	A	386	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	97	ASN	2.1
2	B	77	GLU	2.1
1	A	257	ASN	2.1
1	A	11	VAL	2.1
1	A	162	VAL	2.1
1	A	20	GLY	2.1
1	A	407	GLY	2.1
1	A	185	LYS	2.1
1	A	410	ILE	2.1
1	A	29	GLY	2.1
1	A	51	LEU	2.0
1	A	284	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FLC	A	424	13/13	0.76	0.15	31,38,46,50	0
4	FLC	B	292	13/13	0.78	0.35	70,72,75,75	0
3	ZN	A	423	1/1	0.96	0.22	54,54,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.