



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:06 AM UTC

PDB ID : 7D6I / pdb_00007d6i
Title : A neutralizing MAb targeting receptor-binding-domain of SARS-CoV-2
Authors : Shi, R.; Qi, J.X.; Gao, G.F.; Yan, J.H.
Deposited on : 2020-09-30
Resolution : 3.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1210 (3.48-3.36)
Clashscore	190562	1234 (3.48-3.36)
Ramachandran outliers	187476	1222 (3.48-3.36)
Sidechain outliers	187428	1222 (3.48-3.36)
RSRZ outliers	180081	1210 (3.48-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	
2	B	235	
3	C	217	
4	D	2	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	200	Total	C	N	O	S	0	0	0
			1548	989	255	296	8			

- Molecule 2 is a protein called Heavy chain of GH12-Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1593	1009	268	309	7			

- Molecule 3 is a protein called Light chain of GH12-Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	215	Total	C	N	O	S	0	0	0
			1599	994	261	339	5			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.60Å 150.60Å 121.32Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.30 – 3.41 49.30 – 3.41	Depositor EDS
% Data completeness (in resolution range)	71.8 (49.30-3.41) 71.9 (49.30-3.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.66 (at 3.40Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.226 , 0.255 0.227 , 0.258	Depositor DCC
R_{free} test set	816 reflections (3.69%)	wwPDB-VP
Wilson B-factor (Å ²)	60.8	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 62.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.046 for -h,-k,l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	4768	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/1591	0.40	0/2172
2	B	0.21	0/1632	0.53	1/2224 (0.0%)
3	C	0.18	0/1639	0.50	2/2245 (0.1%)
All	All	0.17	0/4862	0.48	3/6641 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
3	C	0	1
All	All	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	100	GLY	N-CA-C	6.66	120.71	113.58
3	C	52	ASP	CA-C-N	-5.17	114.46	122.67
3	C	52	ASP	C-N-CA	-5.17	114.46	122.67

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	519	HIS	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	B	53	GLN	Peptide
3	C	156	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1548	0	1432	32	0
2	B	1593	0	1547	35	0
3	C	1599	0	1499	23	0
4	D	28	0	25	2	0
All	All	4768	0	4503	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:ALA:HB2	2:B:102:VAL:HG21	1.45	0.95
3:C:52:ASP:OD2	3:C:67:ILE:HD12	1.78	0.84
2:B:125:PRO:HB3	2:B:151:TYR:HB3	1.67	0.76
1:A:485:GLY:H	1:A:488:CYS:HB2	1.51	0.76
3:C:155:ALA:N	3:C:158:SER:O	2.18	0.75
2:B:102:VAL:HG13	2:B:103:ARG:HG2	1.70	0.74
3:C:156:ASP:OD2	3:C:194:ARG:N	2.24	0.69
1:A:521:PRO:HB2	1:A:523:THR:HG23	1.76	0.68
1:A:366:SER:HA	1:A:369:TYR:CZ	2.31	0.65
1:A:520:ALA:HB1	1:A:521:PRO:HD2	1.79	0.65
2:B:52:LYS:HB2	2:B:57:GLU:HG3	1.78	0.64
3:C:28:SER:O	3:C:31:SER:OG	2.11	0.64
2:B:34:MET:HB3	2:B:79:LEU:HD22	1.80	0.64
3:C:50:TYR:O	3:C:54:GLN:HB2	1.98	0.63
2:B:101:TRP:H	2:B:101:TRP:HD1	1.47	0.63
2:B:132:PRO:HD3	2:B:144:LEU:HB3	1.82	0.62
1:A:334:ASN:HB3	1:A:361:CYS:HA	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:388:ASN:HD22	1:A:527:PRO:HG2	1.65	0.62
2:B:199:THR:HG22	2:B:201:ILE:HD11	1.84	0.59
3:C:118:PRO:HB3	3:C:144:PHE:HB3	1.84	0.58
2:B:101:TRP:N	2:B:101:TRP:CD1	2.72	0.57
1:A:372:ALA:HA	2:B:103:ARG:HH11	1.70	0.56
1:A:528:LYS:O	1:A:528:LYS:HD2	2.04	0.56
1:A:384:PRO:HA	1:A:387:LEU:HG	1.87	0.55
2:B:101:TRP:H	2:B:101:TRP:CD1	2.24	0.55
2:B:132:PRO:HB3	2:B:143:ALA:O	2.06	0.55
1:A:388:ASN:HB3	1:A:527:PRO:HG2	1.89	0.55
2:B:101:TRP:CZ2	2:B:103:ARG:HB2	2.43	0.54
2:B:206:HIS:CD2	2:B:208:PRO:HD2	2.42	0.54
1:A:369:TYR:O	2:B:102:VAL:HG11	2.09	0.53
1:A:350:VAL:HG22	1:A:422:ASN:HB3	1.91	0.52
3:C:30:ALA:HB2	3:C:67:ILE:HD13	1.91	0.52
3:C:12:GLU:OE1	3:C:16:LYS:HG3	2.10	0.51
1:A:411:ALA:HB3	1:A:414:GLN:HG3	1.92	0.51
3:C:86:GLU:HG3	3:C:109:THR:HA	1.93	0.50
1:A:393:THR:HA	1:A:522:ALA:HA	1.92	0.50
3:C:99:LEU:HD12	3:C:99:LEU:H	1.76	0.50
2:B:102:VAL:HG22	2:B:102:VAL:O	2.11	0.50
2:B:1:GLU:O	2:B:26:GLY:HA3	2.12	0.50
3:C:15:GLY:O	3:C:16:LYS:HG2	2.13	0.48
2:B:101:TRP:HD1	2:B:101:TRP:N	2.10	0.48
1:A:365:TYR:CD2	1:A:387:LEU:HB3	2.48	0.48
2:B:57:GLU:OE1	2:B:59:TYR:OH	2.27	0.48
1:A:342:PHE:HB2	4:D:1:NAG:H82	1.95	0.48
3:C:56:PRO:HD2	3:C:59:VAL:HG21	1.95	0.48
3:C:199:GLN:HG2	3:C:208:GLU:HG3	1.95	0.47
1:A:341:VAL:HG11	1:A:397:ALA:HB1	1.96	0.47
1:A:528:LYS:O	1:A:528:LYS:CD	2.62	0.47
3:C:109:THR:HG21	3:C:146:PRO:HB3	1.96	0.47
2:B:12:VAL:O	2:B:117:VAL:HA	2.14	0.47
3:C:188:GLU:O	3:C:192:SER:OG	2.25	0.47
3:C:35:GLN:HG3	3:C:50:TYR:HA	1.96	0.46
1:A:401:VAL:HG22	1:A:509:ARG:HG2	1.97	0.46
2:B:33:TRP:CD1	2:B:101:TRP:O	2.68	0.46
1:A:403:ARG:HG3	1:A:406:GLU:HG3	1.98	0.46
3:C:49:ILE:HA	3:C:54:GLN:O	2.16	0.46
2:B:89:GLU:OE2	2:B:89:GLU:N	2.48	0.45
2:B:130:LEU:HD12	2:B:187:VAL:HG12	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:431:GLY:HA2	1:A:515:PHE:CD2	2.52	0.44
2:B:91:THR:HG23	2:B:116:THR:HA	1.99	0.44
1:A:366:SER:HA	1:A:369:TYR:CE2	2.52	0.44
3:C:52:ASP:OD2	3:C:67:ILE:CD1	2.57	0.44
1:A:396:TYR:HB2	1:A:514:SER:HB2	1.99	0.44
1:A:497:PHE:CE2	1:A:507:PRO:HB3	2.52	0.44
2:B:4:LEU:HD23	2:B:24:ALA:HA	2.00	0.44
1:A:379:CYS:HA	1:A:432:CYS:HA	2.01	0.43
2:B:101:TRP:CE2	2:B:103:ARG:HB2	2.53	0.43
2:B:175:VAL:HG13	3:C:167:THR:HG22	2.01	0.43
1:A:365:TYR:CE2	1:A:387:LEU:HB3	2.53	0.43
2:B:53:GLN:H	2:B:53:GLN:HG2	1.59	0.42
3:C:29:ILE:N	3:C:72:ASN:OD1	2.51	0.42
1:A:363:ALA:HB1	1:A:365:TYR:CE1	2.54	0.42
2:B:161:ASN:HD21	2:B:200:TYR:H	1.68	0.42
2:B:98:ARG:NH2	2:B:107:ASP:OD2	2.53	0.42
3:C:125:PRO:HD3	3:C:137:LEU:HG	2.01	0.41
2:B:175:VAL:O	2:B:182:TYR:HA	2.20	0.41
2:B:145:GLY:HA2	2:B:160:TRP:CZ2	2.55	0.41
3:C:55:ARG:HA	3:C:56:PRO:HD3	1.94	0.41
1:A:425:LEU:HD23	1:A:425:LEU:HA	1.94	0.41
3:C:185:LEU:HD23	3:C:190:TRP:HB2	2.03	0.41
2:B:168:GLY:O	2:B:188:VAL:HA	2.20	0.41
1:A:369:TYR:O	2:B:102:VAL:CG1	2.69	0.41
1:A:388:ASN:HB3	1:A:527:PRO:CG	2.51	0.41
2:B:194:SER:O	2:B:198:GLN:HB2	2.20	0.41
4:D:1:NAG:H62	4:D:2:NAG:H82	2.04	0.40
1:A:353:TRP:CD1	1:A:353:TRP:H	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/214 (92%)	198 (100%)	0	0	100	100
2	B	208/235 (88%)	205 (99%)	2 (1%)	1 (0%)	24	54
3	C	213/217 (98%)	210 (99%)	3 (1%)	0	100	100
All	All	619/666 (93%)	613 (99%)	5 (1%)	1 (0%)	43	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	102	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/187 (89%)	166 (100%)	0	100	100
2	B	175/198 (88%)	173 (99%)	2 (1%)	65	73
3	C	182/188 (97%)	181 (100%)	1 (0%)	81	80
All	All	523/573 (91%)	520 (99%)	3 (1%)	78	80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	101	TRP
2	B	103	ARG
3	C	52	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	360	ASN
3	C	199	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	1	4,1	14,14,15	0.39	0	17,19,21	0.60	1 (5%)
4	NAG	D	2	4	14,14,15	0.23	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	C1-O5-C5	2.02	114.89	112.19

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1	NAG	O5-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1	NAG	2	0
4	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/214 (93%)	0.46	11 (5%) 30 23	46, 90, 149, 168	0
2	B	212/235 (90%)	0.08	8 (3%) 44 31	27, 55, 89, 125	0
3	C	215/217 (99%)	0.05	5 (2%) 61 46	25, 58, 104, 138	0
All	All	627/666 (94%)	0.19	24 (3%) 44 31	25, 65, 122, 168	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	104	GLY	4.5
3	C	161	LYS	3.9
2	B	99	ALA	3.4
1	A	366	SER	3.3
2	B	100	GLY	3.2
3	C	214	THR	2.8
3	C	70	SER	2.6
3	C	99	LEU	2.6
2	B	131	ALA	2.6
2	B	3	GLN	2.4
1	A	467	ASP	2.4
1	A	527	PRO	2.4
2	B	132	PRO	2.4
1	A	386	LYS	2.3
1	A	521	PRO	2.2
2	B	199	THR	2.2
2	B	216	ARG	2.2
1	A	445	VAL	2.2
1	A	443	SER	2.1
1	A	483	VAL	2.1
3	C	25	SER	2.1
1	A	478	THR	2.1
1	A	518	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	333	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	D	2	14/15	0.71	0.14	102,119,123,133	0
4	NAG	D	1	14/15	0.80	0.12	71,100,107,108	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.