



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:17 AM UTC

PDB ID : 8D6D / pdb_00008d6d
Title : Crystal Structure of Human Myt1 Kinase domain Bounded with compound 39
Authors : Pau, V.P.T.; Mao, D.Y.L.; Mader, P.; Orlicky, S.; Sicheri, F.
Deposited on : 2022-06-06
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	8% 77% 9% 13%
1	B	311	9% 73% 15% 12%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	0	0
			1982	1273	344	354	11			
1	B	274	Total	C	N	O	S	0	0	0
			2011	1293	337	370	11			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	52	MET	-	initiating methionine	UNP Q99640
A	53	HIS	-	expression tag	UNP Q99640
A	54	HIS	-	expression tag	UNP Q99640
A	55	HIS	-	expression tag	UNP Q99640
A	56	HIS	-	expression tag	UNP Q99640
A	57	HIS	-	expression tag	UNP Q99640
A	58	HIS	-	expression tag	UNP Q99640
A	59	SER	-	expression tag	UNP Q99640
A	60	SER	-	expression tag	UNP Q99640
A	61	GLY	-	expression tag	UNP Q99640
A	62	VAL	-	expression tag	UNP Q99640
A	63	ASP	-	expression tag	UNP Q99640
A	64	LEU	-	expression tag	UNP Q99640
A	65	GLY	-	expression tag	UNP Q99640
A	66	THR	-	expression tag	UNP Q99640
A	67	GLU	-	expression tag	UNP Q99640
A	68	ASN	-	expression tag	UNP Q99640
A	69	LEU	-	expression tag	UNP Q99640
A	70	TYR	-	expression tag	UNP Q99640
A	71	PHE	-	expression tag	UNP Q99640
A	72	GLN	-	expression tag	UNP Q99640
A	73	SER	-	expression tag	UNP Q99640
A	74	MET	-	expression tag	UNP Q99640
B	52	MET	-	initiating methionine	UNP Q99640

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	53	HIS	-	expression tag	UNP Q99640
B	54	HIS	-	expression tag	UNP Q99640
B	55	HIS	-	expression tag	UNP Q99640
B	56	HIS	-	expression tag	UNP Q99640
B	57	HIS	-	expression tag	UNP Q99640
B	58	HIS	-	expression tag	UNP Q99640
B	59	SER	-	expression tag	UNP Q99640
B	60	SER	-	expression tag	UNP Q99640
B	61	GLY	-	expression tag	UNP Q99640
B	62	VAL	-	expression tag	UNP Q99640
B	63	ASP	-	expression tag	UNP Q99640
B	64	LEU	-	expression tag	UNP Q99640
B	65	GLY	-	expression tag	UNP Q99640
B	66	THR	-	expression tag	UNP Q99640
B	67	GLU	-	expression tag	UNP Q99640
B	68	ASN	-	expression tag	UNP Q99640
B	69	LEU	-	expression tag	UNP Q99640
B	70	TYR	-	expression tag	UNP Q99640
B	71	PHE	-	expression tag	UNP Q99640
B	72	GLN	-	expression tag	UNP Q99640
B	73	SER	-	expression tag	UNP Q99640
B	74	MET	-	expression tag	UNP Q99640

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is (1P)-2-amino-5-bromo-1-(3-hydroxy-2,6-dimethylphenyl)-1H-pyrrolo[2,3-b]quinoxaline-3-carboxamide (CCD ID: QGY) (formula: C₁₉H₁₆BrN₅O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	Br	C	N	O	0	0
			27	1	19	5	2		
4	B	1	Total	Br	C	N	O	0	0
			27	1	19	5	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	19	Total	O	0	0
			19	19		
5	B	14	Total	O	0	0
			14	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase



- Molecule 1: Membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.46Å 112.92Å 72.66Å 90.00° 110.34° 90.00°	Depositor
Resolution (Å)	58.33 – 2.35 58.33 – 2.35	Depositor EDS
% Data completeness (in resolution range)	94.8 (58.33-2.35) 95.3 (58.33-2.35)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 2.34Å)	Xtriage
Refinement program	PHENIX 1.19.2	Depositor
R, R_{free}	0.241 , 0.272 0.241 , 0.271	Depositor DCC
R_{free} test set	1541 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	76.6	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 82.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.100 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4110	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, QGY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/2030	0.44	0/2758
1	B	0.21	0/2060	0.39	0/2799
All	All	0.23	0/4090	0.42	0/5557

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1982	0	1888	15	0
1	B	2011	0	1886	25	0
2	A	12	0	18	0	0
2	B	8	0	12	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	27	0	0	0	0
4	B	27	0	0	1	0
5	A	19	0	0	0	0
5	B	14	0	0	0	0
All	All	4110	0	3804	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:320:LEU:HB3	1:B:325:THR:HG21	1.71	0.72
1:A:328:LEU:HD22	1:A:332:LEU:HD23	1.82	0.61
1:B:286:ALA:HB1	1:B:347:ARG:HD2	1.83	0.61
1:B:140:ARG:HH22	2:B:403:EDO:H21	1.66	0.60
1:A:320:LEU:HD21	1:A:340:LEU:HD12	1.86	0.57
1:B:277:GLU:HB2	1:B:282:SER:HB3	1.88	0.56
1:B:129:SER:HB3	1:B:132:ASP:OD1	2.07	0.55
1:A:274:MET:HB2	1:A:278:LEU:HD12	1.90	0.54
1:B:112:ARG:HA	1:B:127:VAL:HG12	1.93	0.51
1:B:331:GLU:HB3	1:B:358:VAL:HG11	1.91	0.51
1:B:155:LEU:HD12	1:B:183:LEU:HD22	1.93	0.51
1:A:112:ARG:HA	1:A:127:VAL:HG12	1.93	0.50
1:B:225:HIS:CG	1:B:285:THR:HB	2.48	0.49
1:A:214:LEU:HD13	1:A:296:ILE:HG13	1.95	0.48
1:B:107:GLN:HG2	1:B:112:ARG:HH12	1.78	0.48
1:A:172:ARG:NH1	1:A:174:GLU:OE1	2.47	0.47
1:A:229:LEU:HD22	1:A:255:LEU:HD21	1.96	0.47
1:A:356:LEU:O	1:A:360:ARG:HG3	2.15	0.47
1:B:187:THR:OG1	4:B:404:QGY:N12	2.47	0.47
1:B:229:LEU:HD22	1:B:255:LEU:HD21	1.96	0.47
1:B:97:TYR:O	1:B:99:PRO:HD3	2.15	0.47
1:B:242:GLY:N	1:B:246:ARG:O	2.45	0.47
1:B:334:SER:O	1:B:338:MET:HG3	2.15	0.47
1:A:334:SER:O	1:A:338:MET:HG3	2.14	0.46
1:A:357:PRO:HA	1:A:360:ARG:NE	2.30	0.46
1:B:122:GLY:HA3	1:B:140:ARG:O	2.15	0.46
1:B:214:LEU:HD13	1:B:296:ILE:HG13	1.97	0.46
1:A:256:VAL:HG11	1:A:266:VAL:HG22	1.97	0.46
1:B:309:GLY:O	1:B:313:GLN:HG2	2.16	0.45
1:B:290:PHE:CZ	1:B:316:ARG:HG2	2.52	0.44
1:A:272:ARG:HD3	1:A:307:HIS:CE1	2.53	0.44
1:B:80:ARG:HB3	1:B:89:SER:OG	2.18	0.43
1:B:286:ALA:C	1:B:347:ARG:HH11	2.26	0.43
1:B:121:TYR:HE2	1:B:124:VAL:HG23	1.83	0.42
1:B:335:VAL:O	1:B:339:MET:HG3	2.19	0.42
1:A:121:TYR:HE1	1:A:124:VAL:HG23	1.85	0.41
1:A:173:LEU:HD12	1:A:186:GLN:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:GLN:O	1:A:317:GLN:HG3	2.21	0.40
1:B:272:ARG:HD3	1:B:307:HIS:CE1	2.56	0.40
1:B:254:LEU:HG	1:B:267:GLN:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/311 (85%)	258 (98%)	6 (2%)	0	100	100
1	B	266/311 (86%)	257 (97%)	9 (3%)	0	100	100
All	All	530/622 (85%)	515 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/256 (74%)	190 (100%)	0	100	100
1	B	191/256 (75%)	189 (99%)	2 (1%)	68	81
All	All	381/512 (74%)	379 (100%)	2 (0%)	81	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	91	THR
1	B	323	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	HIS
1	B	227	GLN
1	B	307	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	QGY	A	405	-	28,30,30	1.60	5 (17%)	36,46,46	1.75	9 (25%)
4	QGY	B	404	-	28,30,30	1.46	6 (21%)	36,46,46	1.59	6 (16%)
2	EDO	A	401	-	3,3,3	0.43	0	2,2,2	0.27	0
3	SO4	A	403	-	4,4,4	0.24	0	6,6,6	0.13	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	402	-	4,4,4	0.21	0	6,6,6	0.11	0
2	EDO	A	404	-	3,3,3	0.42	0	2,2,2	0.30	0
2	EDO	B	403	-	3,3,3	0.38	0	2,2,2	0.41	0
2	EDO	B	401	-	3,3,3	0.44	0	2,2,2	0.33	0
2	EDO	A	402	-	3,3,3	0.42	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	QGY	A	405	-	-	0/8/8/8	0/4/4/4
4	QGY	B	404	-	-	0/8/8/8	0/4/4/4
2	EDO	A	401	-	-	0/1/1/1	-
2	EDO	A	404	-	-	0/1/1/1	-
2	EDO	B	403	-	-	0/1/1/1	-
2	EDO	B	401	-	-	0/1/1/1	-
2	EDO	A	402	-	-	0/1/1/1	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	405	QGY	C27-N26	4.09	1.36	1.31
4	A	405	QGY	C11-N12	2.95	1.40	1.34
4	B	404	QGY	C11-N12	2.73	1.40	1.34
4	A	405	QGY	C14-N15	2.71	1.41	1.33
4	A	405	QGY	C23-C22	2.70	1.43	1.38
4	B	404	QGY	C14-N15	2.69	1.41	1.33
4	B	404	QGY	C27-N26	2.65	1.34	1.31
4	B	404	QGY	C23-C22	2.49	1.43	1.38
4	B	404	QGY	C19-N18	2.39	1.42	1.37
4	B	404	QGY	C17-C27	2.27	1.48	1.43
4	A	405	QGY	C27-N10	2.18	1.41	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	405	QGY	N26-C27-N10	5.43	130.98	126.39
4	B	404	QGY	N26-C27-N10	4.52	130.21	126.39
4	B	404	QGY	C17-C27-N26	-4.03	121.96	125.49
4	A	405	QGY	C17-C27-N26	-3.99	121.99	125.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	405	QGY	C27-C17-N18	-3.27	119.18	122.08
4	A	405	QGY	C13-C17-N18	2.83	133.06	129.16
4	B	404	QGY	C27-C17-N18	-2.82	119.58	122.08
4	B	404	QGY	C13-C17-N18	2.72	132.91	129.16
4	B	404	QGY	O16-C14-N15	-2.71	116.81	122.89
4	A	405	QGY	O16-C14-N15	-2.57	117.13	122.89
4	A	405	QGY	C22-C20-C19	2.53	124.98	121.38
4	A	405	QGY	C24-C25-N26	2.41	122.21	118.69
4	A	405	QGY	C23-C22-C20	-2.32	117.26	120.20
4	B	404	QGY	C23-C24-C25	2.20	123.10	120.09
4	A	405	QGY	C23-C24-C25	2.14	123.02	120.09

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	404	QGY	1	0
2	B	403	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/311 (86%)	0.82	24 (8%) 15 17	67, 98, 146, 200	0
1	B	274/311 (88%)	0.91	28 (10%) 12 13	71, 104, 163, 203	0
All	All	544/622 (87%)	0.86	52 (9%) 13 16	67, 100, 158, 203	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	283	TYR	4.9
1	A	78	GLN	4.1
1	A	110	PHE	3.9
1	A	259	GLY	3.7
1	B	259	GLY	3.6
1	B	201	TRP	3.4
1	B	283	TYR	3.4
1	A	95	PRO	3.3
1	B	84	PHE	3.2
1	A	256	VAL	3.2
1	B	266	VAL	3.2
1	B	361	GLN	3.1
1	B	95	PRO	3.1
1	B	87	GLU	3.0
1	B	97	TYR	3.0
1	B	110	PHE	2.9
1	A	319	TYR	2.9
1	B	88	ALA	2.9
1	A	113	LEU	2.8
1	B	130	LYS	2.8
1	A	321	PRO	2.7
1	B	164	VAL	2.7
1	A	242	GLY	2.7
1	B	105	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	100	SER	2.6
1	B	321	PRO	2.5
1	B	177	TRP	2.5
1	A	105	PHE	2.5
1	A	161	HIS	2.5
1	B	77	LEU	2.4
1	B	252	PHE	2.4
1	B	319	TYR	2.4
1	A	96	GLY	2.4
1	B	194	LEU	2.4
1	A	102	PRO	2.3
1	B	355	ALA	2.3
1	B	108	GLN	2.3
1	B	155	LEU	2.2
1	B	131	GLU	2.2
1	B	299	VAL	2.2
1	A	99	PRO	2.2
1	A	282	SER	2.2
1	B	324	PHE	2.1
1	A	131	GLU	2.1
1	A	190	CYS	2.1
1	A	135	LEU	2.1
1	A	361	GLN	2.1
1	B	176	ALA	2.1
1	A	265	GLU	2.1
1	B	315	LEU	2.1
1	A	155	LEU	2.0
1	A	315	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	401	4/4	0.64	0.18	115,118,119,120	0
2	EDO	B	403	4/4	0.75	0.13	99,108,109,110	0
2	EDO	A	404	4/4	0.77	0.11	86,88,105,107	0
2	EDO	A	402	4/4	0.79	0.13	124,127,129,130	0
3	SO4	A	403	5/5	0.89	0.09	102,128,132,215	0
3	SO4	B	402	5/5	0.90	0.12	100,111,123,155	0
4	QGY	B	404	27/27	0.92	0.11	66,80,105,189	0
2	EDO	B	401	4/4	0.94	0.13	84,84,85,92	0
4	QGY	A	405	27/27	0.96	0.10	77,85,92,126	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around QGY B 404:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around QGY A 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.