



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:25 AM UTC

PDB ID : 1D7Y / pdb_00001d7y
Title : CRYSTAL STRUCTURE OF NADH-DEPENDENT FERREDOXIN REDUCTASE, BPHA4
Authors : Senda, T.; Yamada, T.; Sakurai, N.; Kubota, M.; Nishizaki, T.; Masai, E.; Fukuda, M.; Mitsui, Y.
Deposited on : 1999-10-21
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	408	 78% 16% . .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FERREDONIN REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	0	0
			2976	1865	544	559	8			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FERREDONIN REDUCTASE

Chain A:  78% 16%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.20Å 99.20Å 175.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.10	Depositor
% Data completeness (in resolution range)	97.4 (40.00-2.10)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.183 , 0.228	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3149	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/3028	0.89	7/4129 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184	ALA	N-CA-C	7.16	122.03	113.16
1	A	276	ARG	N-CA-C	-6.28	98.28	108.52
1	A	128	THR	N-CA-C	-5.77	100.73	109.85
1	A	64	ILE	N-CA-C	5.64	117.62	112.43
1	A	185	ALA	N-CA-C	5.31	118.33	109.09
1	A	51	LEU	N-CA-C	5.27	117.71	111.33
1	A	44	ARG	N-CA-C	-5.12	102.10	109.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2976	0	2995	49	0
2	A	53	0	31	0	0
3	A	120	0	0	2	0
All	All	3149	0	3026	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397:THR:HG22	1:A:400:ARG:H	1.24	1.01
1:A:287:ARG:HH11	1:A:287:ARG:HG2	1.53	0.74
1:A:123:THR:HG23	3:A:498:HOH:O	1.87	0.73
1:A:197:HIS:HE1	1:A:323:GLN:HE21	1.37	0.73
1:A:329:GLN:HE22	1:A:371:ALA:HA	1.56	0.71
1:A:197:HIS:HE1	1:A:323:GLN:NE2	1.89	0.69
1:A:279:ASN:C	1:A:279:ASN:HD22	2.02	0.68
1:A:307:ASP:OD1	1:A:309:THR:HB	1.94	0.67
1:A:397:THR:HG22	1:A:400:ARG:N	2.07	0.64
1:A:279:ASN:HD22	1:A:281:LEU:H	1.48	0.62
1:A:275:THR:O	1:A:287:ARG:HG3	2.02	0.60
1:A:145:GLN:CA	1:A:145:GLN:HE21	2.15	0.60
1:A:337:ASP:N	1:A:357:LEU:O	2.34	0.60
1:A:279:ASN:ND2	1:A:281:LEU:H	2.00	0.59
1:A:222:ASP:CG	1:A:224:THR:HG23	2.27	0.59
1:A:222:ASP:OD1	1:A:224:THR:HG23	2.04	0.57
1:A:25:GLU:OE2	1:A:303:ARG:HD3	2.07	0.55
1:A:351:LYS:HE2	1:A:369:ASN:O	2.07	0.55
1:A:183:ARG:HG2	1:A:184:ALA:N	2.22	0.53
1:A:271:LEU:HD22	1:A:298:GLY:HA2	1.91	0.52
1:A:44:ARG:HB3	1:A:65:ARG:HD2	1.91	0.52
1:A:257:VAL:HA	1:A:262:ARG:O	2.09	0.52
1:A:287:ARG:HH11	1:A:287:ARG:CG	2.23	0.50
1:A:196:TYR:O	1:A:200:GLN:HG2	2.13	0.49
1:A:192:PHE:HZ	1:A:350:PRO:HB3	1.77	0.49
1:A:114:ARG:NH1	1:A:238:LEU:HD21	2.28	0.48
1:A:287:ARG:HG2	1:A:287:ARG:NH1	2.27	0.47
1:A:388:ASP:OD2	1:A:390:ALA:HB3	2.15	0.46
1:A:313:TYR:CE2	1:A:315:GLU:HB2	2.51	0.46
1:A:185:ALA:HB1	1:A:189:LEU:HD23	1.98	0.45
1:A:214:VAL:HG23	1:A:219:LEU:HD11	1.98	0.45
1:A:394:ASP:OD1	1:A:395:PRO:HD2	2.17	0.45
1:A:145:GLN:HE21	1:A:145:GLN:HA	1.82	0.45
1:A:173:LEU:HB3	1:A:204:LEU:HD23	1.98	0.45
1:A:197:HIS:CE1	1:A:323:GLN:HE21	2.27	0.45
1:A:180:LEU:O	1:A:181:MET:C	2.60	0.45
1:A:327:ARG:H	1:A:369:ASN:ND2	2.16	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:ARG:NH2	3:A:464:HOH:O	2.34	0.44
1:A:388:ASP:HB3	1:A:391:ALA:HB3	2.00	0.44
1:A:375:ALA:HB3	1:A:376:PRO:CD	2.48	0.42
1:A:287:ARG:CG	1:A:287:ARG:NH1	2.81	0.42
1:A:48:ARG:N	1:A:49:PRO:CD	2.83	0.42
1:A:266:PRO:O	1:A:267:ASP:HB2	2.20	0.41
1:A:44:ARG:O	1:A:45:PRO:C	2.62	0.41
1:A:279:ASN:C	1:A:279:ASN:ND2	2.73	0.41
1:A:125:PRO:HD2	1:A:229:ASP:O	2.21	0.41
1:A:113:PRO:CG	1:A:130:ARG:HB2	2.50	0.41
1:A:375:ALA:HB3	1:A:376:PRO:HD3	2.03	0.40
1:A:375:ALA:N	1:A:376:PRO:HD2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/408 (98%)	381 (96%)	17 (4%)	1 (0%)	36	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	395	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	299/307 (97%)	273 (91%)	26 (9%)	9 7

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	32	GLN
1	A	44	ARG
1	A	70	ARG
1	A	75	GLU
1	A	120	GLN
1	A	145	GLN
1	A	183	ARG
1	A	208	ARG
1	A	225	ARG
1	A	271	LEU
1	A	279	ASN
1	A	286	GLU
1	A	287	ARG
1	A	303	ARG
1	A	309	THR
1	A	327	ARG
1	A	338	GLU
1	A	340	ILE
1	A	344	GLU
1	A	372	ARG
1	A	381	LEU
1	A	386	LYS
1	A	394	ASP
1	A	397	THR
1	A	399	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	GLN
1	A	145	GLN
1	A	197	HIS
1	A	240	ASN
1	A	279	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	ASN
1	A	323	GLN
1	A	329	GLN
1	A	358	GLN
1	A	369	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	449	-	58,58,58	2.18	14 (24%)	85,89,89	1.17	9 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	449	-	-	5/34/50/50	0/6/6/6

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	449	FAD	P-O3P	-10.08	1.48	1.59
2	A	449	FAD	C1'-C2'	4.04	1.58	1.52
2	A	449	FAD	PA-O2A	-3.96	1.37	1.55
2	A	449	FAD	O5'-C5'	3.62	1.58	1.44
2	A	449	FAD	C9A-N10	3.59	1.47	1.41
2	A	449	FAD	P-O2P	-3.39	1.39	1.55
2	A	449	FAD	C4X-N5	3.01	1.37	1.30
2	A	449	FAD	C8-C7	2.72	1.47	1.40
2	A	449	FAD	C5A-C6A	2.44	1.47	1.41
2	A	449	FAD	O2-C2	-2.25	1.19	1.24
2	A	449	FAD	C9A-C5X	2.25	1.44	1.41
2	A	449	FAD	C2-N3	2.23	1.43	1.39
2	A	449	FAD	C5A-C4A	2.08	1.42	1.39
2	A	449	FAD	P-O5'	-2.03	1.51	1.59

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	449	FAD	O2A-PA-O3P	2.70	114.58	107.27
2	A	449	FAD	O5B-PA-O1A	-2.54	98.86	108.94
2	A	449	FAD	C2A-N1A-C6A	2.49	122.82	118.73
2	A	449	FAD	O4B-C1B-C2B	-2.37	101.54	106.62
2	A	449	FAD	O2A-PA-O1A	2.09	122.19	112.44
2	A	449	FAD	C5'-C4'-C3'	-2.07	108.31	112.22
2	A	449	FAD	C4B-O4B-C1B	-2.06	104.93	109.47
2	A	449	FAD	O5'-P-O1P	-2.05	100.80	108.94
2	A	449	FAD	C4-N3-C2	-2.04	122.03	125.64

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	449	FAD	PA-O3P-P-O5'
2	A	449	FAD	P-O3P-PA-O2A
2	A	449	FAD	C2B-C1B-N9A-C8A
2	A	449	FAD	P-O3P-PA-O1A
2	A	449	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.