



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:27 AM UTC

PDB ID : 4D75 / pdb_00004d75
Title : Cytochrome P450 3A4 bound to an inhibitor
Authors : Sevrioukova, I.; Poulos, T.
Deposited on : 2014-11-19
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME P450 3A4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	460	3711	2415	613	659	24	0	1	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	expression tag	UNP P08684
A	22	ALA	-	expression tag	UNP P08684
A	504	HIS	-	expression tag	UNP P08684
A	505	HIS	-	expression tag	UNP P08684
A	506	HIS	-	expression tag	UNP P08684
A	507	HIS	-	expression tag	UNP P08684
A	282	ALA	LYS	engineered mutation	UNP P08684
A	285	ALA	GLU	engineered mutation	UNP P08684

- Molecule 2 is tert-butyl {6-oxo-6-[(pyridin-3-ylmethyl)amino]hexyl}carbamate (CCD ID: PK9) (formula: C₁₇H₂₇N₃O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			23	17	3	3		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	17	Total O 17 17	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYTOCHROME P450 3A4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.09Å 99.75Å 131.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	79.46 – 2.25 79.46 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.7 (79.46-2.25) 89.4 (79.46-2.25)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.43 (at 2.25Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.200 , 0.254 0.212 , 0.262	Depositor DCC
R_{free} test set	1247 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.419	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3794	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, PK9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	1/3805 (0.0%)	1.04	14/5145 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	184	ILE	C-N	-5.01	1.27	1.33

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	LYS	N-CA-C	8.81	129.57	110.80
1	A	211	LEU	N-CA-C	-7.43	98.63	110.14
1	A	205	GLU	CA-C-N	-7.38	110.94	122.67
1	A	205	GLU	C-N-CA	-7.38	110.94	122.67
1	A	257	LYS	CA-C-N	7.22	129.83	120.44
1	A	257	LYS	C-N-CA	7.22	129.83	120.44
1	A	141	LYS	CG-CD-CE	-7.18	94.79	111.30
1	A	141	LYS	CA-CB-CG	-6.12	101.86	114.10
1	A	170	VAL	N-CA-C	5.88	116.40	108.17
1	A	208	LYS	N-CA-C	-5.85	104.64	113.89
1	A	470	GLU	N-CA-C	5.50	117.27	111.28
1	A	238	ILE	CB-CA-C	5.46	119.81	111.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	VAL	CB-CA-C	-5.30	108.66	113.70
1	A	209	LYS	CB-CA-C	-5.05	100.37	110.42

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	206	ASN	Peptide
1	A	207	THR	Peptide
1	A	208	LYS	Peptide
1	A	209	LYS	Peptide
1	A	446	ARG	Sidechain
1	A	470	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3711	0	3797	68	1
2	A	23	0	27	8	0
3	A	43	0	30	2	0
4	A	17	0	0	0	0
All	All	3794	0	3854	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:GLU:HB3	1:A:259:SER:C	1.95	0.91
1:A:258:GLU:N	1:A:259:SER:HA	1.93	0.82
1:A:209:LYS:HB3	1:A:210:LEU:O	1.80	0.81
1:A:209:LYS:CE	1:A:303:ILE:HG21	2.13	0.78
1:A:243:ARG:HA	1:A:246:THR:OG1	1.83	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:PHE:HE2	2:A:600:PK9:H031	1.50	0.76
1:A:209:LYS:HE2	1:A:303:ILE:HG21	1.68	0.73
1:A:212:ARG:HE	1:A:308:GLU:HG2	1.54	0.72
1:A:258:GLU:HB3	1:A:260:ARG:N	2.04	0.71
1:A:181:MET:HE2	1:A:185:THR:HG23	1.76	0.68
1:A:184:ILE:HG21	1:A:303:ILE:HG23	1.78	0.65
1:A:468:CYS:SG	1:A:469:LYS:N	2.69	0.65
1:A:339:LEU:HB3	1:A:343:ALA:HB3	1.80	0.64
1:A:249:LEU:O	1:A:253:VAL:HG23	1.99	0.61
1:A:58:CYS:HB3	1:A:399:TYR:CD1	2.34	0.61
1:A:252:SER:O	1:A:256:MET:HG3	2.00	0.60
2:A:600:PK9:H101	2:A:600:PK9:H033	1.84	0.59
1:A:209:LYS:CE	1:A:303:ILE:CG2	2.81	0.58
1:A:209:LYS:HE3	1:A:303:ILE:CG2	2.33	0.58
1:A:132:LEU:HD21	1:A:288:LYS:HA	1.84	0.58
1:A:259:SER:O	1:A:262:GLU:HB3	2.03	0.58
1:A:269:VAL:HG23	1:A:269:VAL:O	2.03	0.57
1:A:273:GLN:HA	1:A:276:ILE:HG12	1.86	0.56
1:A:184:ILE:HG22	1:A:185:THR:N	2.20	0.56
1:A:475:LEU:HD11	1:A:485:PRO:HB3	1.86	0.56
1:A:209:LYS:HE3	1:A:303:ILE:HG21	1.90	0.54
2:A:600:PK9:H20	2:A:600:PK9:H013	1.90	0.54
1:A:145:MET:O	1:A:149:ILE:HG13	2.09	0.53
1:A:183:VAL:O	1:A:187:THR:OG1	2.11	0.53
1:A:258:GLU:N	1:A:259:SER:CA	2.71	0.53
1:A:247:ASN:O	1:A:248:PHE:C	2.53	0.51
1:A:270:ASP:N	1:A:270:ASP:OD1	2.43	0.51
1:A:136:THR:HG21	1:A:271:PHE:HD1	1.75	0.50
1:A:209:LYS:HE3	1:A:303:ILE:HB	1.93	0.50
1:A:136:THR:CG2	1:A:271:PHE:CD1	2.95	0.50
1:A:179:TYR:CZ	1:A:455:ALA:HB2	2.47	0.50
1:A:490:VAL:O	1:A:491:LEU:HD23	2.12	0.50
1:A:136:THR:CG2	1:A:271:PHE:HD1	2.25	0.50
1:A:209:LYS:HG2	1:A:210:LEU:HG	1.94	0.50
1:A:207:THR:CG2	1:A:209:LYS:HE2	2.41	0.49
1:A:246:THR:HG22	1:A:300:ILE:HD13	1.96	0.48
1:A:71:VAL:HG21	1:A:386:MET:HE1	1.95	0.48
1:A:136:THR:HG22	1:A:271:PHE:CE1	2.49	0.48
1:A:371:MET:HE1	1:A:481:GLY:O	2.13	0.48
1:A:136:THR:HG22	1:A:271:PHE:CD1	2.49	0.47
1:A:209:LYS:HE3	1:A:303:ILE:CB	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ARG:C	1:A:257:LYS:N	2.73	0.47
1:A:304:PHE:CE2	2:A:600:PK9:H031	2.40	0.47
1:A:305:ALA:HB1	2:A:600:PK9:H21	1.97	0.46
1:A:383:ILE:O	1:A:383:ILE:HG13	2.15	0.46
2:A:600:PK9:H032	2:A:600:PK9:H19	1.97	0.45
1:A:209:LYS:HB3	1:A:210:LEU:C	2.41	0.45
1:A:299:SER:O	1:A:303:ILE:HG13	2.15	0.45
2:A:600:PK9:H032	2:A:600:PK9:C19	2.46	0.45
1:A:305:ALA:HA	2:A:600:PK9:C20	2.47	0.45
1:A:275:MET:HE1	1:A:298:GLN:HB2	1.99	0.45
1:A:59:MET:O	1:A:63:GLU:HG3	2.17	0.44
1:A:255:ARG:C	1:A:257:LYS:H	2.24	0.44
3:A:601:HEM:HMC2	3:A:601:HEM:HBC2	2.00	0.44
1:A:267:HIS:O	1:A:268:ARG:HB2	2.18	0.43
1:A:188:SER:HA	1:A:271:PHE:HB2	2.00	0.43
1:A:209:LYS:HZ3	1:A:209:LYS:HG3	1.58	0.43
1:A:280:ASN:OD1	1:A:280:ASN:C	2.62	0.42
1:A:319:TYR:CZ	1:A:475:LEU:HB2	2.54	0.42
1:A:126:TRP:CZ2	1:A:440:ARG:HD2	2.54	0.42
1:A:258:GLU:H	1:A:259:SER:HA	1.79	0.42
1:A:94:LEU:HD21	1:A:396:ILE:HD13	2.02	0.42
1:A:272:LEU:O	1:A:276:ILE:HG23	2.19	0.42
1:A:260:ARG:C	1:A:262:GLU:H	2.29	0.41
1:A:181:MET:HE3	1:A:181:MET:HA	2.02	0.41
1:A:293:LEU:C	1:A:293:LEU:HD23	2.46	0.40
1:A:386:MET:HE3	1:A:386:MET:HB2	1.87	0.40
1:A:211:LEU:HD22	1:A:308:GLU:OE1	2.22	0.40
3:A:601:HEM:HMB2	3:A:601:HEM:HBB2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:LYS:NZ	1:A:168:LYS:NZ[2_445]	1.76	0.44

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/487 (93%)	417 (92%)	34 (8%)	4 (1%)	14	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	288	LYS
1	A	209	LYS
1	A	210	LEU
1	A	405	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	422/441 (96%)	402 (95%)	20 (5%)	23	26

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	111	VAL
1	A	162	ARG
1	A	184	ILE
1	A	207	THR
1	A	209	LYS
1	A	221	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	235	VAL
1	A	244	GLU
1	A	246	THR
1	A	260	ARG
1	A	262	GLU
1	A	270	ASP
1	A	288	LYS
1	A	293	LEU
1	A	382	GLU
1	A	421	LYS
1	A	424	LYS
1	A	459	VAL
1	A	492	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	ASN
1	A	451	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PK9	A	600	3	23,23,23	1.47	2 (8%)	29,29,29	1.48	3 (10%)
3	HEM	A	601	2,1	50,50,50	1.73	7 (14%)	67,82,82	1.39	10 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PK9	A	600	3	-	12/19/19/19	0/1/1/1
3	HEM	A	601	2,1	-	1/14/54/54	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	HEM	C3D-C2D	7.47	1.52	1.36
2	A	600	PK9	O05-C06	6.44	1.47	1.34
3	A	601	HEM	FE-NC	3.47	2.06	1.95
3	A	601	HEM	CAC-C3C	3.38	1.56	1.47
3	A	601	HEM	CAB-C3B	3.01	1.55	1.47
3	A	601	HEM	FE-NA	2.83	2.04	1.95
3	A	601	HEM	CMC-C2C	2.67	1.56	1.50
3	A	601	HEM	CMA-C3A	2.51	1.55	1.50
2	A	600	PK9	O05-C02	-2.05	1.44	1.48

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	PK9	O05-C06-N08	5.11	117.25	110.00
3	A	601	HEM	C4D-ND-C1D	4.54	110.58	105.21
3	A	601	HEM	CHD-C1D-ND	3.25	127.92	124.42
2	A	600	PK9	O05-C06-O07	-3.23	119.91	125.64
3	A	601	HEM	CBD-CAD-C3D	-2.61	105.31	112.53
3	A	601	HEM	CMB-C2B-C1B	-2.41	121.27	125.03
3	A	601	HEM	C3B-C2B-C1B	2.33	108.16	106.41
2	A	600	PK9	C21-N22-C23	2.29	120.86	116.85
3	A	601	HEM	CHA-C4D-ND	2.28	127.19	124.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	HEM	CHB-C1B-NB	2.25	127.14	124.37
3	A	601	HEM	CAD-C3D-C4D	2.22	128.56	124.70
3	A	601	HEM	O2A-CGA-CBA	2.13	120.72	114.00
3	A	601	HEM	CHC-C4B-NB	2.07	126.65	124.42

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600	PK9	C13-C14-N16-C17
2	A	600	PK9	C10-C09-N08-C06
2	A	600	PK9	O05-C06-N08-C09
2	A	600	PK9	N08-C06-O05-C02
2	A	600	PK9	O07-C06-O05-C02
2	A	600	PK9	O15-C14-N16-C17
2	A	600	PK9	O07-C06-N08-C09
2	A	600	PK9	C12-C13-C14-N16
2	A	600	PK9	C12-C13-C14-O15
2	A	600	PK9	N08-C09-C10-C11
2	A	600	PK9	C09-C10-C11-C12
2	A	600	PK9	C10-C11-C12-C13
3	A	601	HEM	CAA-CBA-CGA-O2A


There are no ring outliers.

2 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	PK9	8	0
3	A	601	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand PK9 A 600



Ligand HEM A 601



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	460/487 (94%)	0.70	35 (7%) 20 18	39, 69, 119, 171	1 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	261	LEU	5.9
1	A	191	VAL	4.3
1	A	211	LEU	4.2
1	A	144	GLU	4.2
1	A	209	LYS	4.1
1	A	212	ARG	3.6
1	A	213	PHE	3.6
1	A	269	VAL	3.3
1	A	210	LEU	3.2
1	A	226	PHE	3.1
1	A	203	PHE	3.0
1	A	308	GLU	3.0
1	A	276	ILE	2.9
1	A	267	HIS	2.9
1	A	202	PRO	2.9
1	A	222	SER	2.9
1	A	207	THR	2.8
1	A	259	SER	2.8
1	A	490	VAL	2.7
1	A	208	LYS	2.7
1	A	46	PHE	2.7
1	A	141	LYS	2.7
1	A	192	ASN	2.7
1	A	293	LEU	2.7
1	A	307	TYR	2.4
1	A	201	ASP	2.4
1	A	270	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	215	PHE	2.3
1	A	471	THR	2.3
1	A	304	PHE	2.3
1	A	422	LYS	2.2
1	A	204	VAL	2.2
1	A	257	LYS	2.1
1	A	260	ARG	2.1
1	A	274	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PK9	A	600	23/23	0.82	0.19	56,95,104,107	0
3	HEM	A	601	43/43	0.98	0.07	36,44,50,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PK9 A 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.