



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:18 AM UTC

PDB ID : 5D7G / pdb_00005d7g
Title : Structure of human ATG5 E122D-ATG16L1 complex at 3.0 Angstroms
Authors : Qiu, Y.; Schulman, B.A.
Deposited on : 2015-08-13
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	80% 14% 6%
1	C	280	80% 13% 6%
1	E	280	72% 20% 7%
1	G	280	69% 22% 9%
2	B	71	45% 10% 45%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	71	49%7%44%
2	F	71	46%10%44%
2	H	71	41%13%44%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Autophagy protein 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2033	1330	331	360	12			
1	C	263	Total	C	N	O	S	0	0	0
			2060	1342	341	365	12			
1	E	260	Total	C	N	O	S	0	0	0
			2052	1334	336	370	12			
1	G	256	Total	C	N	O	S	0	0	0
			2011	1313	329	357	12			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q9H1Y0
A	-3	ALA	-	expression tag	UNP Q9H1Y0
A	-2	MET	-	expression tag	UNP Q9H1Y0
A	-1	GLY	-	expression tag	UNP Q9H1Y0
A	0	SER	-	expression tag	UNP Q9H1Y0
A	122	ASP	GLU	engineered mutation	UNP Q9H1Y0
C	-4	GLY	-	expression tag	UNP Q9H1Y0
C	-3	ALA	-	expression tag	UNP Q9H1Y0
C	-2	MET	-	expression tag	UNP Q9H1Y0
C	-1	GLY	-	expression tag	UNP Q9H1Y0
C	0	SER	-	expression tag	UNP Q9H1Y0
C	122	ASP	GLU	engineered mutation	UNP Q9H1Y0
E	-4	GLY	-	expression tag	UNP Q9H1Y0
E	-3	ALA	-	expression tag	UNP Q9H1Y0
E	-2	MET	-	expression tag	UNP Q9H1Y0
E	-1	GLY	-	expression tag	UNP Q9H1Y0
E	0	SER	-	expression tag	UNP Q9H1Y0
E	122	ASP	GLU	engineered mutation	UNP Q9H1Y0
G	-4	GLY	-	expression tag	UNP Q9H1Y0
G	-3	ALA	-	expression tag	UNP Q9H1Y0
G	-2	MET	-	expression tag	UNP Q9H1Y0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-1	GLY	-	expression tag	UNP Q9H1Y0
G	0	SER	-	expression tag	UNP Q9H1Y0
G	122	ASP	GLU	engineered mutation	UNP Q9H1Y0

- Molecule 2 is a protein called Autophagy-related protein 16-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	39	Total	C	N	O	0	0	0
			295	186	58	51			
2	D	40	Total	C	N	O	0	0	0
			333	209	64	60			
2	F	40	Total	C	N	O	0	0	0
			321	199	65	57			
2	H	40	Total	C	N	O	0	0	0
			311	198	55	58			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP Q676U5
B	0	SER	-	expression tag	UNP Q676U5
D	-1	GLY	-	expression tag	UNP Q676U5
D	0	SER	-	expression tag	UNP Q676U5
F	-1	GLY	-	expression tag	UNP Q676U5
F	0	SER	-	expression tag	UNP Q676U5
H	-1	GLY	-	expression tag	UNP Q676U5
H	0	SER	-	expression tag	UNP Q676U5


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Autophagy protein 5



- Molecule 1: Autophagy protein 5




- Molecule 1: Autophagy protein 5




- Molecule 1: Autophagy protein 5

Chain G:  69% 22% 9%




- Molecule 2: Autophagy-related protein 16-1

Chain B:  45% 10% 45%




- Molecule 2: Autophagy-related protein 16-1

Chain D:  49% 7% 44%





- Molecule 2: Autophagy-related protein 16-1

Chain F:  46% 10% 44%



- Molecule 2: Autophagy-related protein 16-1

Chain H:  41% 13% 44%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	217.11Å 84.48Å 151.85Å 90.00° 133.81° 90.00°	Depositor
Resolution (Å)	49.53 – 3.00 49.53 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (49.53-3.00) 92.5 (49.53-3.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.198 , 0.244 0.204 , 0.247	Depositor DCC
R_{free} test set	1921 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	87.5	Xtriage
Anisotropy	0.647	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 65.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for h+2*k,-h-l 0.015 for h,-k,-h-l 0.019 for -h-2*k,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9417	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2098	0.79	0/2873
1	C	0.49	0/2126	0.82	2/2912 (0.1%)
1	E	0.48	0/2117	0.83	1/2897 (0.0%)
1	G	0.49	0/2075	0.79	0/2835
2	B	0.44	0/300	0.92	2/408 (0.5%)
2	D	0.42	0/338	0.93	2/455 (0.4%)
2	F	0.46	0/327	0.79	0/441
2	H	0.46	0/316	0.83	0/429
All	All	0.48	0/9697	0.82	7/13250 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	10	PHE	CA-C-N	7.25	128.23	120.11
2	D	10	PHE	C-N-CA	7.25	128.23	120.11
1	C	260	TYR	N-CA-C	-6.17	103.04	110.13
2	B	10	PHE	CA-C-N	-5.83	113.58	120.11
2	B	10	PHE	C-N-CA	-5.83	113.58	120.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2033	0	1852	31	0
1	C	2060	0	1890	21	0
1	E	2052	0	1882	37	0
1	G	2011	0	1845	40	0
2	B	295	0	252	7	0
2	D	333	0	315	3	0
2	F	321	0	287	5	0
2	H	311	0	278	10	0
3	C	1	0	0	0	0
All	All	9417	0	8601	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:71:GLU:OE1	1:G:103:HIS:HE1	1.50	0.93
1:E:85:LEU:HD13	1:E:266:HIS:HB3	1.63	0.81
2:H:24:ARG:NH1	2:H:25:ASP:OD1	2.21	0.73
1:G:71:GLU:OE1	1:G:103:HIS:CE1	2.41	0.70
1:G:243:ILE:HD13	2:H:21:LEU:HG	1.79	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/280 (92%)	248 (96%)	9 (4%)	0	100	100
1	C	259/280 (92%)	253 (98%)	6 (2%)	0	100	100
1	E	254/280 (91%)	246 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	248/280 (89%)	245 (99%)	2 (1%)	1 (0%)	30	65
2	B	37/71 (52%)	36 (97%)	1 (3%)	0	100	100
2	D	38/71 (54%)	38 (100%)	0	0	100	100
2	F	38/71 (54%)	38 (100%)	0	0	100	100
2	H	38/71 (54%)	38 (100%)	0	0	100	100
All	All	1169/1404 (83%)	1142 (98%)	26 (2%)	1 (0%)	48	80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	271	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/256 (79%)	201 (100%)	1 (0%)	81	89
1	C	208/256 (81%)	202 (97%)	6 (3%)	37	70
1	E	211/256 (82%)	206 (98%)	5 (2%)	43	73
1	G	205/256 (80%)	198 (97%)	7 (3%)	32	66
2	B	24/64 (38%)	24 (100%)	0	100	100
2	D	33/64 (52%)	33 (100%)	0	100	100
2	F	30/64 (47%)	30 (100%)	0	100	100
2	H	29/64 (45%)	26 (90%)	3 (10%)	7	28
All	All	942/1280 (74%)	920 (98%)	22 (2%)	44	74

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	61	ARG
1	G	152	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	101	THR
1	G	211	GLN
1	C	272	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	30	GLN
1	G	237	GLN
1	E	252	GLN
1	E	158	GLN
1	E	257	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/280 (93%)	-0.10	2 (0%) 82 64	60, 89, 133, 159	0
1	C	263/280 (93%)	-0.11	2 (0%) 82 64	55, 88, 127, 161	0
1	E	260/280 (92%)	-0.14	2 (0%) 82 64	57, 92, 145, 165	0
1	G	256/280 (91%)	-0.12	0 100 100	65, 91, 136, 160	0
2	B	39/71 (54%)	-0.12	0 100 100	68, 95, 127, 136	0
2	D	40/71 (56%)	-0.04	0 100 100	59, 94, 130, 133	0
2	F	40/71 (56%)	-0.07	0 100 100	62, 95, 155, 163	0
2	H	40/71 (56%)	0.09	0 100 100	65, 93, 139, 149	0
All	All	1201/1404 (85%)	-0.11	6 (0%) 87 72	55, 91, 136, 165	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	62	GLN	3.1
1	C	227	ILE	2.6
1	A	197	PRO	2.5
1	A	2	THR	2.3
1	C	64	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.