



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 23, 2026 – 09:56 AM UTC

PDB ID : 7D7K / pdb_00007d7k
Title : The crystal structure of SARS-CoV-2 papain-like protease in apo form
Authors : Zhao, Y.; Sun, L.; Yang, H.T.; Rao, Z.H.
Deposited on : 2020-10-04
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	2% 86% 13% .
1	B	312	15% 76% 23% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-structural protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	0	3	0
			2490	1586	409	477	18			
1	B	312	Total	C	N	O	S	0	2	0
			2428	1548	397	467	16			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	111	SER	CYS	engineered mutation	UNP P0DTD1
B	111	SER	CYS	engineered mutation	UNP P0DTD1

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CAFFEINE (CCD ID: CFF) (formula: C₈H₁₀N₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	4	2		
3	B	1	Total	C	N	O	0	0
			14	8	4	2		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	327	Total O 327 327	0	0
6	B	247	Total O 247 247	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Non-structural protein 3



• Molecule 1: Non-structural protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	116.00Å 116.00Å 254.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.28 – 1.90 49.28 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.7 (49.28-1.90) 98.7 (49.28-1.90)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.183 , 0.207 0.185 , 0.206	Depositor DCC
R_{free} test set	2000 reflections (2.48%)	wwPDB-VP
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5599	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CFF, ZN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.93	15/2555 (0.6%)	0.82	0/3468
1	B	1.18	27/2491 (1.1%)	0.90	8/3389 (0.2%)
All	All	1.06	42/5046 (0.8%)	0.86	8/6857 (0.1%)

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	276	ILE	C-O	-8.22	1.15	1.24
1	A	168	THR	C-O	-7.68	1.15	1.24
1	B	258	PHE	C-O	-7.35	1.15	1.23
1	A	279	LYS	C-O	-7.26	1.17	1.23
1	B	279	LYS	C-O	-7.10	1.17	1.23
1	B	300	ILE	C-O	-6.96	1.15	1.24
1	B	265	THR	C-O	-6.75	1.16	1.24
1	A	126	LYS	C-O	-6.51	1.16	1.24
1	A	290	LEU	C-O	-6.45	1.16	1.24
1	B	190	LYS	C-O	-6.38	1.16	1.24
1	B	306	LYS	C-O	-6.37	1.16	1.23
1	A	309	SER	C-O	-6.34	1.16	1.23
1	B	278	SER	C-O	-6.28	1.16	1.24
1	B	303	VAL	C-O	-6.27	1.17	1.24
1	B	264	TYR	C-O	-6.22	1.16	1.24
1	A	65	ARG	C-O	-6.19	1.16	1.24
1	B	140	ARG	C-O	-6.06	1.16	1.24
1	B	228	LYS	N-CA	-6.01	1.38	1.45
1	B	141	ALA	C-O	-5.94	1.16	1.24
1	A	82	ARG	C-O	-5.91	1.17	1.24
1	B	229	GLN	N-CA	-5.77	1.38	1.46
1	B	90	THR	C-O	-5.69	1.17	1.24
1	A	68	ALA	C-O	-5.69	1.17	1.24
1	B	254	LYS	C-O	-5.68	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	289	LEU	C-O	-5.67	1.16	1.23
1	B	284	CYS	C-O	-5.61	1.17	1.24
1	A	170	SER	C-O	-5.55	1.17	1.24
1	A	83	TYR	C-O	-5.55	1.17	1.24
1	B	255	HIS	C-O	-5.54	1.17	1.23
1	B	99	ASN	C-O	-5.54	1.16	1.23
1	B	122	GLN	C-O	-5.51	1.16	1.24
1	B	145	ALA	C-O	-5.50	1.17	1.24
1	B	253	LEU	C-O	-5.48	1.17	1.24
1	B	188	VAL	C-O	-5.45	1.18	1.24
1	A	308	ASN	C-O	-5.34	1.17	1.24
1	A	288	ALA	C-O	-5.34	1.17	1.24
1	B	142	GLY	C-O	-5.33	1.17	1.24
1	A	124	GLU	C-O	-5.29	1.17	1.24
1	B	282	LEU	C-O	-5.25	1.17	1.24
1	A	66	VAL	C-O	-5.14	1.18	1.24
1	B	256	GLY	C-O	-5.06	1.16	1.24
1	B	143	GLU	C-O	-5.04	1.17	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	224	CYS	N-CA-C	-6.17	99.86	109.72
1	B	224	CYS	CA-C-N	6.15	129.14	120.28
1	B	224	CYS	C-N-CA	6.15	129.14	120.28
1	B	144	ALA	N-CA-C	6.06	120.67	113.16
1	B	188	VAL	N-CA-C	5.76	116.16	107.75
1	B	229	GLN	N-CA-C	-5.58	100.20	109.46
1	B	50	HIS	N-CA-C	-5.02	100.11	110.80
1	B	192	CYS	N-CA-C	5.01	119.03	113.01

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2490	0	2435	29	0
1	B	2428	0	2322	41	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	14	0	10	0	0
3	B	14	0	10	0	0
4	A	28	0	42	5	0
4	B	24	0	36	3	0
5	A	15	0	0	1	0
5	B	10	0	0	0	0
6	A	327	0	0	2	0
6	B	247	0	0	1	0
All	All	5599	0	4855	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:VAL:HG23	1:B:227:GLY:HA3	1.59	0.85
1:A:194:GLN:H	4:A:404:EDO:H11	1.48	0.78
1:B:23:MET:HA	1:B:46:PRO:HG2	1.72	0.71
1:B:38:GLY:HA2	1:B:84:MET:HE3	1.75	0.68
1:B:105:LYS:NZ	6:B:502:HOH:O	2.29	0.64
1:B:240:PRO:HA	1:B:307:GLU:O	1.96	0.64
1:A:66:VAL:CG2	1:B:227:GLY:HA3	2.26	0.64
1:B:38:GLY:H	1:B:84:MET:HE1	1.68	0.59
1:A:4:THR:HG23	6:A:749:HOH:O	2.02	0.59
1:A:17:HIS:ND1	4:A:406:EDO:H21	2.18	0.58
1:B:13:ASN:HB2	1:B:56:TYR:OH	2.02	0.58
1:B:61:ASP:HB2	1:B:64:LEU:H	1.68	0.58
1:A:74:THR:OG1	4:A:407:EDO:H11	2.04	0.58
1:A:140:ARG:HH21	4:A:409:EDO:H21	1.69	0.57
1:B:33:PRO:HB2	1:B:58:LEU:HD23	1.86	0.57
1:B:35:TYR:CB	1:B:84:MET:HE2	2.36	0.55
1:A:66:VAL:HG23	1:B:227:GLY:CA	2.35	0.55
1:A:188:VAL:HG22	1:A:194:GLN:HG2	1.89	0.55
1:B:4:THR:N	1:B:21:VAL:O	2.40	0.55
1:B:239:SER:O	1:B:307:GLU:O	2.25	0.55
1:B:8:PHE:CE1	1:B:18:THR:HG22	2.43	0.54
1:A:62:ASP:OD1	1:B:228:LYS:HA	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:TYR:HB2	1:B:84:MET:HE2	1.90	0.53
1:A:165:VAL:O	1:A:169:MET:HG2	2.08	0.53
1:B:8:PHE:HB2	1:B:54:THR:HA	1.90	0.52
1:B:164:ASP:OD2	4:B:405:EDO:H12	2.09	0.52
1:A:63:THR:O	1:A:67:GLU:HG3	2.10	0.52
1:A:89:HIS:HB2	1:A:159:VAL:HG21	1.90	0.52
1:B:38:GLY:H	1:B:84:MET:CE	2.23	0.50
1:A:125:LEU:C	1:A:125:LEU:HD23	2.36	0.50
1:B:115[A]:THR:HG23	1:B:275:HIS:HB2	1.94	0.49
1:A:252:GLU:OE2	1:A:254:LYS:NZ	2.40	0.49
1:A:166:ARG:HA	1:A:243:MET:HE1	1.95	0.48
1:A:133:GLN:NE2	1:A:137:TYR:HE2	2.11	0.48
1:B:28:GLY:HA3	1:B:42:THR:HG23	1.94	0.48
1:A:223:PRO:HB2	1:B:75:THR:HB	1.96	0.48
1:A:194:GLN:N	4:A:404:EDO:H11	2.22	0.47
1:B:263:GLU:O	1:B:273:TYR:HA	2.14	0.47
1:A:210[B]:THR:HG21	1:A:220:VAL:HG11	1.97	0.47
1:B:38:GLY:CA	1:B:84:MET:HE3	2.43	0.47
1:B:194:GLN:NE2	1:B:315:LYS:HE2	2.30	0.46
1:A:294:SER:N	5:A:411:SO4:O1	2.31	0.46
1:B:8:PHE:HE1	1:B:18:THR:HG22	1.79	0.46
1:B:243:MET:HE3	1:B:304:PHE:CZ	2.50	0.46
1:A:66:VAL:CG2	1:B:227:GLY:CA	2.92	0.46
1:B:136:TYR:CZ	4:B:403:EDO:H11	2.51	0.46
1:B:283:TYR:CD1	1:B:290:LEU:HD11	2.51	0.46
1:A:62:ASP:HA	1:B:227:GLY:O	2.16	0.45
1:A:252:GLU:HG2	1:A:254:LYS:HE2	1.98	0.44
1:B:33:PRO:HA	1:B:42:THR:OG1	2.17	0.44
1:B:263:GLU:OE1	1:B:299:PRO:HD2	2.19	0.43
1:A:126:LYS:HB2	1:A:177[B]:ASN:ND2	2.34	0.42
1:A:215:GLN:HG3	1:A:220:VAL:HG12	2.01	0.42
1:A:126:LYS:HG2	1:A:133:GLN:HE22	1.85	0.42
1:B:170:SER:O	1:B:174:GLN:HG2	2.19	0.42
1:A:192:CYS:SG	1:A:226:CYS:HB3	2.59	0.41
1:B:194:GLN:CD	1:B:315:LYS:HE2	2.45	0.41
1:B:34:THR:N	1:B:42:THR:OG1	2.45	0.41
1:B:8:PHE:HB2	1:B:53:LYS:O	2.20	0.41
1:A:215:GLN:HG2	6:A:653:HOH:O	2.20	0.41
1:B:294:SER:OG	4:B:406:EDO:H11	2.21	0.40
1:B:188:VAL:HG22	1:B:194:GLN:HG2	2.04	0.40
1:B:190:LYS:HE3	1:B:190:LYS:HB2	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/312 (100%)	306 (98%)	7 (2%)	0	100	100
1	B	312/312 (100%)	297 (95%)	14 (4%)	1 (0%)	36	29
All	All	625/624 (100%)	603 (96%)	21 (3%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	40	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/272 (101%)	274 (100%)	1 (0%)	84	87
1	B	258/272 (95%)	258 (100%)	0	100	100
All	All	533/544 (98%)	532 (100%)	1 (0%)	87	90

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	289	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	B	99	ASN
1	B	122	GLN
1	B	186	ASN
1	B	194	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 2 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	B	410	-	3,3,3	0.53	0	2,2,2	0.37	0
4	EDO	B	404	-	3,3,3	0.45	0	2,2,2	0.20	0
4	EDO	A	407	-	3,3,3	0.68	0	2,2,2	0.36	0
5	SO4	B	409	-	4,4,4	0.37	0	6,6,6	0.45	0
4	EDO	B	406	-	3,3,3	0.52	0	2,2,2	0.63	0
4	EDO	A	403	-	3,3,3	0.53	0	2,2,2	0.43	0
4	EDO	A	409	-	3,3,3	0.50	0	2,2,2	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	406	-	3,3,3	0.60	0	2,2,2	0.11	0
3	CFF	B	402	-	15,15,15	2.42	5 (33%)	23,23,23	2.03	8 (34%)
4	EDO	B	407	-	3,3,3	0.61	0	2,2,2	0.43	0
4	EDO	A	408	-	3,3,3	0.88	0	2,2,2	0.30	0
4	EDO	A	404	-	3,3,3	0.45	0	2,2,2	0.64	0
5	SO4	A	411	-	4,4,4	0.25	0	6,6,6	0.47	0
5	SO4	B	408	-	4,4,4	0.43	0	6,6,6	0.66	0
3	CFF	A	402	-	15,15,15	2.49	4 (26%)	23,23,23	2.16	8 (34%)
4	EDO	B	403	-	3,3,3	0.69	0	2,2,2	0.24	0
5	SO4	A	410	-	4,4,4	0.49	0	6,6,6	0.44	0
4	EDO	B	405	-	3,3,3	0.54	0	2,2,2	0.25	0
5	SO4	A	412	-	4,4,4	0.32	0	6,6,6	0.22	0
4	EDO	A	405	-	3,3,3	0.57	0	2,2,2	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	410	-	-	0/1/1/1	-
4	EDO	B	406	-	-	1/1/1/1	-
4	EDO	A	403	-	-	0/1/1/1	-
4	EDO	B	404	-	-	0/1/1/1	-
4	EDO	A	409	-	-	1/1/1/1	-
4	EDO	A	406	-	-	1/1/1/1	-
4	EDO	A	405	-	-	0/1/1/1	-
4	EDO	A	407	-	-	0/1/1/1	-
3	CFF	A	402	-	-	-	0/2/2/2
4	EDO	B	403	-	-	1/1/1/1	-
3	CFF	B	402	-	-	-	0/2/2/2
4	EDO	B	405	-	-	1/1/1/1	-
4	EDO	A	408	-	-	0/1/1/1	-
4	EDO	A	404	-	-	1/1/1/1	-
4	EDO	B	407	-	-	0/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	CFF	C4-N3	5.47	1.47	1.38
3	B	402	CFF	C4-N3	5.09	1.46	1.38
3	A	402	CFF	C2-N3	4.81	1.48	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	CFF	C2-N3	4.39	1.47	1.39
3	B	402	CFF	C2-N1	4.29	1.47	1.39
3	A	402	CFF	C2-N1	4.14	1.47	1.39
3	B	402	CFF	C5-C6	3.61	1.52	1.43
3	A	402	CFF	C5-C6	3.41	1.51	1.43
3	B	402	CFF	O13-C6	-2.26	1.18	1.23

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	CFF	C10-N1-C2	4.78	125.67	117.33
3	B	402	CFF	C5-C6-N1	4.47	120.25	112.06
3	A	402	CFF	C5-C6-N1	3.76	118.96	112.06
3	A	402	CFF	N3-C4-N9	3.61	131.94	126.27
3	B	402	CFF	C6-N1-C2	-3.35	120.21	125.66
3	A	402	CFF	C12-N3-C2	3.29	123.06	117.33
3	A	402	CFF	C6-N1-C2	-3.27	120.34	125.66
3	B	402	CFF	C10-N1-C2	3.25	122.99	117.33
3	B	402	CFF	C12-N3-C2	2.89	122.37	117.33
3	B	402	CFF	N3-C4-N9	2.79	130.66	126.27
3	B	402	CFF	C6-C5-C4	-2.39	119.91	122.92
3	A	402	CFF	C10-N1-C6	-2.37	113.97	117.64
3	B	402	CFF	C12-N3-C4	-2.33	117.31	120.75
3	A	402	CFF	N3-C2-N1	2.31	120.02	117.14
3	A	402	CFF	O13-C6-N1	-2.08	116.57	120.36
3	B	402	CFF	O13-C6-C5	-2.04	122.21	126.38

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	406	EDO	O1-C1-C2-O2
4	A	406	EDO	O1-C1-C2-O2
4	A	409	EDO	O1-C1-C2-O2
4	B	405	EDO	O1-C1-C2-O2
4	A	404	EDO	O1-C1-C2-O2
4	B	403	EDO	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	407	EDO	1	0
4	B	406	EDO	1	0
4	A	409	EDO	1	0
4	A	406	EDO	1	0
4	A	404	EDO	2	0
5	A	411	SO4	1	0
4	B	403	EDO	1	0
4	B	405	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/312 (100%)	0.05	7 (2%) 62 66	20, 41, 62, 71	3 (0%)
1	B	312/312 (100%)	0.62	47 (15%) 5 5	22, 44, 102, 128	2 (0%)
All	All	624/624 (100%)	0.33	54 (8%) 16 17	20, 42, 86, 128	5 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	50	HIS	5.2
1	B	4	THR	5.1
1	B	22	ASP	4.9
1	B	5	ILE	4.7
1	B	55	PHE	4.7
1	B	7	VAL	4.6
1	B	47	HIS	4.5
1	B	227	GLY	4.3
1	B	36	LEU	4.3
1	B	20	VAL	4.3
1	B	21	VAL	4.2
1	B	46	PRO	4.2
1	A	125	LEU	4.2
1	B	25	MET	4.1
1	B	44	ILE	4.0
1	B	43	LYS	3.8
1	B	6	LYS	3.8
1	B	57	VAL	3.6
1	B	30	GLN	3.5
1	B	9	THR	3.4
1	B	27	TYR	3.4
1	B	226	CYS	3.3
1	B	39	ALA	3.2
1	B	52	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	8	PHE	2.9
1	B	18	THR	2.8
1	B	48	ASN	2.8
1	B	51	GLU	2.8
1	B	26	THR	2.8
1	B	268	TYR	2.8
1	A	137	TYR	2.7
1	B	23	MET	2.7
1	B	32	GLY	2.7
1	B	31	PHE	2.7
1	B	42	THR	2.5
1	B	35	TYR	2.5
1	B	56	TYR	2.4
1	B	61	ASP	2.4
1	B	24	SER	2.4
1	B	315	LYS	2.4
1	B	225	THR	2.3
1	B	54	THR	2.3
1	B	53	LYS	2.3
1	A	177[A]	ASN	2.3
1	A	180	SER	2.2
1	B	224	CYS	2.2
1	B	58	LEU	2.2
1	B	255	HIS	2.1
1	A	268	TYR	2.1
1	B	28	GLY	2.1
1	A	44	ILE	2.0
1	B	94	LYS	2.0
1	B	289[A]	LEU	2.0
1	A	179	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	409	4/4	0.71	0.15	71,74,74,81	0
5	SO4	A	412	5/5	0.72	0.10	100,100,108,130	0
5	SO4	A	411	5/5	0.75	0.09	99,99,103,113	0
4	EDO	B	406	4/4	0.77	0.15	65,72,73,78	0
5	SO4	B	409	5/5	0.80	0.10	66,77,108,115	0
4	EDO	B	407	4/4	0.82	0.15	58,60,64,75	0
4	EDO	A	407	4/4	0.85	0.14	61,62,63,67	0
4	EDO	A	408	4/4	0.85	0.16	51,53,54,55	0
4	EDO	A	406	4/4	0.88	0.14	50,59,62,68	0
4	EDO	A	404	4/4	0.88	0.11	60,65,67,68	0
5	SO4	A	410	5/5	0.88	0.10	43,59,80,82	0
3	CFF	A	402	14/14	0.91	0.10	30,36,40,40	14
4	EDO	B	405	4/4	0.91	0.11	56,56,59,66	0
4	EDO	A	405	4/4	0.92	0.14	44,49,53,55	0
4	EDO	B	410	4/4	0.92	0.12	49,51,54,64	0
4	EDO	B	403	4/4	0.93	0.10	41,44,47,50	0
5	SO4	B	408	5/5	0.94	0.08	49,54,68,75	0
4	EDO	A	403	4/4	0.95	0.08	46,48,51,53	0
3	CFF	B	402	14/14	0.95	0.10	33,38,43,43	14
4	EDO	B	404	4/4	0.95	0.11	53,58,60,67	0
2	ZN	B	401	1/1	0.97	0.05	45,45,45,45	1
2	ZN	A	401	1/1	0.99	0.03	49,49,49,49	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around ZN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.