



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:53 AM UTC

PDB ID : 1D8F / pdb_00001d8f
Title : CRYSTAL STRUCTURE OF MMP3 COMPLEXED WITH A PIPERAZINE
BASED INHIBITOR.
Authors : Cheng, M.Y.; De, B.; Pikul, S.; Almstead, N.G.; Natchus, M.G.; Anastasio,
M.V.; McPhail, S.J.; Snider, C.E.; Taiwo, Y.O.; Chen, L.Y.
Deposited on : 1999-10-22
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	173	 83% 14% ..
1	B	173	 78% 21% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2798 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STROMELYSIN-1 PRECURSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	169	Total	C	N	O	S	0	0	0
			1346	865	224	255	2			
1	B	173	Total	C	N	O	S	0	0	0
			1376	882	228	264	2			


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		
3	B	3	Total	Ca	0	0
			3	3		

- Molecule 4 is N-HYDROXY-1-(4-METHOXYPHENYL)SULFONYL-4-BENZYLOXYCARBONYL-PIPERAZINE-2-CARBOXAMIDE (CCD ID: SPI) (formula: C₂₀H₂₃N₃O₇S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			31	20	3	7	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	13	Total	O	0	0
			13	13		
5	B	22	Total	O	0	0
			22	22		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: STROMELYSIN-1 PRECURSOR

Chain A:  83% 14% ..



• Molecule 1: STROMELYSIN-1 PRECURSOR

Chain B:  78% 21% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.27Å 78.54Å 106.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.40	Depositor
% Data completeness (in resolution range)	97.7 (8.00-2.40)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.294 , 0.252	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2798	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN, SPI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/1390	1.10	9/1899 (0.5%)
1	B	0.54	0/1421	1.06	12/1941 (0.6%)
All	All	0.56	0/2811	1.08	21/3840 (0.5%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	602	VAL	N-CA-C	6.74	117.52	110.72
1	A	102	VAL	N-CA-C	6.66	117.42	110.62
1	A	158	ASP	N-CA-C	6.33	121.56	111.81
1	A	220	TYR	CA-C-N	6.22	125.92	119.82
1	A	220	TYR	C-N-CA	6.22	125.92	119.82
1	B	634	ARG	N-CA-C	5.88	119.22	109.46
1	B	728	ASP	N-CA-C	5.84	119.22	107.41
1	A	110	LYS	N-CA-C	5.82	118.40	111.71
1	B	715	THR	N-CA-C	5.80	118.39	111.71
1	B	729	LEU	N-CA-C	-5.73	105.95	113.17
1	B	630	LEU	N-CA-C	5.72	118.53	110.23
1	A	244	SER	N-CA-C	-5.60	105.33	111.82
1	A	188	LYS	N-CA-C	-5.57	106.32	113.23
1	A	247	GLY	CA-C-N	5.44	125.98	120.38
1	A	247	GLY	C-N-CA	5.44	125.98	120.38
1	B	607	ASP	N-CA-C	5.42	117.19	111.28
1	B	636	TYR	N-CA-C	-5.36	106.75	113.72
1	B	605	THR	CA-C-N	5.04	124.53	119.19
1	B	605	THR	C-N-CA	5.04	124.53	119.19
1	B	747	GLY	CA-C-N	5.03	125.56	120.38
1	B	747	GLY	C-N-CA	5.03	125.56	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1346	0	1270	14	0
1	B	1376	0	1295	15	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	B	31	0	22	1	0
5	A	13	0	0	0	0
5	B	22	0	0	0	0
All	All	2798	0	2587	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:698:VAL:O	1:B:702:GLU:HG2	1.97	0.65
1:A:89:ILE:O	1:A:89:ILE:HG22	1.99	0.61
1:B:687:THR:HG22	1:B:689:ASP:H	1.69	0.56
1:B:589:ILE:HG22	1:B:589:ILE:O	2.09	0.53
1:A:97:LEU:HD12	1:A:132:PHE:HE2	1.75	0.52
1:B:694:ASN:HD22	1:B:697:LEU:H	1.60	0.50
1:A:122:LYS:O	1:A:126:GLU:HG3	2.14	0.47
1:A:226:LEU:O	1:A:227:THR:HB	2.14	0.47
1:B:622:LYS:O	1:B:626:GLU:HB2	2.15	0.47
1:B:669:ALA:O	1:B:675:ASN:HB3	2.15	0.46
1:B:698:VAL:HG13	4:B:901:SPI:HC51	1.97	0.46
1:A:210:PHE:CG	1:A:211:HIS:N	2.84	0.45
1:A:190:THR:OG1	1:A:227:THR:HA	2.15	0.45
1:A:108:LEU:HD11	1:A:188:LYS:HG3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ASN:ND2	1:A:197:LEU:H	2.14	0.45
1:B:625:GLU:HG3	1:B:630:LEU:O	2.17	0.45
1:A:83:PHE:HA	1:A:212:SER:HB3	2.00	0.44
1:A:194:ASN:HD22	1:A:197:LEU:H	1.67	0.43
1:A:83:PHE:HA	1:A:212:SER:CB	2.49	0.43
1:B:635:LEU:HD12	1:B:640:ALA:HB2	2.02	0.42
1:B:667:ALA:HB3	1:B:705:HIS:HB2	2.01	0.41
1:B:667:ALA:HA	1:B:677:ASP:O	2.20	0.41
1:A:203:ILE:O	1:A:207:LEU:HG	2.21	0.41
1:B:621:LEU:HD23	1:B:703:ILE:HD13	2.03	0.41
1:A:89:ILE:N	1:A:90:PRO:HD3	2.36	0.41
1:B:710:PHE:CG	1:B:711:HIS:N	2.89	0.40
1:A:89:ILE:N	1:A:89:ILE:HD12	2.36	0.40
1:B:628:THR:HB	1:B:629:PRO:CD	2.52	0.40
1:B:687:THR:HG21	1:B:691:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	167/173 (96%)	156 (93%)	10 (6%)	1 (1%)	21	32
1	B	171/173 (99%)	162 (95%)	9 (5%)	0	100	100
All	All	338/346 (98%)	318 (94%)	19 (6%)	1 (0%)	36	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/147 (97%)	142 (99%)	1 (1%)	76	88
1	B	147/147 (100%)	142 (97%)	5 (3%)	32	54
All	All	290/294 (99%)	284 (98%)	6 (2%)	47	69

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	ARG
1	B	607	ASP
1	B	637	GLU
1	B	649	ARG
1	B	707	LEU
1	B	731	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	HIS
1	A	166	HIS
1	A	194	ASN
1	B	596	HIS
1	B	694	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 10 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SPI	B	901	2	32,33,33	2.98	18 (56%)	42,46,46	2.67	15 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SPI	B	901	2	-	4/29/42/42	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	901	SPI	C6-C5	5.18	1.47	1.38
4	B	901	SPI	S1-N1	4.72	1.70	1.63
4	B	901	SPI	C6-C1	4.66	1.46	1.38
4	B	901	SPI	C3-C2	4.62	1.46	1.38
4	B	901	SPI	C2-C1	4.38	1.45	1.38
4	B	901	SPI	C18-C19	4.31	1.46	1.38
4	B	901	SPI	O5-C12	-4.24	1.26	1.34
4	B	901	SPI	C16-C15	3.96	1.45	1.38
4	B	901	SPI	C18-C17	3.94	1.46	1.38
4	B	901	SPI	C17-C16	3.82	1.46	1.38
4	B	901	SPI	C3-C4	3.82	1.45	1.38
4	B	901	SPI	C5-C4	3.76	1.45	1.38
4	B	901	SPI	O4-C12	3.59	1.26	1.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	901	SPI	O3-C7	-3.20	1.33	1.42
4	B	901	SPI	C15-C14	3.15	1.45	1.38
4	B	901	SPI	C19-C14	3.02	1.44	1.38
4	B	901	SPI	C1-S1	2.68	1.80	1.76
4	B	901	SPI	O5-C13	-2.38	1.40	1.45

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	901	SPI	O2-S1-N1	-6.24	96.05	106.97
4	B	901	SPI	O1-S1-C1	6.05	115.62	108.10
4	B	901	SPI	C19-C14-C15	5.54	126.47	118.23
4	B	901	SPI	C18-C19-C14	-4.57	114.19	120.61
4	B	901	SPI	O2-S1-O1	-4.43	112.67	119.59
4	B	901	SPI	C7-O3-C4	4.39	126.91	117.50
4	B	901	SPI	O2-S1-C1	3.84	112.88	108.10
4	B	901	SPI	O5-C12-N2	3.83	118.46	111.77
4	B	901	SPI	C13-O5-C12	3.80	125.07	115.51
4	B	901	SPI	C1-S1-N1	3.54	113.53	107.36
4	B	901	SPI	C8-N1-S1	-3.18	111.45	119.33
4	B	901	SPI	C11-N1-C8	3.12	122.94	114.89
4	B	901	SPI	C17-C16-C15	-2.57	117.07	120.24
4	B	901	SPI	O4-C12-N2	-2.36	119.93	124.30
4	B	901	SPI	C10-C11-N1	2.18	111.38	109.00

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	901	SPI	C3-C4-O3-C7
4	B	901	SPI	C5-C4-O3-C7
4	B	901	SPI	NA-C20-C8-N1
4	B	901	SPI	OB-C20-C8-N1


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	901	SPI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.