



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:03 PM UTC

PDB ID : 5D8G / pdb_00005d8g
Title : A structural view on the dissociation of E. coli Tryptophanase
Authors : Almog, O.
Deposited on : 2015-08-17
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EPE	A	1003	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	0	6	0
			3730	2376	630	697	27			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	MET	VAL	conflict	UNP P0A853

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Mg	0	0
			3	3		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	422	Total	O	0	0
			422	422		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophanase



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	118.29Å 120.27Å 171.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.20 – 1.89 49.20 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.0 (49.20-1.89) 99.0 (49.20-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.174 , 0.214 0.173 , 0.213	Depositor DCC
R_{free} test set	2449 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4173	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CME, EPE, CL, MG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.53	24/3785 (0.6%)	1.23	12/5104 (0.2%)

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	222	ILE	CA-C	8.20	1.59	1.53
1	A	208	ALA	CA-CB	7.74	1.65	1.53
1	A	371	ILE	CA-CB	7.57	1.63	1.54
1	A	425	ALA	CA-CB	7.53	1.64	1.53
1	A	232	ALA	CA-CB	6.52	1.64	1.53
1	A	381	LEU	CA-C	6.41	1.61	1.52
1	A	84	ALA	CA-CB	6.20	1.62	1.53
1	A	397	GLY	N-CA	6.17	1.51	1.45
1	A	380	ALA	CA-CB	5.99	1.62	1.53
1	A	320	VAL	CA-CB	5.72	1.62	1.54
1	A	401	LEU	N-CA	5.61	1.53	1.46
1	A	434	PHE	C-O	5.59	1.30	1.24
1	A	260	TYR	N-CA	5.58	1.52	1.46
1	A	193	ALA	CA-CB	-5.51	1.46	1.53
1	A	17	GLU	CA-C	5.44	1.59	1.52
1	A	282	MET	N-CA	5.43	1.52	1.46
1	A	66	ALA	CA-CB	5.33	1.62	1.53
1	A	41	LEU	N-CA	5.24	1.52	1.45
1	A	112	VAL	N-CA	-5.21	1.40	1.46
1	A	217	ALA	CA-CB	5.12	1.61	1.53
1	A	191	ILE	CA-C	5.12	1.58	1.52
1	A	274	VAL	N-CA	5.10	1.50	1.46
1	A	66	ALA	N-CA	5.07	1.52	1.46
1	A	64	GLN	C-O	5.03	1.30	1.24

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	313	GLY	N-CA-C	-7.15	105.50	114.16
1	A	78	ARG	NE-CZ-NH1	-6.66	114.84	121.50
1	A	455	THR	N-CA-C	-6.34	106.14	114.31
1	A	457	GLU	CA-C-N	-5.81	114.28	120.03
1	A	457	GLU	C-N-CA	-5.81	114.28	120.03
1	A	78	ARG	NE-CZ-NH2	5.78	124.40	119.20
1	A	183	VAL	N-CA-C	5.69	116.47	110.72
1	A	448	ASN	N-CA-C	5.25	119.69	113.28
1	A	267	SER	N-CA-C	-5.22	101.46	109.76
1	A	155	ILE	N-CA-C	5.11	115.62	110.05
1	A	283	LYS	N-CA-C	5.06	116.49	110.97
1	A	255	ARG	NE-CZ-NH2	-5.04	114.67	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3730	0	3706	79	0
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	15	0	17	11	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	A	422	0	0	31	3
All	All	4173	0	3723	82	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:LYS:HE3	7:A:1103:HOH:O	0.90	1.07

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:419:ARG:HH22	4:A:1003:EPE:H61	1.30	0.96
1:A:133:ASN:HD21	1:A:194:THR:H	1.14	0.90
1:A:68[B]:MET:CG	7:A:1344:HOH:O	2.26	0.84
1:A:68[B]:MET:CE	7:A:1344:HOH:O	2.24	0.83
1:A:178:ARG:HD2	7:A:1228:HOH:O	1.78	0.83
1:A:441:HIS:HE1	7:A:1393:HOH:O	1.61	0.82
1:A:68[B]:MET:HG2	7:A:1344:HOH:O	1.79	0.82
1:A:240:GLN:OE1	7:A:1102:HOH:O	2.00	0.79
1:A:412:PRO:HA	7:A:1109:HOH:O	1.81	0.78
1:A:419:ARG:HH12	4:A:1003:EPE:H52	1.50	0.75
1:A:419:ARG:HH22	4:A:1003:EPE:H101	1.52	0.74
1:A:100:HIS:CD2	1:A:309:GLY:HA2	2.23	0.73
1:A:164:ARG:HG2	1:A:164:ARG:HH11	1.53	0.73
1:A:133:ASN:HD22	1:A:133:ASN:H	1.37	0.72
1:A:419:ARG:NH2	4:A:1003:EPE:H61	2.03	0.72
1:A:196:THR:OG1	1:A:358:HIS:HE1	1.73	0.71
1:A:419:ARG:NH1	4:A:1003:EPE:H52	2.06	0.71
1:A:170:ASN:HD21	1:A:206:SER:H	1.37	0.70
1:A:171:PHE:H	1:A:209:ASN:HD21	1.43	0.66
1:A:230:ARG:NH2	7:A:1103:HOH:O	2.12	0.64
1:A:164:ARG:HH11	1:A:164:ARG:CG	2.12	0.62
1:A:342:VAL:HG21	7:A:1265:HOH:O	2.00	0.61
1:A:6:HIS:HE1	1:A:337:GLN:HE21	1.49	0.60
1:A:381:LEU:CD1	1:A:442:VAL:HG11	2.31	0.60
1:A:230:ARG:NH1	7:A:1103:HOH:O	2.27	0.59
1:A:12:ARG:NE	7:A:1107:HOH:O	2.35	0.58
1:A:327:ASN:HD22	1:A:330:TRP:H	1.51	0.58
1:A:228:SER:HB2	1:A:266:MET:HB3	1.87	0.56
1:A:94:GLN:NE2	7:A:1108:HOH:O	2.38	0.56
1:A:68[B]:MET:SD	7:A:1344:HOH:O	2.56	0.56
1:A:12:ARG:NH2	7:A:1105:HOH:O	2.29	0.55
1:A:327:ASN:ND2	1:A:330:TRP:H	2.05	0.54
1:A:230:ARG:HD2	1:A:358:HIS:CE1	2.43	0.54
1:A:230:ARG:CZ	7:A:1103:HOH:O	2.53	0.54
1:A:171:PHE:H	1:A:209:ASN:ND2	2.06	0.54
1:A:292:THR:HG23	7:A:1478:HOH:O	2.08	0.53
1:A:119:GLN:NE2	7:A:1111:HOH:O	2.40	0.53
1:A:264:LEU:HG	1:A:281:CYS:HB2	1.90	0.53
1:A:346:GLU:HB2	1:A:352:CME:HZ2	1.91	0.53
1:A:342:VAL:HG13	1:A:352:CME:SD	2.50	0.52
1:A:63:MET:HE1	1:A:273:MET:HB2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:GLU:HG3	7:A:1304:HOH:O	2.09	0.52
1:A:230:ARG:CZ	1:A:270:LYS:HD3	2.39	0.52
1:A:133:ASN:HD22	1:A:133:ASN:N	2.05	0.52
1:A:301:GLN:CD	7:A:1205:HOH:O	2.54	0.51
1:A:401:LEU:CD2	1:A:413:CYS:HB2	2.41	0.51
1:A:115:LYS:O	1:A:119:GLN:HG3	2.09	0.51
1:A:292:THR:CG2	7:A:1478:HOH:O	2.58	0.51
1:A:419:ARG:NH2	4:A:1003:EPE:H101	2.23	0.50
1:A:198:ASN:OD1	1:A:358:HIS:HD2	1.96	0.49
1:A:407:THR:OG1	1:A:409[A]:LYS:HG3	2.14	0.48
1:A:68[B]:MET:HE3	7:A:1344:HOH:O	2.01	0.48
1:A:5:LYS:HB3	7:A:1384:HOH:O	2.13	0.48
1:A:342:VAL:CG2	7:A:1265:HOH:O	2.58	0.48
1:A:412:PRO:CA	7:A:1109:HOH:O	2.52	0.48
1:A:157:GLU:O	1:A:168:LYS:HE2	2.14	0.47
1:A:170:ASN:ND2	1:A:206:SER:H	2.10	0.47
1:A:5:LYS:HE2	7:A:1386:HOH:O	2.14	0.47
1:A:419:ARG:HH12	4:A:1003:EPE:C5	2.24	0.46
1:A:6:HIS:CE1	1:A:337:GLN:HE21	2.31	0.46
1:A:404:ASP:OD1	1:A:409[A]:LYS:NZ	2.49	0.45
4:A:1003:EPE:O8	4:A:1003:EPE:H51	2.16	0.45
4:A:1003:EPE:H61	4:A:1003:EPE:H101	1.57	0.44
1:A:138:THR:HG23	7:A:1413:HOH:O	2.16	0.44
1:A:271:ASP:O	1:A:273:MET:HG2	2.18	0.44
1:A:401:LEU:HD22	1:A:413:CYS:HB2	2.00	0.44
1:A:394:VAL:HG11	4:A:1003:EPE:H22	1.99	0.44
1:A:409[A]:LYS:HD3	7:A:1143:HOH:O	2.17	0.44
1:A:445:ASN:O	1:A:445:ASN:CG	2.60	0.43
4:A:1003:EPE:O3S	4:A:1003:EPE:H62	2.18	0.43
1:A:413:CYS:N	7:A:1109:HOH:O	2.39	0.43
1:A:236:TYR:CD2	1:A:335:ILE:HD13	2.54	0.42
1:A:123:LEU:C	1:A:123:LEU:HD23	2.45	0.42
1:A:402:GLY:HA2	1:A:462:ARG:CZ	2.49	0.42
1:A:459:LYS:HA	1:A:459:LYS:HD3	1.83	0.42
1:A:276[A]:MET:HB2	1:A:276[A]:MET:HE2	1.52	0.42
1:A:467:LYS:HE2	1:A:467:LYS:HB3	1.76	0.41
1:A:164:ARG:CG	1:A:164:ARG:NH1	2.80	0.41
1:A:142:HIS:HD2	7:A:1399:HOH:O	2.03	0.41
1:A:68[B]:MET:HE1	7:A:1452:HOH:O	2.20	0.40
1:A:100:HIS:HD2	1:A:309:GLY:HA2	1.80	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1500:HOH:O	7:A:1500:HOH:O[11_555]	1.66	0.54
7:A:1161:HOH:O	7:A:1161:HOH:O[14_555]	1.90	0.30
7:A:1472:HOH:O	7:A:1472:HOH:O[4_565]	1.93	0.27

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/467 (100%)	457 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	390/384 (102%)	372 (95%)	18 (5%)	24	16

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	HIS
1	A	107	GLN
1	A	118	GLU
1	A	133	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	145	ILE
1	A	164	ARG
1	A	207	LEU
1	A	218	LYS
1	A	276[A]	MET
1	A	276[B]	MET
1	A	327	ASN
1	A	353	GLN
1	A	366	LYS
1	A	406	LYS
1	A	409[A]	LYS
1	A	409[B]	LYS
1	A	409[C]	LYS
1	A	444	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	61	GLN
1	A	100	HIS
1	A	119	GLN
1	A	133	ASN
1	A	144	GLN
1	A	170	ASN
1	A	187	ASN
1	A	209	ASN
1	A	240	GLN
1	A	327	ASN
1	A	337	GLN
1	A	353	GLN
1	A	354	GLN
1	A	358	HIS
1	A	441	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	A	298	1	8,9,10	0.60	0	6,9,11	1.48	1 (16%)
1	CME	A	352	1	8,9,10	0.61	0	6,9,11	1.34	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	298	1	-	0/5/8/10	-
1	CME	A	352	1	-	2/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	352	CME	CA-CB-SG	-2.19	105.49	114.45
1	A	298	CME	CB-CA-C	-2.10	105.11	110.80

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	352	CME	CZ-CE-SD-SG
1	A	352	CME	CA-CB-SG-SD

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	352	CME	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 6 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	EPE	A	1003	-	15,15,15	1.24	1 (6%)	19,20,20	2.48	9 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	A	1003	-	-	8/9/19/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1003	EPE	C10-S	4.16	1.83	1.77

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	EPE	O1S-S-C10	5.07	114.38	106.73
4	A	1003	EPE	C7-N4-C3	4.23	122.50	111.24
4	A	1003	EPE	C5-N4-C3	4.19	117.86	108.84
4	A	1003	EPE	O3S-S-O1S	-3.62	102.34	111.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	EPE	C2-C3-N4	2.96	116.63	110.65
4	A	1003	EPE	C7-N4-C5	2.85	118.83	111.24
4	A	1003	EPE	C5-C6-N1	2.82	116.34	110.65
4	A	1003	EPE	O3S-S-C10	2.32	110.54	106.00
4	A	1003	EPE	O2S-S-C10	2.01	109.77	106.73

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1003	EPE	C10-C9-N1-C2
4	A	1003	EPE	C10-C9-N1-C6
4	A	1003	EPE	C9-C10-S-O1S
4	A	1003	EPE	C8-C7-N4-C3
4	A	1003	EPE	S-C10-C9-N1
4	A	1003	EPE	C9-C10-S-O3S
4	A	1003	EPE	C9-C10-S-O2S
4	A	1003	EPE	C8-C7-N4-C5

There are no ring outliers.

1 monomer is involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1003	EPE	11	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	463/467 (99%)	-0.10	14 (3%) 52 56	7, 22, 40, 57	6 (1%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	409[A]	LYS	8.0
1	A	308	GLY	3.8
1	A	307	TYR	3.3
1	A	163	VAL	2.8
1	A	162	GLY	2.6
1	A	304	PHE	2.6
1	A	305	PRO	2.5
1	A	104	GLY	2.3
1	A	353	GLN	2.3
1	A	441	HIS	2.3
1	A	302	GLU	2.2
1	A	408	GLY	2.2
1	A	105	ALA	2.2
1	A	164	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	352	10/11	0.91	0.15	31,34,48,50	0
1	CME	A	298	10/11	0.95	0.12	30,31,41,42	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EPE	A	1003	15/15	0.78	0.26	61,65,77,80	0
3	MG	A	1004	1/1	0.84	0.10	62,62,62,62	0
5	NA	A	1006	1/1	0.90	0.10	49,49,49,49	0
3	MG	A	1005	1/1	0.95	0.24	36,36,36,36	0
3	MG	A	1002	1/1	0.99	0.08	18,18,18,18	0
6	CA	A	1007	1/1	0.99	0.18	32,32,32,32	0
2	CL	A	1001	1/1	1.00	0.02	13,13,13,13	1

6.5 Other polymers [i](#)

There are no such residues in this entry.