



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 05:49 PM UTC

PDB ID : 6D8V / pdb_00006d8v
Title : Methyl-accepting Chemotaxis protein X
Authors : Shrestha, M.; Schubot, F.D.
Deposited on : 2018-04-27
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	8% 67% 27% 6%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2044 atoms, of which 12 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable chemoreceptor (Methyl-accepting chemotaxis) trans-membrane protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	269	2012	1269	344	393	6	0	0	0

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	8	4	1	3	0	0

- Molecule 3 is 1,1-DIMETHYL-PROLINIUM (CCD ID: PBE) (formula: $C_7H_{14}NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			22	7	12	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	O	0	0
			2	2		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable chemoreceptor (Methyl-accepting chemotaxis) transmembrane protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.04Å 119.04Å 121.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	78.63 – 2.80 78.63 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.4 (78.63-2.80) 92.8 (78.63-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.62Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.206 , 0.245 (Not available) , 0.241	Depositor DCC
R_{free} test set	671 reflections (4.15%)	wwPDB-VP
Wilson B-factor (Å ²)	74.0	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 87.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2044	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, PBE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	7/2055 (0.3%)	0.77	13/2790 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	227	GLY	CA-C	-6.99	1.45	1.51
1	A	227	GLY	C-O	-6.17	1.16	1.23
1	A	291	LEU	CA-C	-6.12	1.45	1.52
1	A	293	ASP	C-O	-5.62	1.16	1.24
1	A	223	PRO	N-CD	5.29	1.55	1.47
1	A	294	VAL	CA-CB	-5.22	1.47	1.54
1	A	294	VAL	C-O	-5.21	1.16	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	277	ASP	N-CA-C	9.96	124.68	110.23
1	A	276	TYR	N-CA-C	-9.06	92.71	108.20
1	A	292	PRO	N-CA-C	7.57	124.29	113.47
1	A	292	PRO	CB-CA-C	-7.27	101.50	113.06
1	A	225	GLY	N-CA-C	-6.51	97.76	113.18
1	A	291	LEU	N-CA-C	-6.28	102.23	110.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	292	PRO	CA-N-CD	-6.08	103.49	112.00
1	A	227	GLY	N-CA-C	-5.70	102.08	111.38
1	A	281	PRO	CA-N-CD	-5.59	104.17	112.00
1	A	222	LYS	N-CA-C	5.43	118.02	110.20
1	A	280	GLU	CA-C-N	-5.35	113.15	119.84
1	A	280	GLU	C-N-CA	-5.35	113.15	119.84
1	A	292	PRO	CA-C-O	-5.29	112.36	120.03

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	221	VAL	Peptide
1	A	276	TYR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2012	0	1952	72	0
2	A	8	0	11	1	0
3	A	10	12	13	1	0
4	A	2	0	0	0	0
All	All	2032	12	1976	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:LEU:HD12	1:A:282:PHE:HE2	1.23	1.03
1:A:222:LYS:HG2	1:A:223:PRO:HD2	1.45	0.98
1:A:222:LYS:HG2	1:A:223:PRO:CD	1.96	0.95
1:A:274:LEU:HD12	1:A:282:PHE:CE2	2.09	0.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:GLU:HG3	1:A:114:PRO:HD2	1.55	0.86
1:A:224:PHE:HB3	1:A:227:GLY:HA3	1.57	0.85
1:A:277:ASP:OD1	1:A:278:GLY:N	2.14	0.81
1:A:274:LEU:CD1	1:A:282:PHE:HE2	1.94	0.79
1:A:115:LYS:O	1:A:115:LYS:HD2	1.82	0.79
1:A:284:ARG:HD3	1:A:303:ASP:OD1	1.84	0.77
1:A:222:LYS:HG2	1:A:223:PRO:N	1.98	0.77
1:A:232:LEU:HB2	1:A:299:LEU:HB2	1.72	0.72
1:A:270:MET:HG3	1:A:285:VAL:HG22	1.72	0.71
1:A:285:VAL:HB	1:A:302:VAL:HG13	1.71	0.71
1:A:222:LYS:CG	1:A:223:PRO:HD2	2.19	0.70
1:A:55:ILE:CD1	1:A:221:VAL:HG23	2.24	0.67
1:A:43:LEU:HD12	1:A:43:LEU:O	1.95	0.66
1:A:187:THR:HB	1:A:211:LEU:HD22	1.78	0.64
1:A:282:PHE:HB3	1:A:305:PRO:HA	1.79	0.64
1:A:114:PRO:HD3	1:A:157:TYR:OH	1.98	0.62
1:A:247:LEU:O	1:A:248:MET:HB2	2.02	0.58
1:A:110:PHE:CE2	1:A:112:GLU:HB2	2.38	0.58
1:A:223:PRO:HB2	1:A:227:GLY:O	2.04	0.58
1:A:113:GLU:OE1	1:A:162:TYR:OH	2.20	0.57
1:A:114:PRO:O	1:A:115:LYS:HG3	2.03	0.57
1:A:88:ARG:NH1	1:A:128:ARG:O	2.38	0.56
1:A:256:VAL:H	2:A:401:TRS:HN2	1.53	0.56
1:A:282:PHE:HB2	1:A:304:VAL:O	2.05	0.56
1:A:136:PHE:CE1	1:A:138:PRO:HG3	2.40	0.56
1:A:140:TRP:HE3	1:A:148:ILE:HG22	1.71	0.56
1:A:282:PHE:HD2	1:A:282:PHE:C	2.14	0.54
1:A:282:PHE:C	1:A:282:PHE:CD2	2.85	0.54
1:A:114:PRO:HD3	1:A:157:TYR:CE2	2.44	0.53
1:A:225:GLY:O	1:A:226:SER:HB3	2.09	0.53
1:A:114:PRO:HD3	1:A:157:TYR:CZ	2.44	0.53
1:A:226:SER:O	1:A:226:SER:OG	2.25	0.52
1:A:92:ILE:HD11	1:A:136:PHE:HE1	1.75	0.52
1:A:88:ARG:O	1:A:92:ILE:HD12	2.10	0.51
1:A:149:GLN:NE2	1:A:183:ASP:HB2	2.25	0.51
1:A:271:ILE:HG22	1:A:274:LEU:HD23	1.93	0.50
1:A:55:ILE:CD1	1:A:221:VAL:CG2	2.89	0.50
1:A:234:GLN:HG2	1:A:296:ALA:HB1	1.94	0.49
1:A:222:LYS:HD3	1:A:223:PRO:O	2.12	0.49
1:A:218:LEU:HD13	1:A:229:VAL:HG23	1.95	0.49
1:A:123:ILE:O	1:A:123:ILE:HG13	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:GLU:HB2	1:A:274:LEU:HD22	1.95	0.48
1:A:282:PHE:CB	1:A:305:PRO:HA	2.44	0.48
1:A:255:GLY:HA3	1:A:271:ILE:HD13	1.96	0.47
1:A:122:VAL:HG22	1:A:122:VAL:O	2.13	0.47
1:A:88:ARG:HA	1:A:117:TYR:CE2	2.50	0.47
1:A:176:PRO:HB2	1:A:248:MET:HG2	1.98	0.46
1:A:127:GLU:OE1	1:A:127:GLU:N	2.49	0.46
1:A:281:PRO:O	1:A:306:ARG:HG2	2.15	0.46
1:A:230:TYR:HB3	1:A:240:ALA:O	2.16	0.45
1:A:274:LEU:CD1	1:A:282:PHE:CE2	2.85	0.45
1:A:263:LEU:HG	1:A:299:LEU:HD21	1.97	0.45
1:A:224:PHE:CD2	1:A:224:PHE:O	2.70	0.45
1:A:265:THR:HG21	1:A:269:ARG:HH12	1.82	0.45
1:A:289:PHE:CE1	1:A:300:VAL:HG23	2.52	0.44
1:A:237:LYS:HD2	1:A:248:MET:O	2.18	0.44
1:A:271:ILE:CG2	1:A:274:LEU:HD23	2.46	0.44
1:A:140:TRP:CE3	1:A:148:ILE:HG22	2.52	0.44
1:A:55:ILE:HD12	1:A:221:VAL:CG2	2.49	0.43
1:A:231:LEU:HD12	1:A:299:LEU:O	2.18	0.43
1:A:88:ARG:HG3	1:A:89:ALA:N	2.33	0.43
1:A:161:TRP:CD2	3:A:402:PBE:HG2	2.55	0.42
1:A:238:TRP:HB2	1:A:246:LEU:O	2.20	0.42
1:A:55:ILE:HD12	1:A:221:VAL:HG23	2.02	0.42
1:A:131:ASN:HB3	1:A:152:THR:HG22	2.01	0.41
1:A:149:GLN:HE22	1:A:183:ASP:HB2	1.86	0.41
1:A:255:GLY:HA3	1:A:271:ILE:CG2	2.51	0.41
1:A:73:MET:HB2	1:A:98:ASN:HD21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/269 (92%)	231 (94%)	16 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/206 (99%)	182 (89%)	22 (11%)	6	21

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	THR
1	A	76	VAL
1	A	84	GLN
1	A	86	THR
1	A	92	ILE
1	A	95	LEU
1	A	112	GLU
1	A	144	ARG
1	A	150	LEU
1	A	211	LEU
1	A	221	VAL
1	A	223	PRO
1	A	226	SER
1	A	229	VAL
1	A	234	GLN
1	A	252	ASP
1	A	274	LEU
1	A	275	THR
1	A	282	PHE
1	A	286	VAL
1	A	292	PRO
1	A	302	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TRS	A	401	-	7,7,7	0.30	0	9,9,9	0.33	0
3	PBE	A	402	-	9,10,10	3.61	5 (55%)	9,15,15	1.67	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	A	401	-	-	3/9/9/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PBE	A	402	-	-	2/4/17/17	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	PBE	CG-CB	-6.19	1.26	1.51
3	A	402	PBE	CA-C	-5.69	1.45	1.53
3	A	402	PBE	CD-N	-5.20	1.31	1.51
3	A	402	PBE	CD-CG	3.02	1.64	1.51
3	A	402	PBE	CB-CA	2.08	1.58	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	PBE	OXT-C-CA	4.06	121.26	111.88
3	A	402	PBE	OXT-C-O	-2.01	119.53	124.08

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	TRS	C2-C-C1-O1
2	A	401	TRS	C3-C-C1-O1
2	A	401	TRS	N-C-C1-O1
3	A	402	PBE	O-C-CA-CB
3	A	402	PBE	OXT-C-CA-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	TRS	1	0
3	A	402	PBE	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	269/269 (100%)	0.42	22 (8%)	17 13	52, 93, 142, 167	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	224	PHE	4.5
1	A	221	VAL	4.1
1	A	120	LYS	4.1
1	A	222	LYS	3.9
1	A	276	TYR	3.8
1	A	279	ASN	3.8
1	A	123	ILE	3.5
1	A	223	PRO	3.3
1	A	220	ALA	3.1
1	A	145	ASN	2.9
1	A	86	THR	2.9
1	A	278	GLY	2.8
1	A	219	SER	2.7
1	A	277	ASP	2.5
1	A	280	GLU	2.4
1	A	225	GLY	2.3
1	A	84	GLN	2.3
1	A	134	GLY	2.2
1	A	260	LYS	2.1
1	A	226	SER	2.0
1	A	122	VAL	2.0
1	A	183	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	TRS	A	401	8/8	0.85	0.17	112,131,136,138	0
3	PBE	A	402	10/10	0.96	0.10	91,110,132,134	0

6.5 Other polymers [i](#)

There are no such residues in this entry.