



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:56 PM UTC

PDB ID : 4D92 / pdb_00004d92
Title : Salmonella typhimurium D-Cysteine desulphydrase soaked with beta-chloro-D-alanine shows pyruvate bound 4 Å away from active site
Authors : Bharath, S.R.; Shveta, B.; Rajesh, K.H.; Savithri, H.S.; Murthy, M.R.N.
Deposited on : 2012-01-11
Resolution : 2.22 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	2% 91% 5% .
1	B	342	2% 89% 5% 6%
1	C	342	2% 86% 6% 7%
1	D	342	89% 7% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PYR	B	402	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-cysteine desulfhydrase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	P	S	0	3	0
			2465	1574	411	469	1	10			
1	B	321	Total	C	N	O	P	S	0	4	0
			2415	1538	405	462	1	9			
1	C	317	Total	C	N	O	P	S	0	3	0
			2367	1510	395	452	1	9			
1	D	328	Total	C	N	O	P	S	0	1	0
			2459	1568	415	465	1	10			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	expression tag	UNP Q8ZNT7
A	-12	ARG	-	expression tag	UNP Q8ZNT7
A	-11	GLY	-	expression tag	UNP Q8ZNT7
A	-10	SER	-	expression tag	UNP Q8ZNT7
A	-9	HIS	-	expression tag	UNP Q8ZNT7
A	-8	HIS	-	expression tag	UNP Q8ZNT7
A	-7	HIS	-	expression tag	UNP Q8ZNT7
A	-6	HIS	-	expression tag	UNP Q8ZNT7
A	-5	HIS	-	expression tag	UNP Q8ZNT7
A	-4	HIS	-	expression tag	UNP Q8ZNT7
A	-3	GLY	-	expression tag	UNP Q8ZNT7
A	-2	MET	-	expression tag	UNP Q8ZNT7
A	-1	ALA	-	expression tag	UNP Q8ZNT7
A	0	SER	-	expression tag	UNP Q8ZNT7
B	-13	MET	-	expression tag	UNP Q8ZNT7
B	-12	ARG	-	expression tag	UNP Q8ZNT7
B	-11	GLY	-	expression tag	UNP Q8ZNT7
B	-10	SER	-	expression tag	UNP Q8ZNT7
B	-9	HIS	-	expression tag	UNP Q8ZNT7
B	-8	HIS	-	expression tag	UNP Q8ZNT7
B	-7	HIS	-	expression tag	UNP Q8ZNT7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	HIS	-	expression tag	UNP Q8ZNT7
B	-5	HIS	-	expression tag	UNP Q8ZNT7
B	-4	HIS	-	expression tag	UNP Q8ZNT7
B	-3	GLY	-	expression tag	UNP Q8ZNT7
B	-2	MET	-	expression tag	UNP Q8ZNT7
B	-1	ALA	-	expression tag	UNP Q8ZNT7
B	0	SER	-	expression tag	UNP Q8ZNT7
C	-13	MET	-	expression tag	UNP Q8ZNT7
C	-12	ARG	-	expression tag	UNP Q8ZNT7
C	-11	GLY	-	expression tag	UNP Q8ZNT7
C	-10	SER	-	expression tag	UNP Q8ZNT7
C	-9	HIS	-	expression tag	UNP Q8ZNT7
C	-8	HIS	-	expression tag	UNP Q8ZNT7
C	-7	HIS	-	expression tag	UNP Q8ZNT7
C	-6	HIS	-	expression tag	UNP Q8ZNT7
C	-5	HIS	-	expression tag	UNP Q8ZNT7
C	-4	HIS	-	expression tag	UNP Q8ZNT7
C	-3	GLY	-	expression tag	UNP Q8ZNT7
C	-2	MET	-	expression tag	UNP Q8ZNT7
C	-1	ALA	-	expression tag	UNP Q8ZNT7
C	0	SER	-	expression tag	UNP Q8ZNT7
D	-13	MET	-	expression tag	UNP Q8ZNT7
D	-12	ARG	-	expression tag	UNP Q8ZNT7
D	-11	GLY	-	expression tag	UNP Q8ZNT7
D	-10	SER	-	expression tag	UNP Q8ZNT7
D	-9	HIS	-	expression tag	UNP Q8ZNT7
D	-8	HIS	-	expression tag	UNP Q8ZNT7
D	-7	HIS	-	expression tag	UNP Q8ZNT7
D	-6	HIS	-	expression tag	UNP Q8ZNT7
D	-5	HIS	-	expression tag	UNP Q8ZNT7
D	-4	HIS	-	expression tag	UNP Q8ZNT7
D	-3	GLY	-	expression tag	UNP Q8ZNT7
D	-2	MET	-	expression tag	UNP Q8ZNT7
D	-1	ALA	-	expression tag	UNP Q8ZNT7
D	0	SER	-	expression tag	UNP Q8ZNT7

- Molecule 2 is BENZAMIDINE (CCD ID: BEN) (formula: C₇H₈N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			9	7	2		
2	B	1	Total	C	N	0	0
			9	7	2		
2	D	1	Total	C	N	0	0
			9	7	2		

- Molecule 3 is PYRUVIC ACID (CCD ID: PYR) (formula: $C_3H_4O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	164	Total	O	0	0
			164	164		
5	B	145	Total	O	0	0
			145	145		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	106	Total 106	O 106	0	0
5	D	171	Total 171	O 171	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: D-cysteine desulfhyrase




- Molecule 1: D-cysteine desulfhyrase



- Molecule 1: D-cysteine desulfhyrase



- Molecule 1: D-cysteine desulfhyrase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.34Å 165.46Å 67.96Å 90.00° 118.50° 90.00°	Depositor
Resolution (Å)	58.30 – 2.22 58.30 – 2.22	Depositor EDS
% Data completeness (in resolution range)	92.4 (58.30-2.22) 92.4 (58.30-2.22)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.176 , 0.222 (Not available) , 0.199	Depositor DCC
R_{free} test set	3201 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.002 for -h-l,k,h 0.002 for l,k,-h-l 0.027 for h,-k,-h-l 0.035 for -h-l,-k,l 0.025 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10363	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, BEN, PYR, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2497	0.80	0/3404
1	B	0.60	0/2443	0.81	0/3329
1	C	0.58	0/2391	0.81	1/3259 (0.0%)
1	D	0.61	0/2481	0.81	1/3381 (0.0%)
All	All	0.60	0/9812	0.81	2/13373 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	170	VAL	N-CA-C	-5.32	105.35	110.72
1	C	45	ILE	N-CA-C	5.07	115.06	107.51

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2465	0	2461	10	0
1	B	2415	0	2403	11	0
1	C	2367	0	2350	14	0
1	D	2459	0	2463	14	0
2	A	9	0	7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	9	0	7	0	0
2	D	9	0	7	0	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	0	0
3	D	6	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
5	A	164	0	0	1	0
5	B	145	0	0	0	0
5	C	106	0	0	2	0
5	D	171	0	0	1	0
All	All	10363	0	9698	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1:MET:HB3	1:D:2:PRO:HD3	1.32	1.11
1:C:105:THR:OG1	1:C:328[A]:VAL:HG12	1.80	0.81
1:D:1:MET:HB3	1:D:2:PRO:CD	2.16	0.73
1:A:251:HIS:HE1	5:A:559:HOH:O	1.73	0.71
1:B:105:THR:OG1	1:B:328[A]:VAL:HG12	2.00	0.62
1:B:134:ASP:H	1:B:138:GLN:NE2	2.01	0.59
1:B:328[A]:VAL:HG11	1:D:145:ARG:HH12	1.68	0.59
1:D:51:LLP:H6	1:D:192:ALA:HB3	1.87	0.57
1:B:134:ASP:H	1:B:138:GLN:HE21	1.51	0.57
1:D:51:LLP:HE3	1:D:80:HIS:HB2	1.88	0.54
1:C:314:HIS:HD2	1:C:316:GLY:H	1.53	0.54
1:C:71:ILE:HG12	1:C:96:VAL:HB	1.89	0.54
1:B:314:HIS:HD2	1:B:316:GLY:H	1.56	0.53
1:D:94:HIS:HD2	5:D:572:HOH:O	1.91	0.53
1:D:105:THR:OG1	1:D:328[A]:VAL:HG12	2.09	0.53
1:C:257:PHE:HA	1:C:290:LYS:HD3	1.91	0.53
1:A:314:HIS:HD2	1:A:316:GLY:H	1.56	0.52
1:B:314:HIS:CD2	1:B:316:GLY:H	2.28	0.52
1:B:26:ARG:HD3	1:B:278:LEU:O	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:105:THR:HG1	1:C:328[A]:VAL:HG12	1.74	0.51
1:A:51:LLP:HE3	1:A:80:HIS:HB2	1.93	0.50
1:C:134:ASP:H	1:C:138:GLN:HE21	1.58	0.49
1:B:105:THR:CB	1:B:328[A]:VAL:HG12	2.43	0.49
1:A:51:LLP:H6	1:A:192:ALA:HB3	1.96	0.48
1:C:251:HIS:HE1	5:C:598:HOH:O	1.96	0.48
1:D:314:HIS:HD2	1:D:316:GLY:H	1.61	0.47
1:C:51:LLP:H6	1:C:192:ALA:HB3	1.96	0.46
1:C:314:HIS:CD2	1:C:316:GLY:H	2.32	0.46
1:C:112:THR:OG1	1:D:325:HIS:HE1	1.98	0.46
1:C:295:LEU:HD23	1:C:295:LEU:C	2.41	0.46
1:B:72:THR:OG1	1:B:80:HIS:HE1	1.99	0.46
1:A:14:PHE:HB3	1:A:45:ILE:HD12	1.98	0.46
1:C:134:ASP:H	1:C:138:GLN:NE2	2.15	0.44
1:D:166:ALA:CB	1:D:238:ILE:HD11	2.47	0.44
1:A:134:ASP:H	1:A:138:GLN:NE2	2.15	0.44
1:C:228:LYS:N	1:C:229:PRO:HD2	2.34	0.43
1:A:94:HIS:HD2	5:C:513:HOH:O	2.00	0.43
1:D:185:GLY:O	1:D:309:PRO:HD2	2.19	0.43
1:D:216:ILE:HD13	1:D:253:TRP:CZ2	2.54	0.42
1:B:50:ASN:HB2	1:B:169:TYR:CE1	2.55	0.41
1:B:262:GLY:HA2	1:B:286:VAL:HG13	2.03	0.41
1:A:43:THR:OG1	1:A:48:GLY:HA2	2.21	0.41
1:A:195:SER:O	1:A:196:ALA:HB3	2.20	0.41
1:C:42:VAL:O	1:C:42:VAL:HG12	2.20	0.41
1:D:52:LEU:HB2	1:D:83:GLN:HE21	1.85	0.41
1:A:272:VAL:HG22	1:A:283:LEU:HB2	2.03	0.40
1:D:53:ARG:HD2	1:D:169:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/266 (92%)	243 (100%)	1 (0%)	84	91
1	B	247/266 (93%)	244 (99%)	3 (1%)	63	76
1	C	240/266 (90%)	236 (98%)	4 (2%)	53	68
1	D	251/266 (94%)	250 (100%)	1 (0%)	84	91
All	All	982/1064 (92%)	973 (99%)	9 (1%)	70	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	LEU
1	B	6	LEU
1	B	7	THR
1	B	146	ILE
1	C	99	LEU
1	C	273	LYS
1	C	283	LEU
1	C	284	ASP
1	D	174	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	HIS
1	A	83	GLN
1	A	94	HIS
1	A	138	GLN
1	A	149	GLN
1	A	178	GLN
1	A	208	HIS
1	A	241	GLN
1	A	251	HIS
1	A	314	HIS
1	B	80	HIS
1	B	138	GLN
1	B	208	HIS
1	B	314	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	325	HIS
1	C	80	HIS
1	C	138	GLN
1	C	178	GLN
1	C	241	GLN
1	C	251	HIS
1	C	301	GLN
1	C	314	HIS
1	D	4	HIS
1	D	80	HIS
1	D	83	GLN
1	D	94	HIS
1	D	125	GLN
1	D	138	GLN
1	D	149	GLN
1	D	314	HIS
1	D	325	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	A	51	1	23,24,25	0.74	0	25,32,34	1.03	2 (8%)
1	LLP	C	51	1	23,24,25	0.69	0	25,32,34	1.08	1 (4%)
1	LLP	D	51	1	23,24,25	0.70	0	25,32,34	1.12	1 (4%)
1	LLP	B	51	1	23,24,25	0.79	1 (4%)	25,32,34	1.34	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	51	1	-	4/16/17/19	0/1/1/1
1	LLP	C	51	1	-	3/16/17/19	0/1/1/1
1	LLP	D	51	1	-	3/16/17/19	0/1/1/1
1	LLP	B	51	1	-	4/16/17/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	51	LLP	C3-C2	-2.31	1.38	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	51	LLP	OP4-C5'-C5	2.57	114.17	109.36
1	B	51	LLP	OP4-C5'-C5	2.37	113.81	109.36
1	B	51	LLP	CD-CE-NZ	-2.26	104.85	110.83
1	B	51	LLP	C5-C6-N1	-2.17	120.30	123.83
1	C	51	LLP	C5-C6-N1	-2.10	120.41	123.83
1	A	51	LLP	OP3-P-OP2	2.04	115.46	107.80
1	A	51	LLP	C5-C6-N1	-2.03	120.52	123.83

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	51	LLP	O-C-CA-CB
1	C	51	LLP	O-C-CA-CB
1	A	51	LLP	C4-C4'-NZ-CE
1	D	51	LLP	C4-C4'-NZ-CE
1	C	51	LLP	C4-C4'-NZ-CE
1	B	51	LLP	C3-C4-C4'-NZ
1	B	51	LLP	CD-CE-NZ-C4'
1	B	51	LLP	C5-C4-C4'-NZ
1	B	51	LLP	CG-CD-CE-NZ
1	A	51	LLP	C3-C4-C4'-NZ
1	C	51	LLP	C3-C4-C4'-NZ
1	D	51	LLP	C3-C4-C4'-NZ

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	51	LLP	CE-CD-CG-CB
1	D	51	LLP	CE-CD-CG-CB

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	51	LLP	2	0
1	C	51	LLP	1	0
1	D	51	LLP	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BEN	A	401	-	9,9,9	0.88	0	7,11,11	0.77	0
4	SO4	C	402	-	4,4,4	0.43	0	6,6,6	0.21	0
4	SO4	A	403	-	4,4,4	0.43	0	6,6,6	0.28	0
2	BEN	B	401	-	9,9,9	1.30	1 (11%)	7,11,11	0.81	0
3	PYR	D	402	-	5,5,5	1.52	1 (20%)	3,6,6	1.43	0
2	BEN	D	401	-	9,9,9	1.38	2 (22%)	7,11,11	0.90	1 (14%)
4	SO4	B	403	-	4,4,4	0.36	0	6,6,6	0.19	0
4	SO4	D	403	-	4,4,4	0.42	0	6,6,6	0.45	0
3	PYR	C	401	-	5,5,5	1.45	1 (20%)	3,6,6	1.57	1 (33%)
3	PYR	A	402	-	5,5,5	1.62	1 (20%)	3,6,6	0.78	0
3	PYR	B	402	-	5,5,5	1.47	1 (20%)	3,6,6	1.73	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEN	A	401	-	-	0/4/4/4	0/1/1/1
2	BEN	B	401	-	-	0/4/4/4	0/1/1/1
3	PYR	D	402	-	-	4/4/4/4	-
2	BEN	D	401	-	-	0/4/4/4	0/1/1/1
3	PYR	C	401	-	-	3/4/4/4	-
3	PYR	A	402	-	-	2/4/4/4	-
3	PYR	B	402	-	-	4/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	BEN	C-N2	-2.87	1.26	1.33
2	D	401	BEN	C-N2	-2.69	1.27	1.33
3	D	402	PYR	CA-C	-2.57	1.45	1.54
3	B	402	PYR	CA-C	-2.50	1.46	1.54
3	A	402	PYR	CA-C	-2.44	1.46	1.54
3	C	401	PYR	CA-C	-2.30	1.46	1.54
2	D	401	BEN	C1-C	2.11	1.51	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	PYR	OXT-C-CA	2.68	121.01	113.59
3	C	401	PYR	OXT-C-CA	2.26	119.86	113.59
2	D	401	BEN	C1-C-N2	2.19	121.34	118.01

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	PYR	OXT-C-CA-CB
3	B	402	PYR	O-C-CA-CB
3	B	402	PYR	OXT-C-CA-CB
3	C	401	PYR	O-C-CA-CB
3	C	401	PYR	OXT-C-CA-CB
3	D	402	PYR	O-C-CA-O3
3	D	402	PYR	O-C-CA-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	402	PYR	OXT-C-CA-O3
3	D	402	PYR	OXT-C-CA-CB
3	A	402	PYR	O-C-CA-CB
3	B	402	PYR	OXT-C-CA-O3
3	C	401	PYR	OXT-C-CA-O3
3	B	402	PYR	O-C-CA-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/342 (95%)	-0.48	2 (0%) 85 84	11, 17, 30, 46	3 (0%)
1	B	320/342 (93%)	-0.18	5 (1%) 70 68	9, 22, 38, 53	4 (1%)
1	C	316/342 (92%)	0.04	8 (2%) 58 56	9, 25, 42, 56	4 (1%)
1	D	327/342 (95%)	-0.43	1 (0%) 90 89	10, 17, 32, 64	1 (0%)
All	All	1290/1368 (94%)	-0.27	16 (1%) 76 75	9, 20, 39, 64	12 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1	MET	3.4
1	C	7	THR	3.0
1	C	144	THR	3.0
1	C	142	LEU	3.0
1	C	145	ARG	2.7
1	B	4	HIS	2.7
1	C	240	GLY	2.7
1	B	328[A]	VAL	2.6
1	C	143	ALA	2.6
1	C	306	ASP	2.6
1	A	1	MET	2.6
1	B	145	ARG	2.3
1	C	328[A]	VAL	2.3
1	B	181	GLU	2.2
1	B	148	ALA	2.1
1	A	2	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	B	51	24/25	0.95	0.08	17,18,21,21	0
1	LLP	C	51	24/25	0.96	0.07	18,20,21,23	0
1	LLP	D	51	24/25	0.97	0.06	14,15,15,15	0
1	LLP	A	51	24/25	0.98	0.05	13,14,15,16	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PYR	C	401	6/6	0.68	0.20	41,41,42,42	6
3	PYR	B	402	6/6	0.75	0.18	32,33,33,33	6
3	PYR	A	402	6/6	0.75	0.16	31,31,32,33	0
3	PYR	D	402	6/6	0.90	0.11	34,35,36,37	0
4	SO4	C	402	5/5	0.94	0.09	23,23,24,25	5
4	SO4	B	403	5/5	0.95	0.14	16,16,16,16	5
2	BEN	B	401	9/9	0.95	0.07	27,27,28,28	0
4	SO4	D	403	5/5	0.95	0.09	17,17,18,18	5
2	BEN	D	401	9/9	0.96	0.06	13,13,14,14	0
2	BEN	A	401	9/9	0.96	0.06	13,13,13,13	0
4	SO4	A	403	5/5	0.98	0.07	15,15,15,15	5

6.5 Other polymers [i](#)

There are no such residues in this entry.