



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 11:19 AM UTC

PDB ID : 5D96 / pdb_00005d96
Title : Oxidoreductase Fragment of Mouse QSOX1 in Complex with a FAb Fragment from an Antibody Targeting Mouse and Human QSOX1
Authors : Fass, D.; Grossman, I.
Deposited on : 2015-08-18
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	78% 19% .
1	D	244	77% 18% ..
2	B	214	82% 18%
2	I	214	83% 16%
3	C	220	77% 20% .
3	J	220	80% 18% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10605 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfhydryl oxidase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	234	Total	C	N	O	S	0	0	0
			1807	1152	307	341	7			
1	A	237	Total	C	N	O	S	0	1	0
			1846	1177	314	348	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	32	GLY	-	expression tag	UNP Q8BND5
D	33	SER	-	expression tag	UNP Q8BND5
D	34	HIS	-	expression tag	UNP Q8BND5
D	35	MET	-	expression tag	UNP Q8BND5
A	32	GLY	-	expression tag	UNP Q8BND5
A	33	SER	-	expression tag	UNP Q8BND5
A	34	HIS	-	expression tag	UNP Q8BND5
A	35	MET	-	expression tag	UNP Q8BND5

- Molecule 2 is a protein called Light chain of Fab fragment from an antibody targeting mouse and human QSOX1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	214	Total	C	N	O	S	0	1	0
			1656	1030	280	337	9			
2	B	214	Total	C	N	O	S	0	0	0
			1651	1027	279	336	9			

- Molecule 3 is a protein called Heavy chain of Fab fragment from an antibody targeting mouse and human QSOX1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	J	215	Total	C	N	O	S	0	0	0
			1612	1021	267	315	9			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	214	Total	C	N	O	S	0	0	0
			1608	1019	266	314	9			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	75	Total	O	0	0
			75	75		
4	I	77	Total	O	0	0
			77	77		
4	J	71	Total	O	0	0
			71	71		
4	A	81	Total	O	0	0
			81	81		
4	B	66	Total	O	0	0
			66	66		
4	C	55	Total	O	0	0
			55	55		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sulfhydryl oxidase 1




- Molecule 1: Sulfhydryl oxidase 1




- Molecule 2: Light chain of Fab fragment from an antibody targeting mouse and human QSOX1



- Molecule 2: Light chain of Fab fragment from an antibody targeting mouse and human QSOX1






- Molecule 3: Heavy chain of Fab fragment from an antibody targeting mouse and human QSOX1

Chain J: 80% 18% .



- Molecule 3: Heavy chain of Fab fragment from an antibody targeting mouse and human QSOX1

Chain C: 77% 20% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.80Å 204.77Å 44.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.42 – 2.30 27.42 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (27.42-2.30) 96.9 (27.42-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.31Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.167 , 0.228 (Not available) , 0.211	Depositor DCC
R_{free} test set	1998 reflections (3.20%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 21.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.420 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10605	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/1896	0.79	3/2589 (0.1%)
1	D	0.34	0/1850	0.86	3/2522 (0.1%)
2	B	0.31	0/1686	0.78	1/2287 (0.0%)
2	I	0.34	0/1694	0.79	2/2298 (0.1%)
3	C	0.32	0/1648	0.81	2/2250 (0.1%)
3	J	0.31	0/1652	0.82	2/2255 (0.1%)
All	All	0.33	0/10426	0.81	13/14201 (0.1%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	204	HIS	CA-C-N	5.91	125.78	119.28
3	C	204	HIS	C-N-CA	5.91	125.78	119.28
1	D	165	PRO	CA-C-N	5.79	126.18	119.47
1	D	165	PRO	C-N-CA	5.79	126.18	119.47
2	B	30	SER	CB-CA-C	-5.51	110.24	116.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1846	0	1790	33	1
1	D	1807	0	1758	42	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1651	0	1585	28	0
2	I	1656	0	1591	24	1
3	C	1608	0	1592	25	0
3	J	1612	0	1595	27	0
4	A	81	0	0	4	0
4	B	66	0	0	1	0
4	C	55	0	0	2	1
4	D	75	0	0	4	0
4	I	77	0	0	1	0
4	J	71	0	0	1	1
All	All	10605	0	9911	167	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:226:SER:OG	2:B:108:ARG:NH1	2.05	0.89
2:B:149:LYS:HB2	2:B:193:THR:HB	1.55	0.88
1:A:204:ARG:HG3	1:A:220:ARG:HD2	1.53	0.88
1:D:230:ASN:ND2	2:B:169:LYS:O	2.14	0.81
1:D:128:GLN:HE21	1:D:128:GLN:HA	1.48	0.77

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:J:301:HOH:O	4:C:333:HOH:O[2_554]	2.09	0.11
2:I:17:ASP:OD1	1:A:197:ARG:NH1[1_556]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	74/244 (30%)	72 (97%)	2 (3%)	0	100	100
1	D	230/244 (94%)	226 (98%)	4 (2%)	0	100	100
2	B	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
2	I	213/214 (100%)	207 (97%)	6 (3%)	0	100	100
3	C	210/220 (96%)	206 (98%)	4 (2%)	0	100	100
3	J	209/220 (95%)	207 (99%)	2 (1%)	0	100	100
All	All	1148/1356 (85%)	1125 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/244 (97%)	-1.38	0 100 100	20, 35, 51, 70	1 (0%)
1	D	234/244 (95%)	-1.39	0 100 100	25, 34, 51, 69	1 (0%)
2	B	214/214 (100%)	-1.44	0 100 100	22, 34, 49, 57	0
2	I	214/214 (100%)	-1.46	0 100 100	23, 32, 43, 53	1 (0%)
3	C	214/220 (97%)	-1.42	0 100 100	23, 35, 52, 65	0
3	J	215/220 (97%)	-1.42	0 100 100	22, 33, 45, 54	0
All	All	1328/1356 (97%)	-1.42	0 100 100	20, 34, 49, 70	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.